

BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds.

Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A.



BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds



Grey Wagtail is one of the decreasing species highlighted by *BirdTrends 2013* - WBS/WBBS volunteers have recorded a 58% decline since 1975, moving it to the high-alert category

Key findings

Species list

Using the BirdTrends pages

These pages are a one-stop shop for information about the population status of the common breeding birds of the wider UK countryside. The report is based on data gathered by the many thousands of volunteers who contribute to BTO-led surveys. For each of 117 species, users can quickly access the latest information on trends in population size, breeding performance and survival rates, as measured by our long-term monitoring schemes.

The website covers the majority of UK breeding birds but excludes (with a few exceptions) colonial seabirds, which are well covered by the JNCC's [Seabird Monitoring Programme](#), and the rare species that are included in the reports of the [Rare Breeding Birds Panel](#) (e.g. Holling & RBBP 2012). For each species, we provide:

- General information concerning species' conservation listings and UK population sizes
- A summary of observed changes in the size of the population and information concerning the possible causes of these changes
- A series of graphs and tables showing the trends and changes in population size, breeding performance and survival over the longest periods available
- Trends for the BTO/JNCC/RSPB Breeding Bird Survey (BBS) cover not just UK as a whole but also each of its constituent countries (England, Scotland, Wales and Northern Ireland)
- Alerts, drawing attention to population declines in any census scheme of greater than 25%, or greater than 50%, that have occurred over the past five, 10 and 25 years and the maximum period available (usually 44 years).

Text, tables, graphs and presentation are updated annually to include the latest results and interpretative material from the literature. Information on demographic trends and on the causes of change is gradually being expanded. The Key findings page provides a brief overview of our main findings this year.

Supporting pages describe the field and analytical methods that were used to produce the results for each species and to identify alerts. We discuss overall patterns of trends in abundance and breeding success, and compare the latest trend information and alerts with the Birds of Conservation Concern list (Eaton *et al.* 2009). Summary tables list alerts and population changes by scheme, and there is also a facility to select and display [your own tables](#) of population change. A detailed References section lists more than 640 of the most relevant recent publications, with onward links to abstracts or to full text where freely available, and is a valuable key to recent scientific work by BTO and other researchers.

We would value your comments on this report and particularly any suggestions on how it can be improved.

EMAIL YOUR
COMMENTS

Authors

These web pages constitute an annual report that is part of the BTO research report series. Authors were Stephen Baillie, John Marchant, David Leech, Dario Massimino, Sarah Eglington, Alison Johnston, David Noble, Carl Barimore, Allison Kew, Iain Downie, Kate Risely and Rob Robinson. The formal citation for the report is given in the page footer.

Key findings

This section summarises the key findings of the report, under six headings, based on the results presented and discussed in the Summary tables and Discussion sections. It concentrates on the alerts raised by this edition of the report and changes to alerts since previous reports in this series.

Declining species



Losses of Wood Warblers in the UK amount to 69% of the population since 1995.

In the current report, there are 26 species for which our best long-term trends show statistically significant population declines of greater than 50% over periods of 31–44 years.

These are [Grey Partridge](#), [Lapwing](#), [Redshank](#), [Woodcock](#), [Snipe](#), [Turtle Dove](#), [Cuckoo](#), [Little Owl](#), [Willow Tit](#), [Marsh Tit](#), [Skylark](#), [Willow Warbler](#), [Whitethroat](#), [Starling](#), [Song Thrush](#), [Mistle Thrush](#), [Spotted Flycatcher](#), [House Sparrow](#), [Tree Sparrow](#), [Yellow Wagtail](#), [Grey Wagtail](#), [Tree Pipit](#), [Linnet](#), [Lesser Redpoll](#), [Yellowhammer](#) and [Corn Bunting](#) (taxonomic order). Lapwing and Grey Wagtail are additions since *BirdTrends 2012*.

Two further species show non-significant declines greater than 50% over long timescales. Change for [Lesser Spotted Woodpecker](#) is non-significant only because its monitoring period ceased in 1999: further strong decline has since been logged by Atlas data. The [House Martin](#) trend has a broad confidence interval that marginally includes zero change.

The steepest long-term populations declines we have measured are for [Tree Sparrow](#), [Turtle Dove](#), [Snipe](#), [Willow Tit](#) and [Grey Partridge](#), which have all declined by 90% or more since 1967, as almost certainly has Lesser Spotted Woodpecker. Tree Sparrow shows the biggest long-term decline despite its strong increase since 1994. Turtle Dove shows the greatest percentage losses of any species over the recent ten-year period.

These 26 species that have halved outweigh the 20 species found to show an equivalent increase, i.e. a doubling of population size, over similar periods (see Positive changes). The gap between these two groups has widened since *BirdTrends 2012*.

Except for [Little Owl](#), which as an introduced species is not eligible, all these rapidly declining species already benefit from listing as either red or amber Birds of Conservation Concern (PSoB/BoCC3) list, as revised in 2009. Seven species among those that have declined the most are listed only as amber and may be candidates for red listing at the next review: [Redshank](#), [Woodcock](#), [Snipe](#), [Willow Warbler](#), [Whitethroat](#), [Mistle Thrush](#) and [Grey Wagtail](#).

A further seven species raise lower-level concern, as a result of statistically significant long-term declines of between 25% and 50%. These are [Common Sandpiper](#), [Tawny Owl](#), [Dipper](#), [Dunnock](#), [Meadow Pipit](#), [Bullfinch](#) and [Reed Bunting](#). All of these species are already on the amber list on account of their population declines, except for [Dipper](#) and [Tawny Owl](#), which remain for now on the green list.

In addition, [Little Grebe](#) and [Curlew](#) (both amber-listed) have also declined by more than 25%, but raise no formal long-term alerts because the confidence intervals around their change estimates are too wide.

Three scarcer species with much shorter monitoring histories have also decreased by more than half during just a 16-year period: [Wood Warbler](#) (red listed), [Whinchat](#) and [Pied Flycatcher](#). The last two are only amber listed but are declining so steeply that red listing may well be warranted.

Recent changes to alerts



Reed Bunting has declined by 27% across the UK since 1967, and rejoins the species for which alerts for population decline are flagged

The *BirdTrends* report raises species alerts for population change to conservation bodies when the best available estimates of long-term decline are statistically significant and pass criteria set at -25% and -50%. Amber and red listings use similar criteria but are reviewed only at intervals of around seven years: thus this report's annual alerts provide valuable updates of current conservation concern.

Species with declines close to these threshold values often change category between years. The latest data indicates, however, that just three adjustments are needed to the long-term alerts since *BirdTrends 2012*. No species leaves the alert category as a whole, nor moves to the lower level, but one species re-enters and two move to the higher alert level.

[Lapwing](#) dropped to the lower alert level in *BirdTrends 2012*, when its long-term UK population decline was estimated at -45%. With a further steep phase of decline currently under way, it now re-enters the higher alert category. Its decline now measures -54% since 1967.

[Grey Wagtail](#) was a green-listed species until amber listed in 2002 but has since undergone accelerating decline. It now moves into the higher alert category for the first time, with a decline of -58%, as measured by WBS/WBBS since 1975.

[Reed Bunting](#) is a species at the borderline of the alert category. After severe declines during the late 1970s it appears to be fluctuating around a much lower population level than when our monitoring began in 1967. In this report it once again raises an alert at the lower level.

Positive changes



Although Red Grouse is amber listed for population decline, its current ten-year trend of -8% represents a positive shift in its rate of decline

Although much of this report focuses on declines and their conservation significance, there are many species that are increasingly strongly as UK breeding birds.

In the current report, there are 20 species for which our most representative long-term trends show a statistically significant doubling in population size over periods of 30–44 years.

These are [Mute Swan](#), [Canada Goose](#), [Shelduck](#), [Mallard](#), [Sparrowhawk](#), [Buzzard](#), [Coot](#), [Stock Dove](#), [Woodpigeon](#), [Collared Dove](#), [Green Woodpecker](#), [Great Spotted Woodpecker](#), [Magpie](#), [Jackdaw](#), [Carrion Crow](#), [Great Tit](#), [Long-tailed Tit](#), [Blackcap](#), [Reed Warbler](#) and [Nuthatch](#) (taxonomic order).

The steepest long-term increases we have measured have been for [Buzzard](#) and [Great Spotted Woodpecker](#), which have both increased by more than 400% since 1967.

These 20 species are directly equivalent to the 26 that have halved in number over similar periods (see Declining species). The gap between these two totals has widened since *BirdTrends 2012*.

Four further species, monitored only over shorter periods have also more than doubled (while three equivalent species have more than halved) [Greylag Goose](#) numbers have risen by 486% since 1993 and during the period since 1995 increases above 100% have been recorded for [Barn Owl](#) (+279%), [Red Kite](#) (+676%) and the non-native [Ring-necked Parakeet](#) (+1057%). [Ring-necked Parakeet](#) has maintained exponential growth since 1995 and has by far the highest rate of increase of any species in this report.

For ten species that are listed red or amber for population decline over the long term – [Red Grouse](#), [Little Grebe](#), [Skylark](#), [Willow Warbler](#), [Song Thrush](#), [House Sparrow](#), [Linnet](#), [Lesser Redpoll](#), [Yellowhammer](#) and [Corn Bunting](#) – decline has started to level off, or has ceased, during the recent ten-year period. Five species have joined this category since *BirdTrends 2012*, as their ten-year trends have risen marginally above -11%.

Six further formerly declining species – [Whitethroat](#), [Grasshopper Warbler](#), [Duncock](#), [Tree Sparrow](#), [Bullfinch](#) and [Reed Bunting](#) – have reversed their population trend to show significant increases over the last ten years. For most of these species, however, population levels remain severely depleted despite the recent increases.

Reduced breeding success



A reduction in breeding success has been the main driver of Linnet population declines.

Our best overall measure of breeding success is Fledglings Per Breeding Attempt (FPBA), calculated from brood sizes and nest failure rates, which indicates the mean number of young leaving each nest in a given year.

Nine species exhibit reduced FPBA over the past 43 years, indicating that their productivity has decreased over time: three red-listed species ([Nightjar](#), [Tree Pipit](#) and [Linnet](#)), two amber-listed species ([Willow Warbler](#) and [Bullfinch](#)) and three green-listed species ([Moorhen](#), [Coal Tit](#), [Chaffinch](#) and [Greenfinch](#)). While productivity of [Moorhen](#), [Nightjar](#), [Willow Warbler](#) and [Linnet](#) has fallen consistently, trends for the other five species are curvilinear, increasing between the mid 1960s and mid 1980s and decreasing thereafter.

Productivity declines in the migrant [Nightjar](#), [Tree Pipit](#) and [Willow Warbler](#) may be driven by changes in habitat or climate on the African wintering grounds or by declining insect numbers in the UK. Alternatively, climatic warming may have resulted in a developing asynchrony between laying dates and the availability of insect prey on the breeding grounds. Long-distance migrants are thought to be particularly susceptible to such disjunction but residents may also be affected, particularly those reliant on seasonal peaks in caterpillars, such as [Chaffinch](#) and to a lesser extent, [Coal Tit](#). However, for species such as these and [Greenfinch](#), where numbers have increased over the same period, we cannot exclude the possibility that reduced breeding success is due to a density dependent increase in intraspecific competition. Lack of food for nestling and breeding female Linnet due to a paucity of stubbles and weeds in more intensively farmed agricultural habitats may have contributed to the reduction in the species' breeding success, while loss of cover and food due to increased grazing pressure from deer may be implicated in Bullfinch productivity declines. The driver for increased [Moorhen](#) nest failure is at present unclear, but increases in aquatic mammalian predators and [Coot](#) populations have been proposed

CES ringing data integrate productivity across the whole season, including juvenile survival in the first few weeks or months after fledging. According to this measure, productivity has fallen significantly for 10 of the species monitored. The apparent breeding success of [Blue Tit](#), [Garden Warbler](#), [Sedge Warbler](#), [Goldfinch](#) and [Reed Bunting](#) has fallen by more than 50% over the last 25 years, while [Great Tit](#), [Willow Warbler](#), [Blackcap](#), [Blackbird](#) and [Song Thrush](#) show reductions in relative productivity of between 25% and 50%. For species such as [Blackcap](#), [Goldfinch](#), [Blue Tit](#) and [Great Tit](#) where population increase has occurred, reductions in productivity may be driven by density-dependent processes, with increased competition for resources in an expanding population reducing the mean breeding success per pair. [Song Thrush](#), [Blackbird](#) and [Sedge Warbler](#), have experienced significant declines in abundance, either on CES sites or more widely (based on CBC/BBS figures), but previous analyses suggest that falling survival rates are likely to have been a more important contributor to population changes than reduced productivity. There is some evidence that a reduction in the number of offspring produced may be the driver of [Willow Warbler](#) declines, however, and may also be preventing recovery of the UK [Reed Bunting](#) population.

Increased breeding success



Many trans-Saharan migrants are exhibiting population declines but Reed Warbler is bucking the trend, possibly due to a substantial increase in breeding success.

Our best overall measure of breeding success is Fledglings Per Breeding Attempt (FPBA), calculated from brood sizes and nest failure rates, which indicates the mean number of young leaving each nest in a given year.

FPBA has increased significantly over the last 43 years for 31 species, across a wide range of taxonomic groups. Population trends are also upward for 16 of these species, including raptors ([Sparrowhawk](#), [Buzzard](#), [Merlin](#), [Peregrine](#), [Barn Owl](#)), pigeons ([Stock Dove](#), [Woodpigeon](#), [Collared Dove](#)), corvids ([Maggie](#), [Jackdaw](#), [Carrion Crow](#)), and some small passerines ([Reed Warbler](#), [Nuthatch](#), [Wren](#), [Robin](#) and [Pied Wagtail](#)). It is therefore possible that increasing productivity has contributed to the population growth exhibited by these species over recent decades.

Conversely, 13 species ([Kestrel](#), [Tawny Owl](#), [Skylark](#), [Starling](#), [Dipper](#), [Blackbird](#), [Wheatear](#), [Duncock](#), [House Sparrow](#), [Tree Sparrow](#), [Grey Wagtail](#), [Meadow Pipit](#) and [Yellowhammer](#)), have declined in number as FPBA has increased, suggesting that a density-dependent reduction in intraspecific competition may have enabled breeding success to rise.

CES ringing data integrate productivity across the whole season, including juvenile survival in the first few weeks or months after fledging. According to this measure, productivity has risen significantly for two of the species monitored, [Reed Warbler](#) and [Chaffinch](#).

Early breeding



Short-distance migrants have exhibited some of the biggest advances in laying dates; Blackcap (pictured) and Chiffchaff are breeding a fortnight earlier than they did 40 years ago

Data from the Nest Record Scheme provide strong evidence of shifts towards earlier laying in a range of species, linked to climate change. We have now identified 42 species that, on average, are laying between one and 31 days earlier, on average, than in the mid 1960s.

The species now laying earlier in the year represent a wide range of taxonomic and ecological groups, including raptors ([Kestrel](#) – 9 days), waterbirds ([Moorhen](#) – 6 days), waders ([Oystercatcher](#) – 5 days), migrant insectivores ([Pied Flycatcher](#) – 12 days, [Swallow](#) – 8 days), resident insectivores ([Robin](#) – 8 days, [Blue Tit](#) – 9 days), corvids ([Magpie](#) – 31 days) and resident seed-eaters ([Greenfinch](#) – 17 days).

For some species these shifts towards earlier laying may be insufficient to match seasonal advances in the peaks of food availability. Recent research has shown that significantly stronger phenological responses to climate change are displayed at lower trophic levels (such as the food birds eat) than at higher levels (such as the birds themselves), increasing the potential for disjunction and resulting productivity declines. However, the evidence for a population-level effect of reduction in breeding success is mixed and more research is needed to determine the extent to which declines in abundance will result.

Only five species demonstrate a significant delay in average laying dates: [Woodpigeon](#), [Skylark](#), [Blackbird](#), [Bullfinch](#) and [Yellowhammer](#). All of these species initiate multiple breeding attempts per season and there is increasing evidence that birds less reliant on seasonal peaks in resource availability may be able to extend their breeding seasons further into the summer, resulting in a later mean value for laying date.

Introduction

Since its formation in 1933, BTO has been deeply committed to gathering quantitative information on the bird populations of the UK. Its nationwide network of skilled volunteers, many of whom are long-term contributors to survey schemes, provides the ideal way to monitor the bird populations that are widely distributed across the countryside. BTO data, from such schemes as the Common Birds Census, Nest Record Scheme and BTO/JNCC/RSPB Breeding Bird Survey, have been increasingly influential in determining nature conservation policy in the UK. The partnership between JNCC and BTO has ensured that these schemes are operated and developed so as to provide high-quality information for nature conservation.

The value of the monitoring work undertaken by the BTO was recognised in the Government's Biodiversity Steering Group report (Anon. 1995). The BTO's results, particularly those regarding declining farmland species, are highlighted as an example of the way in which broad-scale surveillance techniques can identify significant new trends. More generally, the report states that monitoring is essential if the broad aims, specific objectives and precise targets of the Government's Biodiversity Action Plans are to be achieved. It notes that:

- baselines must be established;
- regular and systematic recording must be made, to detect change; and
- the reasons for change should be studied, to inform action.

The BTO's monitoring schemes fulfil a considerable portion of these needs for a wide range of bird species in the UK.

The system of alerts, derived from the BTO's census and nest record data, ensures that conservation bodies are quickly made aware of important demographic changes. Multi-species *indicators*, making extensive use of BTO census data, track how bird populations are faring generally across the countryside, UK-wide and within specific regions or habitats. These indicators were developed in association with Government and some have been adopted by them as policy drivers. More recently, [European bird indicators](#) have been developed (PECBMS 2013b).

Monitoring UK breeding birds

Long-running bird surveys operated by BTO contribute to an overall programme of Integrated Population Monitoring (IPM) that has been developed by the BTO, in partnership with JNCC, to monitor the numbers, breeding performance and survival rates of a wide range of bird species. IPM has the following specific aims (Baillie 1990, 1991):

1. to establish thresholds that will be used to notify conservation bodies of requirements for further research or conservation action;
2. to identify the stage of the life cycle at which demographic changes are taking place;
3. to provide data that will assist in identifying the causes of such changes; and
4. to distinguish changes in population sizes or demographic rates induced by human activities from those that are due to natural fluctuations.

Changes in numbers of breeding birds have been measured by:

- the BTO/JNCC/RSPB Breeding Bird Survey (BBS) – which began in 1994 and replaced the CBC (below) as the major monitoring scheme for landbirds, after a seven-year overlap. BBS is based on around 3,000 1-km squares, within each of which birdwatchers count and record birds in a standardised manner along a 2-km transect. Because the survey squares are chosen randomly, the results are representative of all habitats and regions. Combined CBC/BBS indices now provide long-running and ongoing population monitoring for many common birds.
- the *Common Birds Census* (CBC) – which ran from 1962 to 2000. This scheme mapped the breeding territories of common birds through intensive fieldwork on 200–300 mainly farmland and woodland plots each year, averaging about 70 and 20 ha respectively.
- the Waterways Breeding Bird Survey (WBBS) – which began in 1998 and replaced the WBS (below) as the major monitoring scheme for breeding birds along rivers and canals, after a ten-year overlap. It is a transect scheme akin to BBS but with the transects running alongside linear waterways. Transects comprise up to ten 500-m sections and cover typically 3–3.5 km of bird-rich habitat. Around 250–300 sites are covered each year, mostly randomly selected. Combined WBS/WBBS indices now provide long-running and ongoing population monitoring for many common waterside birds.
- the *Waterways Bird Survey* (WBS) – which ran from 1974 to 2007. WBS observers mapped the territories of birds along rivers, streams and canals on 80–130 plots each year, each on average 4.5 km in length. Around 70 of these sites are currently incorporated within WBBS.
- the Constant Effort Sites scheme (CES) – which began in 1983 and is based on breeding-season bird ringing at over 100 sites. The catching effort is kept constant at each site during each year, so that changes in numbers of birds caught will reflect population changes and not variation in catching effort.
- the Heronries Census – through which counts of 'apparently occupied nests' have been collected from a high proportion of the UK's heronries every year since 1928.


Changes in breeding performance are measured by:

- the Nest Record Scheme – which began in 1939 and collates standardised information on up to 35,000 individual nesting attempts per year. This allows the measurement of:
 - laying dates
 - clutch sizes
 - brood sizes
 - nesting success during egg and chick stages
 - fledglings per breeding attempt (integrating success across all nesting stages).
- CES (see above) – which provides information on overall productivity for a range of species by measuring the ratio of juveniles to adults caught each year.

Changes in survival are measured by:

- the British and Irish Ringing Scheme – which provides information on the finding circumstances and longevity of ringed birds found dead by members of the public.
- CES also provides information on survival rates, based on the recapture of ringed birds at constant-effort sites.
- Further information on survival rates is provided through the Retrapping Adults for Survival scheme (RAS).

The ways in which the schemes fit together are shown in the diagram below, which also demonstrates the way in which the BTO aims to combine all this information, using population models, to elucidate the mechanisms behind the changes we observe in population size.



Combining results from different schemes

Monitoring the changes in the size of a population does not in itself provide sufficient information on which to base an effective conservation strategy (Goss-Custard 1993, Furness & Greenwood 1993). Concurrent monitoring of breeding performance and survival rates is necessary to allow changes in population size to be properly interpreted (Temple & Wiens 1989, Crick *et al.* 2003) and, for long-lived species, can provide early warning of impending conservation problems (Pienkowski 1991).

Where good long-term data sets for breeding performance and survival are lacking, conservation action might have to be taken without an adequate understanding of the mechanisms involved or might need to wait years for detailed research to be undertaken. For many species, however, BTO already holds the necessary data, collected by its volunteers over periods of several decades (Greenwood 2000).

For a long-lived species, a decline in population may not begin until a long period of low survival or reduced reproductive output has already passed. The classic example is that of the [Peregrine](#), which in the UK suffered from poor breeding performance during the 1940s and 1950s due to sub-lethal DDT contamination. This drop in productivity decreased the capacity of the non-breeding section of the population to buffer the severe mortality of breeding adults that occurred due to cyclodiene poisoning from the mid 1950s onwards (Ratcliffe 1993). Monitoring of breeding performance gave an early warning of impending numerical decline (Pienkowski 1991). Another example of a decline in breeding performance that presaged population decline is the catastrophic breeding failures of seabirds, particularly Arctic Terns, in Shetland (Monaghan *et al.* 1989, 1992, Walsh *et al.* 1995, Mavor *et al.* 2003, 2004, Wanless *et al.* 2005).

Farmland birds

During the mid 1980s, the BTO identified rapid declines in the population sizes of several farmland bird species (O'Connor & Shrubbs 1986, Fulløe *et al.* 1995). The BTO has since been able to investigate the demographic mechanisms underlying these declines, using its long-term historical data sets (Siriwardena *et al.* 1998a, 2000a).

This investigation, which was funded by Government and undertaken jointly with Oxford University, looked at changes in population size, breeding performance and survival rates of a variety of species in relation to changing farming practice. It showed that species responded to different aspects of the agricultural environment, but that typically these aspects were linked to intensification or regional specialisation. Declines in survival rates were found to be the main factor driving population decline in these species, with the exception of [Linnet](#), for which the main factor appears to have been a decline in nesting success at the egg stage (Siriwardena *et al.* 2000b). The study was therefore able to eliminate some possible causes of change, and identify areas for future research, thus helping conservation bodies to use their scarce resources productively. This work made an important contribution to the wider programme of work on farmland birds undertaken by many research and conservation organisations (Aebischer *et al.* 2000, Vickery *et al.* 2004).

This report describes a number of other cases where the combined analysis of BTO data sets has helped to identify the causes of population declines, for example on the pages for [Lapwing](#) (Peach *et al.* 1994), [Song Thrush](#) (Baillie 1990, Thomson *et al.* 1997, Robinson *et al.* 2004), [Sedge Warbler](#) (Peach *et al.* 1991), [Willow Warbler](#) (Peach *et al.* 1995a), [Spotted Flycatcher](#) (Freeman & Crick 2003), [Starling](#) (Freeman *et al.* 2002, 2007b), and [House Sparrow](#) (Freeman & Crick 2002). A fully integrated approach, estimating trends in numbers and demographic parameters through a single model containing data from various BTO surveys, has been introduced by Besbeas *et al.* (2002). More recently, the use of state-space models and Bayesian techniques for integrated monitoring has been pioneered by Baillie *et al.* (2009).

Biodiversity Action Plans

The ability to quickly determine the stage of the life-cycle most heavily involved during population declines is particularly important for the conservation agencies when considering the plight of species on the [lists of conservation concern](#). Analysis of BTO data sets, which has already helped to build these lists, is a key point in several of the UK Government's [biodiversity action plans](#) for rapidly declining species. Once conservation actions have been initiated, the BTO's Integrated Population Monitoring programme has a further function, because the success of these actions will be measured and assessed by continued BTO monitoring.

The aims of this report

The report is the latest in a series, begun in 1997, produced under the BTO's partnership with the Joint Nature Conservation Committee (on behalf of Natural England, Scottish Natural Heritage, Natural Resources Wales, and the Council for Nature Conservation and the Countryside) as part of its programme of research into nature conservation.

Only the first two reports were published on paper, with subsequent ones being produced solely as web documents. A complete list of all the previous reports and links to those published online can be found [here](#). The first 12 reports were titled *Breeding Birds in the Wider Countryside: their conservation status* with 'birdtrends' as the link to the web pages and 'wider countryside report' as their informal title. 'BirdTrends' is now the informal title of the report, matching the web link.

The current report in the series is used by conservation practitioners as a ready reference to changes in status among breeding birds in the UK. By publishing it on the BTO website, we aim to make it available to a much wider audience, especially to BTO members and the general birdwatching public. We hope that it also provides a useful resource for schools, colleges and universities, the media, ecological consultants, decision-makers, local government, and the more general world of industry and commerce. In summary, its aims are:

1. To provide, to as wide a readership as possible, a species-by-species overview of the trends in breeding population and reproductive success of birds covered by BTO monitoring schemes since the 1960s, at the UK or UK-country scale.
2. To provide warning alerts to JNCC and country agencies and to other conservation bodies about worrying declines in population size or reproductive success, with special reference to species on the UK red and amber lists.

This document is the result of the sustained fieldwork of many thousands of the BTO's volunteer supporters. Without their enthusiasm, the cause of conservation in the UK would be very much the poorer. The data we present here include information on distributions, from breeding-season and winter atlas projects, and on estimates of the absolute size of breeding populations, which are reported at intervals by the Avian Population Estimates Panel (Stone *et al.* 1997, Baker *et al.* 2006, Musgrove *et al.* 2013). Colonial seabirds, which are well covered by the results of Seabird 2000 (Mitchell *et al.* 2004) and by the JNCC's [Seabird Monitoring Programme](#) (Heubeck 2013), and the majority of species covered by the [Rare Breeding Birds Panel](#) (Holling & RBBP 2007b, 2008, 2009, 2010a, 2010b, 2011b, 2012, 2013), are not included here. Wintering populations of waterfowl are covered by the Wetland Bird Survey annual reports (Austin *et al.* 2014) and by the *WeBS alerts* system (Cook *et al.* 2013).

The main emphasis of this report is on trends in the abundance and demography of individual species. The data on trends in abundance also provide the basis for multi-species *indicators* of bird population changes (Gregory *et al.* 2004). Four indicators of trends in breeding birds are part of the UK Government's 18 [Biodiversity Indicators](#), which track the UK's progress towards [international targets](#) set by the Convention on Biological Diversity in October 2010. This approach has been extended more widely through a collaboration between EBCC, BirdLife and RSPB to produce [pan-European bird indicators](#).

Acknowledgements

Volunteer fieldwork

Our biggest thankyou is to the volunteers who collected the data on which this website is based. The population trends and other results that we present rely on the sustained, long-term fieldwork of many thousands of BTO volunteers. Our knowledge of the conservation status of the UK's bird populations is possible only as a result of their dedication and enthusiasm. The conservation community owes them all an enormous debt of gratitude for their work. We are also very grateful to the many farmers, land managers and landowners who permitted census work, nest recording and ringing to take place on their land.

Report production and analysis

This website presents the latest in a series of reports, prepared within the partnership between the British Trust for Ornithology (BTO) and the Joint Nature Conservation Committee (JNCC) (on behalf of the Council for Nature Conservation and the Countryside, Natural England, Natural Resources Wales and Scottish Natural Heritage), as part of its programme of research into nature conservation.

Mr and Mrs J A Pye's Charitable Settlement provided additional support towards the development of the website.

Our report includes results from the Breeding Bird Survey, which is funded jointly by BTO, JNCC and RSPB. The BBS partners are very grateful to the Environment and Heritage Service in Northern Ireland (now Northern Ireland Environment Agency) and to the Royal Society for the Protection of Birds in Scotland for supporting professional surveys in areas that would otherwise be difficult to cover.

Deborah Procter of JNCC provided helpful discussions, comments and support during the production of this report. Helen Baker, Jacquie Clark, Nigel Clark, David Gibbons, Jeremy Greenwood, Rowena Langston, Ian McLean, Ian Mitchell, David Stroud, Pierre Tellier, Malcolm Vincent and Lawrence Way provided helpful comments on earlier editions of this publication.

The analyses would not have been possible without the hard work of many past and present BTO staff who have organised schemes, collated data sets or overseen analyses, including: Sue Adams, Dawn Balmer, Jeremy Blackburn, Jacquie Clark, Mark Collier, Greg Conway, Rachel Coombes, Humphrey Crick, Steve Freeman, Mark Grantham, Bridget Griffin, Andrew Joys, Mike Raven, Brenda Read, Anna Renwick, Richard Thewlis, Anne Trehitt and Jane Waters.

The work is also heavily dependent on the BTO's computer and database systems overseen jointly by Karen Wright (with Iain Downie). Susan Waghorn, Laura Smith and Mandy Andrews exercised great skill and effort in helping to design and build the website. The site is now maintained by William Skellorn.

We are very grateful to all of the above organisations and individuals for their contributions to this report.

Methods

Eight monitoring schemes have contributed data to this report. Six provide data on changes in abundance: these are the Breeding Bird Survey, Common Birds Census, Waterways Breeding Bird Survey, Waterways Bird Survey, Heronries Census and the Constant Effort Sites ringing scheme. Two schemes, the Nest Record Scheme and Constant Effort Sites, provide data on changes in breeding productivity. Data on survival rates come from detailed analyses of the retrappings and recoveries of ringed birds, from Retrapping Adults for Survival, Constant Effort Sites and the general Ringing Scheme. In addition, information on waterbirds from the [Wetland Bird Survey](#) is included where relevant.

The methodologies of the monitoring schemes are described in turn, including information on fieldwork, data preparation, sampling considerations and the statistical methods used in analysis. Most of the analyses and the preparation of tables and graphs were undertaken using SAS software (SAS 2009, 2010).

The two final parts of the methods section concern the alert system. These deal, first in descriptive terms and second in statistical detail, with the system by which the results of monitoring surveys raise alerts and thereby are brought to the attention of conservation bodies.

Breeding Bird Survey

The BTO/JNCC/RSPB Breeding Bird Survey (BBS) was launched in 1994, following two years of extensive pilot work and earlier desk-based studies. The introduction of the BBS was a move designed to overcome the limitations of the Common Birds Census (CBC), which had monitored bird populations since 1962. In particular, it improves the geographical spread of UK bird monitoring, thus boosting coverage of species and of habitats.

The BBS uses line transects rather than the more intensive territory-mapping method that had been used by the CBC. This change makes the survey relatively quick to undertake, and has been successful in encouraging a large number of volunteers to take part. The average time observers spend per visit on counting birds is only around 90 minutes. Sampling units are the 1x1-km squares of the Ordnance Survey national grid, of which there are some 254,000 in the UK. From these we make random selections for inclusion in the scheme (see Square selection, below). The BBS requires a relatively large sample of survey squares, and the initial aim was to achieve coverage of about 2,500 squares.

An important aspect of BBS is its coordination through a network of volunteer BBS Regional Organisers. Each year, information and survey forms are distributed first to these organisers, who contact volunteers willing to conduct the fieldwork. After the field season, forms are returned to BTO headquarters again via the Regional Organisers, but an alternative, online method for submission of BBS data was introduced in 2003 is now used by a high proportion of observers – see the BBS pages of the main BTO website for details.

Fieldwork involves three visits to each survey square each year. The first is to record details of habitat and to establish or re-check the survey route, while the second and third (termed 'early' and 'late') are to count birds. A survey route is composed of two roughly parallel lines, each 1 km in length, although for practical reasons routes typically deviate somewhat from the ideal. Each of these lines is divided into five sections, making a total of ten 200-m sections, and birds and habitats are recorded within these ten units. The two bird-count visits are made about four weeks apart (ideally in early May and early June), ensuring that late-arriving migrants are recorded. Volunteers record all the birds they see or hear as they walk along their transect routes. Birds are noted in three distance categories (within 25 m, 25–100 m, or more than 100 m on either side of the line, measured at right angles to the transect line), or as in flight. Recording birds within distance bands provides a measure of bird detectability in different habitats and thus allows population densities to be estimated more accurately. The total numbers of each species, excluding juveniles, are recorded in each 200-m transect section and distance category, as well as the timing of the survey and weather conditions.

By 1998, more than 2,300 BBS squares were being surveyed annually, close to the original target of 2,500. Only around a quarter of these plots were covered in 2001, owing to Foot & Mouth Disease access restrictions, but (thanks to our keen observers) the sample recovered immediately to over 2,205 in 2002 and had increased further to 2,327 squares in 2003, 2,532 in 2004, 2,891 in 2005 and 3,307 in 2006. The sample soared to 3,656 in 2007 and is currently running at over 3,400 (Risely *et al.* 2013). Squares are distributed throughout the UK and cover a broad range of habitats, including uplands and urban areas. There are 108 species that are present on 40 or more BBS squares annually and so can be monitored with good precision at the UK scale (Joys *et al.* 2003, Risely *et al.* 2013), although a few present special difficulties because of their colonial or flocking habit or their wide-ranging behaviour. For most of these 108 species, BBS can also assess annual population changes within England alone, using data from 30 or more squares, and for about half the species also within Scotland and Wales as separate units. Sample sizes in Northern Ireland already allow more than 30 species to be indexed annually.

Square selection

Survey squares are chosen randomly using a stratified random sampling approach from within 83 sampling regions. These sampling regions, which in most cases are the standard BTO regions, are the 'strata' (literally layers) of the sample. Survey squares are chosen at random within each region, to a density that varies with the number of BTO members resident there. Regions with larger numbers of potential volunteers are thereby allotted a larger number of squares, enabling more birdwatchers to become involved in these areas. This does not introduce bias into the results because the analysis takes the differences in regional sampling density into account (see below).

Data analysis

Change measures between years are assessed using a log-linear model with Poisson error terms. For each species and square, counts are summed across all sections and distance bands for each visit ('early' and 'late') and the higher value is used in the model (or the single count if the square was visited only once). Counts are modelled as a function of square and year effects. Each observation is weighted by the number of 1-km squares in each region divided by the number of squares counted in that region, to correct for the differences in sampling density between regions. The upper and lower confidence limits of the changes indicate the certainty that can be attached to each change measure. When the limits are both positive or both negative, we can be 85% confident that a real change has taken place (see here for details).

Trends are presented as graphs in which annual population indices are shown alongside a smoothed trend and its 85% confidence limits. A caveat, 'small sample', is provided against the trends for England, Northern Ireland, Wales and Scotland where the mean sample size is between 30 and 40 plots per year. A minimum average sample size of 40 plots is required for the UK trends.

[Go to the BBS section of the main BTO website.](#)

Common Birds Census

The Common Birds Census (CBC) ran from 1962 to 2000 and was the first of the BTO's schemes for monitoring population trends among widespread breeding birds. It has now been superseded for this purpose by BBS.

The CBC was instigated to provide sound information on farmland bird populations in the face of rapid changes in agricultural practice. Although the original emphasis was on farmland, woodland plots were added by 1964. Fieldwork was carried out by a team of 250–300 volunteers. The same observers surveyed the same plots using the same methods year after year. On average, plots were censused for around seven consecutive years but a few dedicated observers surveyed the same sites for more than 30 years. Farmland plots averaged around 70 hectares in extent. Woodland plots were generally smaller, averaging just over 20 hectares. A small number of plots of other habitats, including heathlands and small wetlands, were also surveyed annually, especially before 1985.

A territory-mapping approach was used to estimate the number and positions of territories of each species present on each survey plot during the breeding season. Volunteers visited their survey plots typically eight to ten times between late March and early July and all contacts with birds, either by sight or sound, were plotted on outline maps at a standard scale of 1:2,500. Codes were used to note each bird's species, with sex and age where possible, and also to record activity such as song or nest-building. The registrations were then transferred to species maps and returned to BTO headquarters for analysis. The pattern of registrations on the species maps reveals the numbers of territories for each species. All assessments of territory number were made by a small team of trained BTO staff, applying rigorous guidelines, for maximum consistency between estimates across sites and years. Observers also provided maps and other details of the habitat on their plots. This makes it possible to match the distribution of bird territories with contemporaneous habitat features, providing the potential for detailed studies of bird–habitat relationships.

In 1990, the results from the Common Birds Census were brought together in the book *Population Trends in British Breeding Birds* (Marchant *et al.* 1990). This landmark publication discussed long-term population trends for the years 1962 to 1988 for 164 species, with CBC or Waterways Bird Survey population graphs for around two-thirds of these.

The results from the Common Birds Census (CBC) provided reliable population trends for more than 60 of the UK's commoner breeding species and, through the linking of CBC with BBS, continue to be hugely influential in determining conservation priorities in the UK countryside. The store of detailed maps of almost a million birds' territories, collected through the CBC and maintained by BTO since the early 1960s, is a uniquely valuable resource for investigating the relationships between breeding birds and their environment, over wide temporal and spatial scales.

The weaknesses of the CBC as a monitor of UK bird populations were largely related to the time-consuming nature of both fieldwork and analysis. This inevitably limited the number of volunteers able to participate in the scheme, with the result that areas with few birdwatchers were under-represented. Constrained by its relatively small sample size, CBC concentrated on farmland and woodland habitats. Bird population trends in built-up areas and the uplands were therefore poorly represented. Furthermore, as the plots were chosen by the observers, some may not have been representative of the surrounding countryside and some bias towards bird-rich habitats might be suspected. It is for these reasons that the BBS was introduced in 1994. The two surveys were run in parallel for seven years to allow calibration between the results: for many species, CBC and BBS trends can be linked to form joint CBC/BBS trends that provide ongoing monitoring, continuous since the 1960s (Freeman *et al.* 2003, 2007a).

Validation studies

The CBC was the first national breeding bird monitoring scheme of its kind anywhere in the world and its value has been widely recognised internationally. The territory-mapping method adopted by the CBC is acknowledged as the most efficient and practical way of estimating breeding bird numbers in small areas, and has been well validated. Although intensive nest searches may sometimes reveal more birds, a comparison by Snow (1965) concluded that mapping censuses were a good measure of the true breeding population for 70% of species. Experiments to test differences between observers' abilities to detect birds found that, although there was considerable variation between individual abilities, the observers were consistent from year to year (O'Connor & Marchant 1981). As the CBC relies on data from plots covered by the same observer in consecutive years, this source of bias has no implications for the CBC's ability to identify population trends. It has also been confirmed that the sample of plots from which CBC results are drawn changed little in composition or character over the years (Marchant *et al.* 1990) and that the results of territory analysis are not affected by changes in analysts, once trained (O'Connor & Marchant 1981). Fuller *et al.* (1985) found that farmland CBC plots were representative of ITE lowland land-classes throughout England (excluding the extreme north and southwest), and closely reflected the agricultural statistics for southern and eastern Britain.

Data analysis

Population changes are modelled using a generalised additive model (GAM), a type of log–linear regression model that incorporates a smoothing function (Fewster *et al.* 2000). This has replaced the Mountford model that employed a six-year moving window (Mountford 1982, 1985, Peach & Baillie 1994) and was used to produce annual population indices until 1999, but the principles are similar. These models are also very similar to log–linear Poisson regression as implemented by program TRIM (Pannekoek & van Strien 1996). Counts are modelled as the product of site and year effects on the assumption that between-year changes are homogeneous across plots. Smoothing is used to remove short-term fluctuations (e.g. those caused by periods of severe weather or by measurement error) and thus reveal the underlying pattern of population change. This is achieved by setting the degrees of freedom to about 0.3 times the number of years in the series. Confidence limits on the indices are estimated by bootstrapping (a resampling method; Manly 1991) to avoid making any assumptions about the underlying distribution of counts.

A small number of CBC-only graphs and tabulated trends are presented in this report for species too depleted for annual monitoring to continue. Smoothed indices are plotted as the blue line on these graphs. The two green lines on the graphs, above and below the index line, are the upper and lower 85% confidence limits. Caveats are provided to show where the data suffer from a 'Small sample' if the mean number of plots was less than 20. Data are flagged as 'Unrepresentative?' if the average abundance of a species in 10-km squares containing CBC plots was less than that in other 10-km squares of the species' distribution in the UK (as measured from 1988–91 Breeding Atlas data (Gibbons *et al.* 1993)) or, where average abundances could not be calculated, if expert opinion judged that CBC data may not be representative.

In practice nearly all CBC data included in this report have been combined with BBS data to provide joint CBC/BBS trends, using the methods described in the next section. These methods for producing joint trends represent an extension of those described above.

More information on the

[Common Birds Census](#) (PDF, 87.11 KB)

.

CBC/BBS trends

CBC and BBS have been described separately in earlier sections. This page describes how the results have been combined to derive joint CBC/BBS trends, extending from the 1960s to the present.

As previously noted, the CBC has been an enormously influential project, providing the main source of information on national population levels in the UK since its inception in 1962. Coverage was predominantly in lowland England, where the numbers of potential volunteers are greatest, while coverage was more patchy in more sparsely populated regions and especially the uplands (Marchant *et al.* 1990). CBC plots were situated in a limited number of habitats, predominantly farmland and woodland. Within a large rectangle of southeastern Britain (covering England and Wales south and east from Seascale, Scarborough and Exeter), the plots are nevertheless believed to be broadly representative, at least of lowland land-classes (Fuller *et al.* 1985). For species such as Wood Warbler and Meadow Pipit that have the greater part of their numbers in the far west or north of Britain, however, the CBC may not have accurately reflected UK trends.

The BBS, on account of its more rigorous, stratified random sampling design, and its simplicity in the field, produces better coverage of the previously under-represented regions and habitats. In some early editions of 'Breeding Birds in the Wider Countryside' (e.g. Baillie *et al.* 2002), separate indices were published from CBC and BBS data, for those species with sufficiently large sample sizes. There being no new CBC data since 2000, however, it is unnecessary to present a CBC-only trend – except for those few species that are now so rare that BBS has been unable to contribute.

For most purposes, the presentation and analysis of longer time-series is required, dating back to before the establishment of the BBS but coming right up to the present day. The calculation of 25-year alert designations, as in this report, provides just one example. This need led the BTO to research the compatibility of indices from BBS and CBC data in various years and regions, and the possibility of deriving trustworthy long-term indices from the two data sources in combination (Freeman *et al.* 2003, 2007a). This research suggested that for the vast majority of species considered there was no significant difference between population trends, calculated from the two surveys, based on that part of the country where CBC data are sufficient to support a meaningful comparison. Where a statistically significant difference was found, this was sometimes for very abundant species for which the power to detect even a biologically insubstantial difference was considerable. Within this region, therefore, long-term trends based on CBC and BBS data can be produced for almost all species previously monitored by the CBC alone. For (Freeman *et al.* 2003, 2007a) this was the area covered by Fuller *et al.* (1985), because CBC plots in that region were shown to be representative of lowland farmland there. As this region covers the bulk of England, and for consistency with the rest of this report, we have produced joint indices for CBC/BBS for the whole of England (the CBC/BBS England index), rather than just the English part of the 'Fuller rectangle'.

A second question then is whether one can obtain reliable trends over the same period for the entire UK. That is, since prior to 1994 only CBC data are available, are the population trends within the region well covered by the CBC typical of those for the UK as a whole? The shortage of CBC data in the north and west means that the only way of investigating this is via the BBS data. Significant differences in trends between the area well covered by the CBC and the rest of the UK were found for approximately half the species (see Freeman *et al.* 2003, 2007a, for full details). For such species, a regional bias in CBC data means that no reliable UK index can be produced prior to 1994. In summary, joint population indices dating back to the start of the CBC can continue to be produced for that part of the country well served by the CBC (essentially England) for almost all common species. However, a similar UK index can be produced for only about 50% of species (CBC/BBS UK index).

Data analysis

This report presents joint CBC/BBS trends for the UK and/or England, as appropriate. Ideally the trends would have been estimated using generalised additive models (Fewster *et al.* 2000) but these were too computationally intensive, given the large number of sites involved. Therefore we fitted a generalised linear model, with counts assumed to follow a Poisson distribution, and a logarithmic link function, to the combined CBC/BBS data. Standard errors were calculated via a bootstrapping procedure and there is therefore no need to model overdispersion, as it does not affect the parameter estimates. BBS squares were weighted as in standard BBS trend analyses. CBC plots were assigned the average weight of all BBS squares as this allows them to be incorporated within the analysis while retaining the convention of not applying weights within the BBS sample. The population trend was smoothed using a thin-plate smoothing spline with degrees of freedom about one third the total number of years. Confidence intervals were calculated via a bootstrap procedure. Bootstrap samples were generated by resampling sites from the original data set, with replacement. A generalised linear model was then fitted to each bootstrap replicate and a smoothing spline fitted to the annual population indices as described above. Confidence limits were then calculated as the appropriate percentiles from the sets of smoothed estimates. The overall result is a smoothed trend that is mathematically equivalent to that produced from a generalised additive model. The method of estimation is less statistically efficient because the smoothing is not incorporated within the estimation procedure, and is likely to have resulted in more conservative statistical tests and wider confidence limits. However this compromise was necessary to make it possible to fit the trends within a reasonable amount of computer time (still several weeks).

Data presentation

Indices are plotted on the graphs as annual estimates, with a smoothed trend and its 85% confidence interval. The CBC started on farmland in 1962 and on woodland in 1964. However, the early years of the CBC population indices are strongly influenced by the effects of the unusually severe winters of 1961/62 and 1962/63, as well as by developments in methodology (Marchant *et al.* 1990). Joint CBC/BBS indices have been calculated using only the data from 1966 onwards, therefore, and population changes are calculated back to 1967.

Waterways Bird Survey & Waterways Breeding Bird Survey

Waterways Bird Survey 1974–2007

The Waterways Bird Survey (WBS) monitored the population trends of up to 24 riparian bird species on canals and rivers throughout the UK during the period 1974–2007. WBS used a territory-mapping method like that of its parent scheme, the Common Birds Census, to estimate the breeding population of waterbirds on each of a number of observer-selected survey plots. Detailed territory maps were prepared alongside habitat data that show which features of linear waterways are important to breeding birds. The plots averaged 4.4 km in length. Almost half were slow-flowing lowland rivers with the rest either fast-flowing rivers/streams or canals. In the scheme's closing years there were around 90 plots distributed throughout the UK. The north and west of Britain were better represented by WBS than by the CBC although, as with CBC, coverage outside England was relatively poor (Marchant *et al.* 1990).

All fieldwork was carried out by BTO volunteers. Observers were asked to survey their plots on nine occasions between March and July, mapping all the birds seen or heard onto 1:10,000 ('six-inch') maps. Registrations were then transferred to species maps, which were analysed to reveal the numbers and positions of territories for each species. For the first 20 years all territory analysis was performed by trained headquarters staff but, during 1994–2007, observers mostly completed their own territory analysis, based on the scheme's written guidelines, with results checked and corrected by BTO staff. As WBS employed very similar methods to those of CBC, the validation studies carried out for the latter generally held true for WBS (see CBC section). Marchant *et al.* (1990) found that there had been little change by 1988 in the composition of the WBS sample, in terms of waterway type or geographical spread.

Population changes along waterways have been reported historically for up to 25 riparian species. For specialist waterbirds, including [Little Grebe](#), [Mute Swan](#), [Common Sandpiper](#), [Kingfisher](#), [Sand Martin](#), [Grey Wagtail](#), [Dipper](#) and [Reed Warbler](#), targeted surveys along waterways can provide a better precision of monitoring than is possible through the more generalised BBS surveys. [Goosander](#) is not covered at all as yet by BBS monitoring. Waterways indices can also add a new perspective on trends in waterbirds that are monitored, largely in different habitats, by CBC/BBS. For [Lapwing](#), populations declined rapidly on arable farmland during the late 1980s while numbers on WBS plots, typically representing populations along river floodplains, were more stable. [Yellow Wagtails](#) have declined much more steeply alongside rivers and canals than elsewhere.

Waterways Breeding Bird Survey and joint indices

WBS had similar limitations as a monitoring scheme that led to the CBC's replacement by BBS. In particular, plot distribution was biased geographically and possibly also towards sites that were good for birds, and an intensive survey method was used that severely limited the sample size (Marchant *et al.* 1990). A drawback specific to WBS was that it covered only waterbirds.

BTO addressed these issues by setting up the Waterways Breeding Bird Survey (WBBS), which ran in parallel with WBS from 1998 to 2007 and now outlives it as a permanent annual survey, supplementing BBS. WBBS uses BBS-style transect methods along random waterways, and includes all species of birds (and mammals, too). WBBS has received some of its funding from the Environment Agency.

In a similar development to joint CBC/BBS indices, it has proved possible to link the two waterways schemes to provide joint WBS/WBBS indices, some dating back to 1974, for the species previously covered by WBS (see below).

Data analysis and presentation

Population trends are generated from the combined WBS and WBBS data using a generalised linear model with counts assumed to follow a Poisson distribution and a logarithmic link function. Standard errors were calculated via a bootstrapping procedure involving 199 replications. For presentation in the figures, both the population trend and its confidence limits were also subsequently smoothed using a thin-plate smoothing spline. The overall result is a smoothed trend that is mathematically equivalent to that produced from a generalised additive model, as previously used for the WBS data alone.

More information on

[WBS](#) (PDF, 77.53 KB)
and [WBBS](#).

Heronries Census

As predators at the top of the freshwater food chain, Grey Herons are excellent indicators of environmental health in the countryside. They build large stick nests, mostly in colonies at traditional sites, thus lending themselves to censuses of active nests.

The BTO Heronries Census began in 1928 and is the longest-running breeding-season bird monitoring scheme in the world. The aim of this census is to collect annual nest counts of Grey Herons from as many sites as possible in the United Kingdom. Volunteer observers make counts of 'apparently occupied nests' at heron colonies each year. Changes in the numbers of nests, especially over periods of several years, provide a clear measure of the population trend.

In recent seasons, observers have also counted the nests of Little Egrets *Egretta garzetta*, which have been appearing in an increasing number of southern heronries since the first breeding records in 1996, and even of Cattle Egrets *Bubulcus ibis*, Night-herons *Nycticorax nycticorax* and Spoonbills *Platalea leucorodia*. Since egrets are fully included in the Heronries Census, data are required from all breeding sites, whether or not Grey Herons are also present. Counts of [Cormorant](#) colonies, which often occur alongside heronries, are also welcome (Newson *et al.* 2007, 2013).

Coverage is coordinated through a network of regional organisers. A core of birdwatchers and ringers monitor their local colonies annually, providing a backbone of regular counts. The number of heronries counted each year has grown in recent years to more than 600. Around two-thirds of the heronries in England and Wales are currently counted each year, with more-complete censuses carried out in 1929, 1954, 1964, 1985 and 2003. Historically rather few counts have been made of heronries in Scotland and Northern Ireland, except during the special surveys, but support there for the Heronries Census has been growing fast in recent years. More than 90 heronries in Northern Ireland are currently reported each year.

Data analysis

Population changes are estimated using a ratio-estimators approach derived from that described by Thomas (1993). Essentially, the ratios of the populations in any two (not necessarily consecutive) years of the survey are estimated from counts at sites visited in each of those years. These ratios can be used to estimate the counts at sites that were not visited, and hence build an estimate of the total population. Further modifications have been made to allow for the extinction of colonies and the establishment of new ones (Marchant *et al.* 2004).

Data presentation

On the [Grey Heron](#) page of this report, the UK trend is presented graphically as annual estimates of apparently occupied nests, with a smoothed trend and its 85% confidence limits. The smooth trend line is based on a non-parametric regression model, using thin-plate smoothing splines with degrees of freedom one-third the number of years in the model. Trends are also shown for England and Wales together, and for England, Wales and Scotland alone.

Visit the Heronries Census page of the BTO website.

Constant Effort Sites scheme

The [Constant Effort Sites](#) (CES) scheme uses changes in catch sizes across a network of standardised mist-netting sites to monitor changes in the abundance and breeding success of common passerines in scrub and wetland habitats. At each constant effort site, licensed ringers erect a series of mist nets in the same positions, for the same amount of time, during 12 visits evenly spaced between 1 May and 31 August (Peach *et al.* 1996). Year-to-year changes in the number of adults caught provide a measure of changing population size, while the ratio of young birds to adults in the total catch is used to monitor annual productivity (breeding success). By summing the abundance of young birds between May and August, the CES method should integrate contributions to annual productivity from the entire nesting season, including second and third broods for multi-brooded species, but will also include a small component of mortality during the immediate post-fledging period. More detailed information about analytical methods is given below and were also provided by Peach *et al.* (1998) (abundance) and Robinson *et al.* (2007) (productivity). Between-year recaptures of ringed birds are also used to calculate annual survival rates of adult birds using specialised analytical techniques (Peach 1993).

The CES scheme began in 1983 with 46 sites and now has over 120. The distribution of CES sites tends to reflect the distribution of ringers within Britain and Ireland. The majority are operated in England, and there are small numbers in Scotland, Wales, Northern Ireland and the Republic of Ireland. The CES routinely monitors the populations of 25 species of passerines in scrub and wetland habitats.

Data analysis

Smoothed trends in the abundance of adults and young are separately assessed using a generalised additive model (GAM), with 85% confidence intervals calculated by bootstrapping (Fewster *et al.* 2000). At sites where catching effort in a year falls below the standard 12 visits, but no more than four visits have been missed, annual catch sizes are corrected according to experience during years with complete coverage, by incorporating an offset into the model (see Peach *et al.* 1998 for full details). Sites with fewer than eight visits in a given year are omitted for the year in question.

Annual indices of productivity (young per adult) are estimated from logistic regression models applied to the proportions of juvenile birds in the catch, the year-effects then being transformed to measures of productivity relative to an arbitrary value of 100 in the most recent year. As above, catch sizes are corrected where small numbers of visits have been missed. It should be noted that these indices are only relative figures, and are not estimates of the actual numbers of young produced per adult (Robinson *et al.* 2007).

Annual estimates of adult survival are derived from a form of the standard Cormack–Jolly–Seber capture–mark–recapture model (Lebreton *et al.* 1992), modified to account for the presence of transient birds. Transients are birds passing through the site, or perhaps living on its periphery, and which therefore have a much lower probability of capture than resident birds living in the vicinity of the net rides. The presence of transients thus tends to decrease the estimated survival rates. We allow for this by introducing an additional 'survival period' in the year of first capture (Hines *et al.* 2003). As with our other schemes, we assume survival probabilities vary annually in a similar fashion across all sites, though mean survival probabilities may differ between sites. Because of the standardised capture protocol, we assume that recapture probabilities are site-specific, but constant through time. For each bird we also insert an additional period after the first capture, indicating whether the bird was caught subsequently in the same season. The probability of surviving this period can be regarded as the probability that the bird is resident on the site (that is the probability that it is available for recapture). The survival and recapture probabilities for this initial period are assumed constant across years and sites. Note that the annual estimates of annual survival presented are in fact the probability that adult birds return to the same CE site the following year; this will be lower (to a small but unknown extent) than the true survival rate. We do not estimate survival rates for juvenile birds, because of their much greater propensity to disperse.

Data presentation

Abundance and productivity data are presented graphically with a smoothed trend and its 85% confidence limits. No trend is currently fitted to the survival data, but the individual estimates are presented with 95% confidence limits. A caveat is provided for 'Small samples' when the average number of plots per year is between 10 and 20.

Visit the CES section of the BTO website.

Retrapping Adults for Survival scheme

RAS aims to provide information on adult survival for a range of species in a variety of habitats, particularly those not caught in sufficient numbers on CES sessions or during more general mist-netting. As with CES, between-year recaptures of ringed birds are used to calculate annual survival rates of adults (Peach 1993).

Each RAS project targets an individual species and operates within a defined study area, aiming to catch or resight the majority of the adults breeding within the site each year. RAS ringers often employ colour rings to increase the probability of detecting returning individuals. The minimum annual sample size should ideally be sufficient to include 30 individuals retrapped or resighted from previous years, whilst maintaining a constant trapping/resighting effort. Each RAS study must run for a minimum of five years, but preferably much longer, to allow calculation of long-term trends in survival rate. Examples of analyses of RAS data have been published by Robinson *et al.* (2008, 2010).

The RAS scheme was launched in 1998 and around 130 projects are currently active, covering 50 species in total, and data for nine of these are presented in this report. Study sites are well distributed throughout the UK.

Data analysis and presentation

Annual estimates of adult survival are derived from a form of the standard Cormack–Jolly–Seber capture–mark–recapture model (Lebreton et al. 1992). As with our other schemes, we assume survival probabilities vary annually in a similar fashion across all sites, though mean survival probabilities may differ between sites. Where individuals can be sexed we include a sex-specific intercept, but assume survival varies similarly across years for both sexes; where few individuals of one sex are caught, we exclude these from the models. We model the annual recapture probabilities as a function of either the number days on which the RAS project operated in that year or the amount of effort recorded, choosing the one that best fits the data. Note that the annual estimates of annual survival presented are in fact the probability that adult birds are found to have returned to the same RAS site the following year; this will be lower (to a small but unknown extent) than the true survival rate. We do not estimate survival rates for juvenile birds, because of their much greater propensity to disperse.

Visit the RAS section of the BTO website.

Nest Record Scheme

The BTO's Nest Record Scheme is the largest, longest-running and most highly computerised of such schemes in the world and employs the most advanced and efficient techniques of data gathering, data capture and analysis (Crick *et al.* 2003). BTO now holds more than 1.3 million nest records, of which 69% are already computerised.

The primary aim of the Nest Record Scheme is to monitor the breeding performance of a wide range of UK birds annually as a key part of the BTO's data collection. Periodic reports are published in *BTO News* (e.g. Leech & Barimore 2008) and the significant results communicated immediately to JNCC. Another primary aim is to undertake detailed analyses of breeding performance of species of conservation interest (e.g. Crick *et al.* 1994, Brown *et al.* 1995, Peach *et al.* 1995a, Crick 1997, Chamberlain & Crick 1999, Siriwardena *et al.* 2001, Crick *et al.* 2002, Chamberlain & Crick 2003, Freeman & Crick 2003, Browneet *al.* 2005, Tryjanowski *et al.* 2006, Douglas *et al.* 2010).

The Nest Record Scheme gathers data on the breeding performance of birds in the UK through a network of volunteer ornithologists. Each observer is given a code of conduct that emphasises the responsibility of recorders towards the safety of the birds they record and explains their legal responsibilities. These observers complete standard nest record cards for each nest they find, or submit computerised data, giving details of nest site, habitat, contents of the nest at each visit and evidence for success or failure. When cards are received by the BTO staff, they are checked, sorted and prepared for input and analysis. Data are prioritised for computer input according to their potential for population monitoring and for specific research projects. Those for Schedule 1 species are kept confidential. (These are species protected from disturbance at the nest by Schedule 1 of the [Wildlife and Countryside Act 1981](#): they are generally rare species and the location of their nests may need to be protected from egg collecting (an illegal activity for every wild bird) or other potential disturbance. A special licence is required to visit any nest of a Schedule 1 species.) Computer programs developed by BTO check the data for errors and calculate first-egg date, clutch size, nest loss rates at egg and chick stages.

Currently the BTO collects a total of more than 40,000 records each year for around 180 species. Typically, there are more than 150 records for 50 species and more than 100 for a further 20 species. The quality of records improved substantially in 1990 with the introduction of a new recording card, which promotes greater standardisation and clarity in the information recorded by observers. Nest recording has subsequently become a module within IPMR, the program through which ringing data are currently collected. The general distribution of completed nest records is patchy at the county scale but is more even over larger regions of the UK. Overall, Northern Ireland and parts of Scotland (southeast, Western Isles) and parts of England (West Midlands, southwest) have relatively low coverage, often reflecting observer density. A major analysis of trends over time in various aspects of breeding performance found relatively few differences between major regions, when analysed using analysis of covariance (Crick *et al.* 1993). The scheme receives records from all the UK's major habitats. Most records come from woodland, farmland and freshwater sites, but the scheme also receives data from scrub, grassland, heathland and coastal areas.

Data analysis

Five different variables are analysed for this report: laying date (where day 1 = January 1); clutch size; brood size; and daily nest failure rates during egg and nestling stages, calculated using the methods of Mayfield (1961, 1975) and Johnson (1979) (see Crick *et al.* 2003 for a review).

To minimise the incidence of errors and inaccurately recorded nests, a set of rejection criteria was applied to the data: laying date included only cases where precision was within ± 5 days; clutch size was not estimated for nests which had been visited only once, for nests which were visited when laying could still have been in progress, or for nests which were visited only after hatching; and maximum brood size was calculated only for nests which were observed after hatching. The last variable is an underestimate of brood size at hatching, because observers may miss early losses of individual chicks; it differs from clutch size because some eggs may be lost during incubation or fail to hatch.

Daily failure rates of whole nests were calculated using a formulation of Mayfield's (1961, 1975) method as a logit-linear model with a binomial error term, in which success or failure over a given number of days (as a binary variable) was modelled, with the number of days over which the nest was exposed during the egg and nestling periods as the binomial denominator (Crawley 1993, Etheridge *et al.* 1997, Aebischer 1999). Numbers of exposure days during the egg and nestling periods were calculated as the midpoint between the maximum and minimum possible, given the timing of nest visits recorded on each nest record (note that exposure days refer only to the time span for which data were recorded for each nest and do not represent the full length of the egg or nestling periods). Each calculation assumes that failure rates were constant during the period considered. Violations of this assumption of the Mayfield method can lead to biased estimates if sampling of nests is uneven over the course of each period. It is unlikely that any such bias would vary from year to year so, although absolute failure rates may be biased, annual comparisons should be unaffected (Crick *et al.* 2003). In this report, therefore, we present only temporal trends in daily nest failure rates.

As the combined influence of concurrent trends in these individual breeding parameters on overall productivity is difficult to assess, the estimates produced are used to derive an annual mean estimate of the number of 'fledglings produced per breeding attempt' (FPBA) according to the equation below (Crick *et al.* 2003):

$$FPBA = CS \times HS \times (1 - EF)EP \times (1 - YF)YP$$

where CS represents clutch size, HS represents hatching success, EF and YF represent egg- and chick-stage daily failure rates and EP and YP represent the length of the egg and nestling periods. Standard errors were derived using the formula given by Siriwardena *et al.* (2000b).

Statistical analyses of nest record data were undertaken using SAS programs (SAS 2009). Regressions through annual mean laying dates, clutch sizes and brood sizes were weighted by sample size. Nest survival was analysed by logistic regression. Quadratic regressions were used when the inclusion of a quadratic term provided a significant improvement over linear regression. These are described as 'curvilinear' in the tables on species pages. Significant linear trends are described as 'linear'. The best-fitting regressions (i.e. quadratic or linear) are presented on the figures in this report. Where neither regression is significant, the linear regression line is shown for illustrative purposes.

Data presentation

Results are presented only if the mean sample size of records for a particular variable and species exceeds ten per year, and are presented with a caveat for small sample sizes if the mean number of records contributing data was between ten and 30 per year.

Visit the Nest Record Scheme section of the BTO website.

Alert system

[General approach](#)

[Smoothing population trends](#)

[Years used for analysis](#)

[Confidence limits and statistical testing](#)

[Data-deficient species](#)

General approach

The alert system used within this report is designed to draw attention to developing population declines that may be of conservation concern, and has been described in detail by Baillie & Rehfisch (2006). It also identifies cases where long-term declines have reversed, leading to an improvement in conservation status. It must be stressed that the alerts and reversals reported here are advisory and do not supersede the agreed, longer-term UK conservation listings (Eaton *et al.* 2009; see [PSoB](#) pages). They are based on similar criteria to *Birds of Conservation Concern*, however, and so provide an indication of likely changes at future revisions.

The system is based on statistical analyses of the population trend data for individual species. Alerts seek to identify rapid declines (>50%) and moderate declines (>25% but <50%). These declines are measured over a number of time-scales, depending on the availability of data – the full length of the available time series, and the most recent 25 years, 10 years and 5 years for which change can be estimated. The conservation emphasis is particularly on the longer periods, but short-term changes help to separate declines that are continuing – or accelerating – from those that have ceased or reversed.

The alerts are calculated annually using standard automated procedures. Where species are at the margin of two categories (e.g. a decline of about 25%) they may raise alerts in some years but not others, or different levels of alert in different years.

Data for some species might be biased, owing to possibly unrepresentative monitoring, or imprecise, owing to small sample sizes. Because these data often provide the only information that is available, our general approach is to report all the alerts raised but to flag up clearly any deficiencies in the data.

Smoothing population trends

Bird populations typically show long-term changes that are complex and do not follow simple mathematical trajectories. In addition to the long-term trends, annual population indices also show short-term fluctuations resulting from a combination of natural population variability and statistical error. We use smoothing techniques that aim to extract the long-term pattern of population change, without forcing it to follow any particular shape (such as a straight line or a polynomial curve). These methods remove most of the effects of short-term fluctuations, including natural year-to-year variability, so that the long-term trend is revealed more clearly.

Technical details available here

Years used for analysis

Once a smoothed population trend has been calculated, change measures are calculated from the ratio of the smoothed population indices for the two years of interest. Population indices for the first and last years of a smoothed time series are less reliable than the others, and so we always drop them before calculating alerts. Because the latest year is not included, the alerts are therefore less up-to-date than they could be, but fewer false alarms are generated. The latest year's data points do contribute, however, to the smoothed curve and are dropped only after the smoothing has taken place.

The time it takes BTO to collate and analyse each year's intake of bird monitoring data is another factor affecting the years that can be included in these analyses. Full analyses of data sets are not usually all available until 12–15 months after the end of a particular breeding season. Thus for a report prepared in year x (e.g. 2013) we have analyses of monitoring data up to year $x-1$ (e.g. 2012). As we drop the final year of the smoothed time series, we report here on change measures up to year $x-2$ (e.g. 2011).

Long-term changes for most of the species included in this report are calculated from joint Common Birds Census and Breeding Bird Survey data (CBC/BBS indices), with population changes calculated back to 1967.

Confidence limits and statistical testing

We show 90% confidence limits for population change measures wherever possible. Any decline where the confidence limits do not overlap zero (no change) is regarded as statistically significant and will trigger an alert if it is of sufficient magnitude. Note that, because we are seeking to detect only declines, we are using a one-tailed test – with a P value of 0.05. These confidence limits therefore do not indicate whether increases are statistically significant.

The graphs of population trends show 85% confidence limits because these allow an approximate visual test of whether the difference between the index values for any two given years is statistically significant: if the index values for two given years are assumed to be independent, and normally distributed with standard errors of comparable size (standard errors differing by a factor of up to about 2 are quite acceptable), then to a good approximation the difference between them is significant at the 5% level if there is no overlap in their 85% confidence intervals (Buckland *et al.* 1992, Anganuzzi 1993). This test is fairly robust, and the independence assumption is reasonable if the years are well separated.

Technical details available here

Data-deficient species

There is uncertainty about the reliability of the results for some species, either because data may be unrepresentative or because they are based on a very

small sample of plots. In these cases the cause of the uncertainty is recorded in the comment column of the population change table.

Unrepresentative data

In this report we present joint UK or England CBC/BBS trends only if there was no substantial or statistical difference between the trends from the two schemes over the period when they ran in parallel (Freeman *et al.* 2007a). Thus, since BBS results are drawn from a random sample, the trends are always considered to be representative of the region concerned.

For CBC data representativeness was assessed using the criteria developed by Gibbons *et al.* (1993). Data from the 1988–91 Breeding Atlas were used to compare the average abundance of a given species in 10-km squares with and without CBC plots. If average abundance is higher in squares without CBC plots, it is likely that much of the population is not well sampled by the CBC. In past reports, CBC data for such species were labelled as "unrepresentative". Where there are insufficient data to undertake such calculations, expert opinion was used instead.

Sample size

Sample size is assessed from the average number of plots contributing to the population indices for a given species in each year. A plot with a zero count would be included provided that the species had been recorded there in at least one year and that records for that plot were available for at least two years. Plots where a species has never been recorded do not enter the index calculations. These average sample sizes are shown in column four ('plots') of the population change tables. For CBC, WBS and CES, a mean of between 10 and 20 plots (when rounded to a whole number) is flagged as a small sample. For BBS indices for individual countries a mean in the range 30–40 plots is flagged as a small sample. UK BBS indices are presented only where samples reach at least 40 plots.

Statistical methods for alerts

The alert system page presents an overview of how the alert system works. More detail is given below about the statistical methods used to estimate population changes and their confidence intervals.

General structure of the data

The data for all of the schemes reported here consist of annual counts made over a period of years at a series of sites. They can thus be summarised as a data matrix of sites x years, within which a proportion of the cells contain missing values because not all of the sites are covered every year. Such data can be represented as a simple model:

$$\log(\text{count}) = \text{site effect} + \text{year effect}$$

Each site has a single site-effect parameter. These site parameters are not usually of biological interest but they are important because abundance is likely to differ between sites. The main parameters of interest are the year effects. These can be modelled either with the same number of parameters as years (an annual model), or with a smaller number of parameters, representing a smoothed curve.

A simple annual model would be fitted as a generalised linear model with Poisson errors and a log link function. This is the main model provided by the program TRIM (Pannekoek & van Strien 1996), which is widely used for population monitoring.

Fitting smoothed trends

Our preferred method for generating a smoothed population trend is to fit a smoothed curve to the data directly using a generalised additive model (GAM) (Hastie & Tibshirani 1990, Fewster *et al.* 2000). Thus the model from the previous section becomes:

$$\log(\text{count}) = \text{site effect} + \text{smooth}(\text{year})$$

where smooth (year) represents some smoothing function of the year effect. It was not straightforward to fit GAMs to the bird census data and we have therefore fitted smoothed curves with a similar degree of smoothing to the annual indices (details below).

The non-parametric smoothed curve fitted in our models is based on a smoothing spline. The degree of smoothing is specified by the number of degrees of freedom (df). A simple linear trend has df = 1, whereas the full annual model has df = t-1, where t is the number of years in the time series. Here we set df to be approximately 0.3 times the number of years in the time series (Fewster *et al.* 2000). The degrees of freedom used for the main data sets presented in this report are summarised below.

	Years	Length of time series	df for smoothed index
CBC/BBS	1966–2012	47	15
WBS/WBBS	1974–2012	39	13
Breeding Bird Survey	1994–2012	19	6
Heronries Census	1928–2012	85	28
Constant Effort Sites	1983–2012	30	10

Note that the numbers of years shown here are different from those available for calculating change measures, because we use the whole time series available for analysis (i.e. prior to the truncation of end points), and because we count the number of years in the time series rather than the number of annual change measures.

CBC/BBS, WBS/WBBS and BBS trends

The model fitted to the combined CBC/BBS and WBS/WBBS data is that historically employed for the BBS – a generalised linear model with counts assumed to follow a Poisson distribution and a logarithmic link function. Standard errors were calculated via a bootstrapping procedure involving 199 replications. For presentation in the figures, both the population trend and its confidence limits were also subsequently smoothed using a thin-plate smoothing spline. The overall result is a smoothed trend that is mathematically equivalent to that produced from a generalised additive model.

Heronries Census trends

The Heronries Census data were analysed using a modified sites x years model based on ratio estimation which incorporates information about new colonies (sites) that have been established and other colonies from the sample that are known to have become extinct. The method was developed by Thomas (1993) specifically in relation to the heronries data set. Since then the heronries database has been substantially upgraded and the method has been applied to the full data set (Marchant *et al.* 2004).

Such a method of analysis cannot be easily applied within a GAM framework. Therefore we fitted a smooth curve to the annual population estimates. This was done using PROC TSPLINE of SAS (SAS 2009). This procedure should give very similar estimates to a GAM analysis but it does not provide confidence intervals for the smoothed population trend or the change measures derived from it. Bootstrapped confidence intervals, where available, are thus presented instead for the [Grey Heron](#) trend.

Constant Effort Sites trends

GAMs were fitted to the CES data for catches of adults and juveniles separately with the addition of an offset to correct for missing visits. Confidence limits were fitted using a bootstrap technique to avoid restrictive assumptions about the distribution of the data. Bootstrap samples were drawn from the data by sampling plots with replacement. We generated 199 bootstrap samples from each data set and fitted a GAM to each of them. Confidence limits for the smoothed population indices (85% ci) and change measures (90% ci) were determined by taking the appropriate percentiles from the distributions of the bootstrap estimates, in a similar manner to that employed for the WBS/WBBS trends.

Species

Access the page for a species by clicking its link on the list below. Each species page has alphabetical and taxonomic listings giving access to all the others.

Jump to

[Wildfowl](#)
[Gamebirds](#)
[Seabirds](#)
[Waterbirds](#)

[Hawks](#)
[Waders](#)
[Pigeons](#)
[Owls](#)

[Crows](#)
[Tits](#)
[Larks](#)
[Warblers](#)

[Thrushes](#)
[Sparrows](#)
[Finches](#)
[Buntings](#)

List of species (in [BOU taxonomic order](#))

WILDFOWL

[Mute Swan](#)
[Greylag Goose](#)
[Canada Goose](#)
[Shelduck](#)
[Mallard](#)

[Tufted Duck](#)

[Goosander](#)

GAMEBIRDS

[Red-legged Partridge](#)

[Red Grouse](#)

[Grey Partridge](#)

[Pheasant](#)

WATERBIRDS

[Red-throated Diver](#)

[Cormorant](#)

[Grey Heron](#)

[Little Grebe](#)

[Great Crested Grebe](#)

HAWKS, etc

[Red Kite](#)

[Hen Harrier](#)

[Sparrowhawk](#)

[Buzzard](#)

[Moorhen](#)

[Coot](#)

WADERS

[Oystercatcher](#)

[Golden Plover](#)

[Lapwing](#)

[Ringed Plover](#)

[Curlew](#)

[Common Sandpiper](#)

[Redshank](#)

[Woodcock](#)

[Snipe](#)

PIGEONS, etc

[Feral Pigeon](#)

[Stock Dove](#)

[Woodpigeon](#)

[Collared Dove](#)

[Turtle Dove](#)

[Cuckoo](#)

OWLS, etc

[Barn Owl](#)

[Little Owl](#)

[Tawny Owl](#)

[Nightjar](#)

[Swift](#)

[Kingfisher](#)

[Green Woodpecker](#)
[Great Spotted Woodpecker](#)
[Lesser Spotted Woodpecker](#)
[Kestrel](#)
[Merlin](#)
[Hobby](#)
[Peregrine](#)
[Ring-necked Parakeet](#)
CROWS, etc
[Magpie](#)
[Jay](#)
[Jackdaw](#)
[Rook](#)
[Carrion Crow](#)
[Hooded Crow](#)
[Raven](#)
TITS, etc
[Goldcrest](#)
[Blue Tit](#)
[Great Tit](#)
[Coal Tit](#)
[Willow Tit](#)
[Marsh Tit](#)
LARKS, etc
[Woodlark](#)
[Skylark](#)
[Sand Martin](#)
[Swallow](#)
[House Martin](#)
WARBLERS, etc
[Cetti's Warbler](#)
[Long-tailed Tit](#)
[Wood Warbler](#)
[Chiffchaff](#)
[Willow Warbler](#)
[Blackcap](#)
[Garden Warbler](#)
[Lesser Whitethroat](#)

Whitethroat
[Grasshopper Warbler](#)
[Sedge Warbler](#)
[Reed Warbler](#)
[Nuthatch](#)
[Treecreeper](#)
[Wren](#)
[Starling](#)
[Dipper](#)
THRUSHES, etc
[Ring Ouzel](#)
[Blackbird](#)
[Song Thrush](#)
[Mistle Thrush](#)
[Spotted Flycatcher](#)
[Robin](#)
[Nightingale](#)
[Pied Flycatcher](#)
[Redstart](#)
[Whinchat](#)
[Stonechat](#)
[Wheatear](#)
SPARROWS, etc
[Duncock](#)
[House Sparrow](#)
[Tree Sparrow](#)
[Yellow Wagtail](#)
[Grey Wagtail](#)
[Pied Wagtail](#)
[Tree Pipit](#)
[Meadow Pipit](#)
FINCHES
[Chaffinch](#)
[Bullfinch](#)
[Greenfinch](#)
[Linnet](#)
[Lesser Redpoll](#)
[Common Crossbill](#)
[Goldfinch](#)
[Siskin](#)
BUNTINGS
[Yellowhammer](#)

[Reed Bunting](#)
[Corn Bunting](#)

Information to aid interpretation of the pages for individual species can be found on the [Key to species texts](#) page.

The following seabird species are not covered by *BirdTrends* but full trend information is available from the [Seabird population trends and causes of change](#) report (JNCC 2012), a separate web site produced by a partnership of which both BTO and JNCC are a part.

SEABIRDS

[Fulmar](#)
[Manx Shearwater](#)
[Storm Petrel](#)
[Leach's Petrel](#)
[Gannet](#)
[Shag](#)
[Arctic Skua](#)
[Great Skua](#)

[Kittiwake](#)
[Black-headed Gull](#)
[Mediterranean Gull](#)

[Common Gull](#)
[Lesser Black-backed Gull](#)
[Herring Gull](#)
[Great Black-backed Gull](#)
[Sandwich Tern](#)

[Common Tern](#)
[Roseate Tern](#)
[Arctic Tern](#)
[Little Tern](#)
[Guillemot](#)
[Razorbill](#)
[Black Guillemot](#)
[Puffin](#)

Key to species texts

The 117 species in this report can be accessed in any order, via alphabetic and taxonomic lists. The taxonomic sequence is that maintained by the British Ornithologists' Union and updated in its current [British List](#). The vernacular and scientific names we use are also drawn from that list. Given this report's limited geographical scope, we use the British rather than the international English names. Depending on the availability of data, the following will be found beneath each species heading:

1. Conservation listings: First, the European conservation category is given, according to current listings by BirdLife International in *Birds in Europe: population estimates, trends and conservation status* (BirdLife International 2004). These update the original listings of Tucker & Heath (1994). For SPECs (Species of European Conservation Concern), the European Threat Status is also given. The current SPEC categories are as follows:
 - SPEC 1
Species of global conservation concern, according to the latest assessments by BirdLife International (2013) (www.birdlife.org/datazone/species/search)
 - SPEC 2
Species with an unfavourable European conservation status, and with more than half of the global breeding or wintering population concentrated in Europe
 - SPEC 3
Species with an unfavourable European conservation status, but with less than half of the global breeding or wintering population within Europe

Other species, not considered to be of European conservation concern, and assessed as 'secure', have no SPEC category but are placed into two further groupings:

- Species with a favourable European conservation status, and with less than half of the breeding or wintering population within Europe (Non-SPEC)
- Species with a favourable European conservation status, but with more than half of the global breeding or wintering population concentrated in Europe (Non-SPEC^E)

The UK conservation listing, given next, is taken from *The Population Status of Birds in the UK* (Eaton *et al.* 2009 (BoCC3); see PSoB pages). These assessments supersede two earlier Birds of Conservation Concern listings (Gibbons *et al.* 1996, Gregory *et al.* 2002). There are three categories, as follows:

- Red – high conservation concern
- Amber – medium conservation concern
- Green – all other species (except introduced species, which are not classified)

The main reason or reasons for listing as red or amber are also given. NB:

- SPEC 1 (globally threatened) species are automatically red listed, and SPEC 1(near threatened), SPEC 2 or SPEC 3 species are amber listed (unless they are introduced or a red-list criterion applies)
- Red or amber listing may stem from decline, localisation or international importance of non-breeding as well as breeding populations in the UK
- Rates of population decline used to assess red and amber listing were generally derived from CBC/BBS results for the 25-year period 1981–2006 or for 1969–2006, and do not take more recent changes into account
- Range declines were generally calculated from the numbers of 10-km squares occupied in the 1968–72 and 1988–91 national breeding atlases (Gibbons *et al.* 1993) but made use of more recent material where available
- Historical decline (in UK over the period 1800–1995) was assessed by literature review

For the first time, BoCC3 has undertaken to classify races, for polytypic species, where two or more races occur regularly in the UK. On occasion the listing for a race may differ from that for the species as a whole. These race-level assessments are given alongside those for species level in our species pages although, since our report is mainly about breeding birds in UK, we have omitted races that occur only as migrants or winter visitors.

Following the signing of the Convention on Biological Diversity at the 'Earth Summit' in Rio de Janeiro in 1992, the statutory conservation bodies in the UK compiled [Biodiversity Action Plans \(BAPs\)](#) for 26 rare or threatened bird species, of which 12 are covered by this report. [ABAP review](#) published in 2007 has concluded that 56 UK bird species now qualify for BAPs and has recommended that certain subspecies (e.g. Fair Isle and St Kilda [Wrens](#)) should now be included as BAP priorities. Our report covers 31 of those species.

Where a UK BAP exists, we give the link to the latest available version. For 'priority species', you will find an onward link to the relevant [INCC priority species page](#). A note appears in this section if the species is one for which the [Rare Breeding Birds Panel](#) requires all breeding records to be submitted.

2. Long-term trend: This summarises the headline trend in population size since 1967 from CBC/BBS, 1975 from WBS/WBBS data, or 1984 from CES data. If there are no data available from these schemes, any assessment of trends covers the period since about the mid 1960s, but may also take historical data into account. Increases and declines that are described as 'shallow', 'moderate' or 'rapid' are generally statistically significant. The following terms are used:
 - Rapid decline: >50% population decline according to CBC/BBS, WBS/WBBS or CES
 - Moderate decline: 25–50% population decline according to CBC/BBS, WBS/WBBS or CES
 - Shallow decline: 10–25% population decline according to CBC/BBS, WBS/WBBS or CES
 - Decline/Increase: information has been derived from sources other than CBC/BBS, WBS/WBBS or CES
 - Probable/Possible increase/decline: information has been derived from sources other than CBC/BBS, WBS/WBBS or CES, and the information is uncertain – see the status summary for details
 - Stable/Fluctuating, with no long-term trend no overall change, or change <10%
 - Uncertain: the information from two monitoring schemes conflicts, or the data are unrepresentative of the species' total UK population – see the status summary for details

- Unknown: no information on the UK population trend is available
 - Shallow increase: 10–50% population increase according to CBC/BBS, WBS/WBBS or CES
 - Moderate increase: 50–100% population increase according to CBC/BBS, WBS/WBBS or CES
 - Rapid increase: >100% population increase according to CBC/BBS, WBS/WBBS or CES
3. UK population size: Periodic reports on population sizes of birds in Britain and in the UK, for the breeding season and for winter, are agreed by the Avian Population Estimates Panel (APEP), on which BTO, GWCT, JNCC, RSPB and WWT are represented. UK population estimates from the Panel's third report (Musgrove *et al.* 2013) are given for each of our species, with a shortened reference (APEP13) and a summary of how each estimate was derived. In a handful of cases, new information potentially superseding APEP13 is also presented.
 4. Key facts table: For 43 species only, there follows a table giving a summary of key facts for migration, habitat and diet.
 5. Status summary: This section provides a brief summary of the trends detailed for the species. Unless there is a separate Causes of change section for the species (see 10, below), it also indicates why such changes might have occurred, if this is known, with reference to any published information.
 6. Population trend graphs: The first, headline graph shows the most representative long-term trend in abundance for the species, and is followed under the 'Population changes in detail' header by further graphs from other schemes, including BBS graphs for separate UK countries, as available. Generally for these graphs there are annual estimates (dots), with a smoothed trend line and its 85% confidence interval. The Methods section provides details about how the trend data are calculated for each scheme. Index values provide a relative measure of population size on an arithmetic scale relative to an arbitrary value of 100 in one of the years of the sequence. If an index value increases from 100 to 200, the population has doubled; if it declines from 100 to 50, it has halved. A narrow confidence interval indicates that the index series is estimated precisely, and a wider one that it is less precise, though the scale of the y-axis must be taken into account. The use of 85% confidence limits allows relatively straightforward comparison of points along the modelled line: non-overlap of the 85% confidence limits is equivalent to a statistically significant difference at approximately the 5% level (Anganuzzi 1993).
 7. Population trends table: This table provides details of summarised percentage changes in population size, over the maximum period from each source, and from the past 25 years, 10 years and 5 years, where these figures are available. Further columns indicate the years included, the average number of census plots included in the analysis for each year, the percentage change (an increase if presented with no sign) and the upper and lower 90% confidence limits of that change. Note that positive and negative percentage changes are not directly equivalent: for example, a decrease of 20% would require an increase of 25% to restore the population to its former level. Where the confidence interval does not include zero, population declines are regarded as statistically significant. The 'Alert' column indicates where a statistically significant population decline is estimated to be of 50% or more (>50) or between 25% and 50% (>25) (see the Alerts section for further details). The 'Comment' column lists any caveats that must be considered when interpreting the estimates. The caveats include:
 - Small sample: For CBC/BBS, WBS/WBBS and CES data, a mean sample size of less than 20 (but more than 10) census plots was available; for BBS data from individual countries, a mean sample of less than 40 (but more than 30) plots was available.
 - Unrepresentative?: Some trends may be marked as possibly unrepresentative of the stated region, owing to the original CBC plots being self-selected by observers and thus potentially a biased sample. This judgment was made either because the species' average abundance in 10-km squares containing CBC plots was less than that in other occupied 10-km squares, as measured by 1988–91 Breeding Atlas timed counts or frequency indices (Gibbons *et al.* 1993) or, where these figures could not be calculated, on expert opinion.
 8. Demographic graphs: Graphs from Constant Effort Sites or Nest Record Scheme data illustrate trends in productivity and survival. NRS graphs show annual means, with error bars to denote ± 1 standard error; and quadratic or linear regression lines with their 95% confidence interval. For CES data, the smoothed trends are plotted with their 85% confidence limits (see CES section for details). CES survival graphs show annual estimates, ± 1 standard error, but trends have not been assessed.
 9. Demography table: This provides details of changes in demographic variables since 1968 (or a more recent year, depending on the availability of data). It lists the period of years concerned, the mean annual sample, the type of trend ('curvilinear' is for a significant quadratic trend, 'linear' is for a significant linear trend, 'none' is where the linear trend is not significantly different from horizontal), the modelled values (from the appropriate regression) for the first and last years and their difference (provided only where the trend is significant), and any caveats that must be considered when interpreting the data. Changes are presented either in the units given or as percentages, and are increases unless a minus sign is shown. The caveat 'Small sample' is given when the mean number of nest record cards contributing annually was in the range 10–30, or when the mean annual number of CES plots recording the species was less than 20 (but more than 10).
 10. Causes of change: For a selection of species (currently 43), information on the causes of the demographic changes we have observed has been removed from the Status summary paragraph and expanded under this heading.
 11. Additional information: Links to atlas maps and tables from previous atlas surveys, and the relevant pages of BirdFacts, BirdTrack and Garden BirdWatch, as available from the BTO web site, are provided on the side bar of each species page.

Summary tables

[Tables of alerts and population increases from CBC/BBS](#)

[Tables of alerts and population increases from WBS/WBBS](#)

[Tables of alerts and population increases from CES](#)

[Tables of population declines and increases from BBS](#)

[Tables of breeding performance](#)

Tables of alerts and population increases from CBC/BBS

- 1a. CBC/BBS UK alerts – long term
- 1b. CBC/BBS England alerts – long term
- 2a. CBC/BBS UK alerts – 25 years
- 2b. CBC/BBS England alerts – 25 years
- 3a. CBC/BBS UK alerts – 10 years
- 3b. CBC/BBS England alerts – 10 years
- 4a. CBC/BBS UK alerts – 5 years
- 4b. CBC/BBS England alerts – 5 years
- 5a. CBC/BBS UK population increases of >50% – long term
- 5b. CBC/BBS England population increases of >50% – long term

Tables of alerts and population increases from WBS/WBBS

1. WBS/WBBS alerts – long term
2. WBS/WBBS alerts – 25 years
3. WBS/WBBS alerts – 10 years
4. WBS/WBBS alerts – 5 years
5. WBS/WBBS population increases of >50% – long term

Tables of alerts and population increases from CES

1. CES adults alerts – long term
2. CES adults alerts – 25 years
3. CES adults alerts – 10 years
4. CES adults alerts – 5 years
5. CES adults population increases of >50% – long term

Tables of population declines and increases from BBS

1. BBS – UK alerts – long term
2. BBS – England alerts – long term
3. BBS – Scotland alerts – long term
4. BBS – Wales alerts – long term
5. BBS – Northern Ireland alerts – long term
6. BBS – UK alerts – 10 years
7. BBS – England alerts – 10 years
8. BBS – Scotland alerts – 10 years
9. BBS – Wales alerts – 10 years
10. BBS – Northern Ireland alerts – 10 years
11. BBS – UK alert – 5 years
12. BBS – England alerts – 5 years
13. BBS – Scotland alerts – 5 years
14. BBS – Wales alerts – 5 years
15. BBS – Northern Ireland alerts – 5 years
16. BBS – UK population increases of >50%
17. BBS – England population increases of >50%
18. BBS – Scotland population increases of >50%
19. BBS – Wales population increases of >50%
20. BBS – Northern Ireland population increases of >50%

Tables of breeding performance

1. Clutch size
2. Brood size
3. Egg-stage nest failure rate

WBS/WBBS alerts & population increases

1. [WBS/WBBS alerts – long term](#)
2. [WBS/WBBS alerts – 25 years](#)
3. [WBS/WBBS alerts – 10 years](#)
4. [WBS/WBBS alerts – 5 years](#)
5. [WBS/WBBS population increases of >50% – long term](#)

1. Table of alerts for WBS/WBBS waterways 1975-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Yellow Wagtail	36	24	-95	-99	-89	>50	Small sample
Snipe	36	13	-93	-99	-76	>50	
Pied Wagtail	36	112	-67	-75	-60	>50	
Redshank	36	24	-63	-89	-36	>50	
Reed Bunting	36	85	-59	-71	-41	>50	
Grey Wagtail	36	96	-58	-67	-44	>50	
Common Sandpiper	36	47	-44	-58	-32	>25	
Sedge Warbler	36	71	-37	-57	-9	>25	
Moorhen	36	123	-30	-45	-4	>25	
Dipper	36	64	-30	-47	-8	>25	

2. Table of alerts for WBS/WBBS waterways 1986-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Yellow Wagtail	25	22	-93	-97	-85	>50	Small sample
Snipe	25	15	-91	-98	-80	>50	
Lapwing	25	76	-67	-79	-50	>50	
Redshank	25	26	-62	-81	-45	>50	
Common Sandpiper	25	57	-49	-60	-40	>25	
Pied Wagtail	25	132	-41	-52	-30	>25	
Curlew	25	51	-37	-55	-5	>25	
Dipper	25	76	-26	-40	-7	>25	
Grey Wagtail	25	115	-25	-36	-9	>25	

3. Table of alerts for WBS/WBBS waterways 2001-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Snipe	10	23	-50	-71	-20	>25	
Yellow Wagtail	10	23	-49	-70	-10	>25	
Lapwing	10	111	-45	-52	-34	>25	
Curlew	10	77	-44	-52	-33	>25	
Grey Wagtail	10	170	-41	-49	-33	>25	
Redshank	10	31	-39	-56	-20	>25	
Pied Wagtail	10	197	-30	-38	-22	>25	

4. Table of alerts for WBS/WBBS waterways 2006-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
<u>Grey Wagtail</u>	5	150	-42	-48	-34	>25	
<u>Lapwing</u>	5	102	-32	-42	-23	>25	
<u>Snipe</u>	5	21	-29	-58	-3	>25	

5. Table of population increases for WBS/WBBS waterways 1975-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
<u>Oystercatcher</u>	36	47	62	26	158		
<u>Coot</u>	36	61	83	13	228		
<u>Mute Swan</u>	36	79	88	36	187		
<u>Whitethroat</u>	36	80	160	5	344		
<u>Mallard</u>	36	164	214	137	288		

CBC/BBS alerts & population increases

- 1a. [CBC/BBS UK alerts – long term](#)
- 1b. [CBC/BBS England alerts – long term](#)
- 2a. [CBC/BBS UK alerts – 25 years](#)
- 2b. [CBC/BBS England alerts – 25 years](#)
- 3a. [CBC/BBS UK alerts – 10 years](#)
- 3b. [CBC/BBS England alerts – 10 years](#)
- 4a. [CBC/BBS UK alerts – 5 years](#)
- 4b. [CBC/BBS England alerts – 5 years](#)
- 5a. [CBC/BBS UK population increases of >50% – long term](#)
- 5b. [CBC/BBS England population increases of >50% – long term](#)

1a. Table of population alerts for CBC/BBS UK 1967-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Turtle Dove	44	106	-94	-97	-91	>50	
Willow Tit	44	43	-92	-96	-86	>50	
Grey Partridge	44	132	-90	-93	-86	>50	
Spotted Flycatcher	44	127	-89	-93	-85	>50	
Corn Bunting	44	73	-87	-94	-77	>50	
Yellow Wagtail	44	80	-73	-88	-39	>50	
Marsh Tit	44	98	-73	-81	-64	>50	
Little Owl	44	60	-64	-80	-40	>50	
Mistle Thrush	44	552	-59	-65	-51	>50	
Song Thrush	44	895	-57	-63	-49	>50	
Whitethroat	44	596	-56	-68	-42	>50	
Yellowhammer	44	550	-55	-65	-44	>50	
Lapwing	44	301	-54	-71	-29	>50	
Bullfinch	44	328	-36	-48	-22	>25	
Dunnock	44	923	-33	-42	-24	>25	
Reed Bunting	44	248	-27	-42	-10	>25	

1b. Table of population alerts for CBC/BBS England 1967-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Tree Sparrow	44	93	-95	-98	-92	>50	
Turtle Dove	44	105	-94	-96	-91	>50	
Spotted Flycatcher	44	97	-91	-94	-88	>50	
Willow Tit	44	40	-91	-97	-85	>50	
Grey Partridge	44	118	-90	-93	-86	>50	
Starling	44	626	-88	-92	-84	>50	
Tree Pipit	44	49	-86	-92	-74	>50	
Corn Bunting	44	70	-85	-94	-74	>50	
Lesser Redpoll	44	47	-83	-94	-53	>50	
Cuckoo	44	292	-73	-81	-63	>50	
Yellow Wagtail	44	79	-72	-88	-46	>50	
Marsh Tit	44	90	-72	-80	-62	>50	
Linnet	44	449	-71	-79	-63	>50	
Skylark	44	601	-62	-70	-55	>50	

<u>Mistle Thrush</u>	44	Period	449	-61	-68	-54	>50	
Species	(yrs)	Plots	(n)	Change	Lower	Upper	Alert	Comment
Yellowhammer	44		480	-59	-68	-49	>50	
<u>Little Owl</u>	44		57	-58	-75	-34	>50	
<u>Willow Warbler</u>	44		477	-57	-70	-44	>50	
<u>Whitethroat</u>	44		516	-56	-69	-38	>50	
<u>Song Thrush</u>	44		711	-55	-62	-44	>50	
<u>Meadow Pipit</u>	44		188	-48	-75	-22	>25	
<u>Bullfinch</u>	44		264	-39	-52	-22	>25	
<u>Dunnock</u>	44		763	-37	-45	-25	>25	
<u>Lapwing</u>	44		252	-36	-68	-7	>25	

2a. Table of population alerts for CBC/BBS UK 1986-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
<u>Turtle Dove</u>	25	132	-91	-94	-87	>50	
<u>Willow Tit</u>	25	46	-90	-94	-84	>50	
<u>Spotted Flycatcher</u>	25	160	-81	-86	-75	>50	
<u>Grey Partridge</u>	25	176	-76	-82	-71	>50	
<u>Little Owl</u>	25	83	-65	-75	-53	>50	
<u>Yellow Wagtail</u>	25	117	-65	-78	-51	>50	
<u>Corn Bunting</u>	25	106	-65	-82	-46	>50	
<u>Lapwing</u>	25	481	-60	-69	-47	>50	
<u>Yellowhammer</u>	25	855	-50	-56	-43	>25	
<u>Mistle Thrush</u>	25	855	-44	-50	-36	>25	
<u>Marsh Tit</u>	25	129	-41	-55	-22	>25	
<u>Tawny Owl</u>	25	95	-27	-45	-9	>25	

2b. Table of population alerts for CBC/BBS England 1986-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
<u>Turtle Dove</u>	25	130	-91	-93	-88	>50	
<u>Willow Tit</u>	25	41	-89	-94	-84	>50	
<u>Lesser Redpoll</u>	25	48	-88	-95	-78	>50	
<u>Tree Pipit</u>	25	60	-86	-92	-77	>50	
<u>Spotted Flycatcher</u>	25	117	-84	-89	-78	>50	
<u>Starling</u>	25	1013	-80	-84	-77	>50	
<u>Grey Partridge</u>	25	158	-75	-81	-68	>50	
<u>Cuckoo</u>	25	437	-72	-76	-67	>50	
<u>Yellow Wagtail</u>	25	114	-64	-77	-48	>50	
<u>Corn Bunting</u>	25	101	-63	-81	-44	>50	
<u>Tree Sparrow</u>	25	103	-60	-81	-33	>50	
<u>Little Owl</u>	25	81	-59	-70	-46	>50	
<u>Willow Warbler</u>	25	713	-57	-66	-50	>50	
<u>Yellowhammer</u>	25	747	-56	-61	-51	>50	
<u>Lapwing</u>	25	402	-50	-62	-37	>50	
<u>Mistle Thrush</u>	25	689	-49	-55	-42	>25	

House Sparrow	Species	25	Period (yrs)	903	Plots (n)	-43	Change (%)	-57	Lower limit	-23	Upper limit	>25	Alert	Comment
Marsh Tit		25		118		-40		-53		-21		>25		
Skylark		25		970		-32		-39		-22		>25		
Meadow Pipit		25		302		-26		-44		-9		>25		

3a. Table of population alerts for CBC/BBS UK 2001-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
<u>Turtle Dove</u>	10	133	-77	-81	-72	>50	
<u>Willow Tit</u>	10	45	-64	-74	-57	>50	
<u>Little Owl</u>	10	103	-50	-59	-38	>50	
<u>Mistle Thrush</u>	10	1285	-36	-39	-31	>25	
<u>Spotted Flycatcher</u>	10	186	-35	-49	-21	>25	
<u>Lapwing</u>	10	766	-34	-39	-26	>25	
<u>Greenfinch</u>	10	2071	-33	-36	-30	>25	
<u>Grey Partridge</u>	10	225	-31	-42	-23	>25	
<u>Marsh Tit</u>	10	163	-28	-39	-15	>25	
<u>Yellow Wagtail</u>	10	152	-25	-34	-13	>25	

3b. Table of population alerts for CBC/BBS England 2001-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
<u>Turtle Dove</u>	10	131	-78	-81	-73	>50	
<u>Willow Tit</u>	10	40	-62	-72	-55	>50	
<u>Little Owl</u>	10	101	-48	-59	-35	>25	
<u>Spotted Flycatcher</u>	10	131	-47	-56	-36	>25	
<u>Starling</u>	10	1539	-46	-50	-43	>25	
<u>Cuckoo</u>	10	532	-44	-49	-40	>25	
<u>Mistle Thrush</u>	10	1015	-38	-42	-34	>25	
<u>Greenfinch</u>	10	1739	-33	-36	-30	>25	
<u>Tree Pipit</u>	10	77	-28	-43	-9	>25	
<u>Pied Wagtail</u>	10	1072	-28	-32	-24	>25	
<u>Lapwing</u>	10	651	-26	-32	-19	>25	
<u>Marsh Tit</u>	10	147	-25	-37	-13	>25	

4a. Table of population alerts for CBC/BBS UK 2006-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
<u>Turtle Dove</u>	5	113	-63	-69	-57	>50	
<u>Willow Tit</u>	5	46	-47	-59	-37	>25	
<u>Lapwing</u>	5	839	-35	-38	-28	>25	
<u>Greenfinch</u>	5	2283	-35	-38	-34	>25	
<u>Little Owl</u>	5	108	-34	-42	-21	>25	

<u>Mistle Thrush</u>	5	1363	-26	-30	-22	>25	
Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment

4b. Table of population alerts for CBC/BBS England 2006-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
<u>Turtle Dove</u>	5	111	-63	-68	-58	>50	
<u>Willow Tit</u>	5	42	-43	-59	-32	>25	
<u>Greenfinch</u>	5	1928	-35	-38	-34	>25	
<u>Little Owl</u>	5	106	-34	-42	-23	>25	
<u>Starling</u>	5	1669	-32	-36	-29	>25	
<u>Lapwing</u>	5	720	-31	-34	-25	>25	
<u>Spotted Flycatcher</u>	5	132	-31	-44	-20	>25	
<u>Cuckoo</u>	5	553	-27	-32	-23	>25	

5a. Table of population increases of >50% for UK CBC/BBS 1967-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
<u>Wren</u>	44	1080	54	35	76		
<u>Coal Tit</u>	44	386	79	1	188		
<u>Chiffchaff</u>	44	644	91	56	140		
<u>Magpie</u>	44	825	103	68	148		
<u>Reed Warbler</u>	44	66	109	24	333		
<u>Great Tit</u>	44	974	110	82	140		
<u>Jackdaw</u>	44	690	123	38	235		
<u>Mallard</u>	44	567	167	106	249		
<u>Woodpigeon</u>	44	1004	173	55	490		
<u>Coot</u>	44	123	182	75	550		
<u>Mute Swan</u>	44	108	240	47	642		
<u>Nuthatch</u>	44	226	251	162	365		
<u>Blackcap</u>	44	697	272	218	398		
<u>Great Spotted Woodpecker</u>	44	463	405	265	630		

5b. Table of population increases of >50% for England CBC/BBS 1967-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
<u>Wren</u>	44	859	63	43	85		
<u>Reed Warbler</u>	44	63	87	20	234		
<u>Pheasant</u>	44	655	90	53	154		
<u>Goldfinch</u>	44	550	91	41	139		
<u>Chiffchaff</u>	44	549	94	55	155		
<u>Great Tit</u>	44	801	98	72	130		
<u>Magpie</u>	44	699	107	67	173		
<u>Jackdaw</u>	44	554	108	36	262		
<u>Long-tailed Tit</u>	44	401	122	63	240		

Carrion Crow	Species	44Period (yrs)	83Plots (n)	125Change (%)	82Lower limit	186Upper limit	Alert	Includes Hooded Crow	Comment
<u>Coot</u>		44	112	170	72	490			
<u>Stock Dove</u>		44	322	187	102	307			
<u>Woodpigeon</u>		44	807	192	57	533			
<u>Green Woodpecker</u>		44	339	199	119	335			
<u>Mallard</u>		44	479	203	126	283			
<u>Mute Swan</u>		44	93	209	45	611			
<u>Blackcap</u>		44	606	234	181	326			
<u>Nuthatch</u>		44	195	263	163	418			
<u>Great Spotted Woodpecker</u>		44	411	360	230	600			
<u>Buzzard</u>		44	236	712	433	1829			

CES alerts & population increases

1. [CES adults alerts – long term](#)
2. [CES adults alerts – 25 years](#)
3. [CES adults alerts – 10 years](#)
4. [CES adults alerts – 5 years](#)
5. [CES adults population increases of >50% – long term](#)

1. Table of alerts for CES adults 1984-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Willow Warbler	27	89	-66	-73	-57	>50	Small sample
Lesser Whitethroat	27	38	-64	-82	-49	>50	
Reed Bunting	27	59	-62	-72	-48	>50	
Willow Tit	27	18	-57	-85	-21	>50	
Sedge Warbler	27	66	-42	-59	-26	>25	
Song Thrush	27	82	-33	-47	-17	>25	
Reed Warbler	27	56	-29	-45	-9	>25	

2. Table of alerts for CES adults 1986-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Lesser Whitethroat	25	39	-67	-79	-55	>50	Small sample
Willow Tit	25	18	-63	-87	-38	>50	
Willow Warbler	25	93	-62	-68	-52	>50	
Reed Bunting	25	62	-59	-70	-44	>50	
Sedge Warbler	25	70	-50	-62	-38	>25	

3. Table of alerts for CES adults 2001-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Greenfinch	10	48	-38	-53	-24	>25	
Sedge Warbler	10	71	-35	-43	-27	>25	

4. Table of alerts for CES adults 2006-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Greenfinch	5	45	-36	-52	-25	>25	
Robin	5	99	-25	-30	-20	>25	
Reed Bunting	5	65	-25	-35	-14	>25	

5. Table of population increases for CES adults 1984-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
<u>Blackcap</u>	27	91	116	79	163		
<u>Chiffchaff</u>	27	73	261	136	501		

BBS population declines & increases

1. [BBS – UK alerts – 16 years](#)
2. [BBS – England alerts – 16 years](#)
3. [BBS – Scotland alerts – 16 years](#)
4. [BBS – Wales alerts – 16 years](#)
5. [BBS – Northern Ireland alerts – 16 years](#)
6. [BBS – UK alerts – 10 years](#)
7. [BBS – England alerts – 10 years](#)
8. [BBS – Scotland alerts – 10 years](#)
9. [BBS – Wales alerts – 10 years](#)
10. [BBS – Northern Ireland alerts – 10 years](#)
11. [BBS – UK alert – 5 years](#)
12. [BBS – England alerts – 5 years](#)
13. [BBS – Scotland alerts – 5 years](#)
14. [BBS – Wales alerts – 5 years](#)
15. [BBS – Northern Ireland alerts – 5 years](#)
16. [BBS – UK population increases of >50% – 16 years](#)
17. [BBS – England population increases of >50% – 16 years](#)
18. [BBS – Scotland population increases of >50% – 16 years](#)
19. [BBS – Wales population increases of >50% – 16 years](#)
20. [BBS – Northern Ireland population increases of >50% – 16 years](#)

1. Table of declines >25% for BBS UK 1995-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Turtle Dove	16	158	-85	-88	-82	>50	
Willow Tit	16	50	-82	-87	-76	>50	
Wood Warbler	16	52	-69	-78	-54	>50	
Whinchat	16	75	-60	-71	-48	>50	
Grey Partridge	16	225	-53	-60	-45	>50	
Starling	16	1745	-53	-56	-50	>50	
Pied Flycatcher	16	40	-52	-67	-35	>50	
Cuckoo	16	721	-50	-55	-46	>50	
Spotted Flycatcher	16	194	-49	-59	-38	>25	
Curlew	16	510	-45	-51	-39	>25	
Yellow Wagtail	16	157	-45	-53	-34	>25	
Little Owl	16	99	-44	-54	-32	>25	
Redshank	16	84	-42	-56	-25	>25	
Lapwing	16	666	-41	-47	-33	>25	
Swift	16	1026	-39	-46	-32	>25	
Kingfisher	16	54	-39	-54	-15	>25	
Dipper	16	58	-36	-54	-12	>25	
Corn Bunting	16	142	-34	-45	-23	>25	
Grey Wagtail	16	221	-31	-41	-12	>25	
Mistle Thrush	16	1176	-31	-37	-24	>25	
Kestrel	16	648	-30	-37	-23	>25	

2. Table of declines >25% for BBS England 1995-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Turtle Dove	16	156	-85	-88	-81	>50	
Willow Tit	16	45	-81	-87	-74	>50	
Cuckoo	16	562	-65	-67	-62	>50	

Starling	Species	16	Period (yrs)	1429	Plots (n)	-58	Change (%)	-61	Lower limit	-55	Upper limit	>50	Alert	Comment
Spotted Flycatcher		16		135		-56		-64		-45		>50		
Grey Partridge		16		201		-49		-55		-41		>25		
Whinchat		16		32		-48		-66		-27		>25		
Tree Pipit		16		74		-45		-61		-22		>25		
Yellow Wagtail		16		153		-45		-52		-34		>25		
Little Owl		16		96		-42		-52		-29		>25		
Swift		16		887		-39		-45		-31		>25		
Mistle Thrush		16		922		-38		-42		-33		>25		
Kingfisher		16		48		-33		-49		-11		>25		
Curlew		16		326		-32		-38		-23		>25		
Redshank		16		60		-32		-49		-8		>25		
Corn Bunting		16		136		-30		-43		-19		>25		
Willow Warbler		16		913		-29		-34		-23		>25		
Marsh Tit		16		133		-26		-37		-12		>25		

3. Table of declines >25% for BBS Scotland 1995-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Kestrel	16	43	-57	-72	-35	>50	
Swift	16	50	-57	-68	-37	>50	
Lapwing	16	87	-56	-64	-44	>50	
Curlew	16	120	-56	-63	-47	>50	
Starling	16	146	-40	-53	-27	>25	
Rook	16	109	-34	-48	-11	>25	
Grey Wagtail	16	33	-32	-53	0	>25	
Oystercatcher	16	127	-30	-39	-20	>25	
Meadow Pipit	16	200	-29	-37	-21	>25	

4. Table of declines >25% for BBS Wales 1995-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Starling	16	80	-70	-80	-60	>50	
Curlew	16	35	-58	-70	-43	>50	
Goldcrest	16	80	-49	-63	-21	>25	
Yellowhammer	16	35	-48	-64	-33	>25	
Swift	16	66	-44	-61	-11	>25	
Cuckoo	16	56	-34	-48	-21	>25	
Linnet	16	92	-30	-48	-8	>25	

5. Table of declines >25% for BBS Northern Ireland 1995-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
---------	-----------------	--------------	---------------	----------------	----------------	-------	---------

Skylark	16	34	-50	-60	-41	>50	
Species	Period	Plots	Change	Lower	Upper	Alert	Comment
Meadow Pipit	16	64	-29	-41	-7	>25	
	(yrs)	(n)	(%)	limit	limit		

6. Table of declines >25% for BBS UK 2001-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
<u>Turtle Dove</u>	10	132	-77	-81	-72	>50	
<u>Willow Tit</u>	10	45	-64	-73	-57	>50	
<u>Grey Wagtail</u>	10	260	-53	-58	-44	>50	
<u>Little Owl</u>	10	102	-49	-59	-38	>25	
<u>Stonechat</u>	10	203	-49	-56	-37	>25	
<u>Starling</u>	10	1866	-46	-50	-42	>25	
<u>Kingfisher</u>	10	61	-45	-63	-26	>25	
<u>Whinchat</u>	10	69	-44	-55	-29	>25	
<u>Wood Warbler</u>	10	49	-40	-57	-18	>25	
<u>Dipper</u>	10	64	-39	-57	-18	>25	
<u>Mistle Thrush</u>	10	1285	-36	-40	-32	>25	
<u>Goldcrest</u>	10	888	-34	-40	-25	>25	
<u>Spotted Flycatcher</u>	10	186	-34	-49	-17	>25	
<u>Lapwing</u>	10	733	-33	-37	-25	>25	
<u>Curlew</u>	10	545	-33	-41	-26	>25	
<u>Greenfinch</u>	10	2048	-33	-36	-31	>25	
<u>Grey Partridge</u>	10	222	-32	-42	-23	>25	
<u>Cuckoo</u>	10	686	-32	-38	-27	>25	
<u>Redshank</u>	10	93	-30	-44	-13	>25	
<u>Marsh Tit</u>	10	159	-29	-40	-18	>25	
<u>Swift</u>	10	1111	-28	-35	-19	>25	

7. Table of declines >25% for BBS England 2001-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
<u>Turtle Dove</u>	10	130	-78	-81	-72	>50	
<u>Willow Tit</u>	10	40	-63	-72	-51	>50	
<u>Little Owl</u>	10	99	-47	-58	-35	>25	
<u>Starling</u>	10	1520	-47	-50	-43	>25	
<u>Spotted Flycatcher</u>	10	126	-45	-54	-33	>25	
<u>Stonechat</u>	10	95	-44	-56	-30	>25	
<u>Cuckoo</u>	10	519	-43	-47	-38	>25	
<u>Grey Wagtail</u>	10	177	-42	-49	-32	>25	
<u>Redshank</u>	10	68	-40	-53	-22	>25	
<u>Mistle Thrush</u>	10	992	-38	-41	-35	>25	
<u>Whinchat</u>	10	32	-37	-53	-21	>25	
<u>Kingfisher</u>	10	55	-33	-46	-16	>25	
<u>Greenfinch</u>	10	1724	-33	-35	-30	>25	
<u>Snipe</u>	10	106	-32	-43	-16	>25	
<u>Swift</u>	10	957	-28	-35	-21	>25	

<u>Tree Pipit</u>	10	77	-28	-44	-11	>25	
Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
<u>Pied Wagtail</u>	10	1072	-27	-32	-23	>25	
<u>Lapwing</u>	10	622	-26	-30	-19	>25	
<u>Feral Pigeon/Rock Dove</u>	10	615	-26	-33	-18	>25	
<u>Marsh Tit</u>	10	145	-26	-36	-12	>25	

8. Table of declines >25% for BBS Scotland 2001-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
<u>Grey Wagtail</u>	10	37	-64	-74	-45	>50	
<u>Stonechat</u>	10	48	-56	-65	-36	>50	
<u>Goldcrest</u>	10	104	-50	-61	-32	>25	
<u>Starling</u>	10	158	-44	-54	-34	>25	
<u>Mistle Thrush</u>	10	87	-42	-54	-20	>25	
<u>Kestrel</u>	10	44	-40	-62	-12	>25	
<u>Lapwing</u>	10	87	-40	-51	-25	>25	
<u>Curlew</u>	10	115	-40	-50	-30	>25	
<u>Wren</u>	10	248	-37	-42	-27	>25	
<u>Rook</u>	10	114	-37	-51	-19	>25	
<u>Grey Heron</u>	10	52	-31	-47	-11	>25	Non-breeders included
<u>Song Thrush</u>	10	199	-26	-37	-12	>25	

9. Table of declines >25% for BBS Wales 2001-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
<u>Starling</u>	10	81	-52	-61	-43	>50	
<u>Curlew</u>	10	33	-43	-57	-25	>25	
<u>Goldcrest</u>	10	86	-43	-52	-30	>25	
<u>Swift</u>	10	70	-41	-55	-13	>25	
<u>Linnet</u>	10	100	-40	-54	-29	>25	
<u>Greenfinch</u>	10	132	-39	-47	-26	>25	
<u>Grey Heron</u>	10	47	-35	-45	-21	>25	Non-breeders included
<u>Treecreeper</u>	10	42	-31	-49	-7	>25	
<u>Yellowhammer</u>	10	34	-27	-42	-11	>25	
<u>Robin</u>	10	216	-26	-31	-19	>25	

10. Table of declines >25% for BBS Northern Ireland 2001-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
<u>Skylark</u>	10	35	-57	-68	-50	>50	
<u>Greenfinch</u>	10	63	-52	-63	-41	>50	
<u>Meadow Pipit</u>	10	73	-51	-58	-46	>50	
<u>Goldcrest</u>	10	52	-46	-53	-27	>25	

<u>Starling</u>	10	94	-41	-51	-31	>25	
Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
<u>Rook</u>	10	88	-40	-51	-24	>25	
<u>Mistle Thrush</u>	10	70	-38	-52	-20	>25	
<u>Wren</u>	10	108	-34	-37	-23	>25	

11. Table of declines >25% for BBS UK 2006-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
<u>Turtle Dove</u>	5	112	-63	-68	-56	>50	
<u>Stonechat</u>	5	231	-62	-67	-53	>50	
<u>Grey Wagtail</u>	5	279	-51	-54	-40	>50	
<u>Willow Tit</u>	5	45	-46	-59	-35	>25	
<u>Kingfisher</u>	5	65	-43	-52	-33	>25	
<u>Greenfinch</u>	5	2245	-35	-37	-33	>25	
<u>Whinchat</u>	5	76	-34	-47	-18	>25	
<u>Starling</u>	5	1996	-33	-39	-27	>25	
<u>Lapwing</u>	5	784	-32	-36	-26	>25	
<u>Little Owl</u>	5	106	-32	-41	-21	>25	
<u>Wood Warbler</u>	5	53	-32	-49	-10	>25	
<u>Dipper</u>	5	74	-29	-45	-5	>25	
<u>Goldcrest</u>	5	960	-28	-34	-20	>25	
<u>Redshank</u>	5	101	-27	-41	-13	>25	
<u>Mistle Thrush</u>	5	1363	-26	-30	-22	>25	

12. Table of declines >25% for BBS England 2006-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
<u>Turtle Dove</u>	5	110	-63	-68	-57	>50	
<u>Stonechat</u>	5	117	-56	-64	-44	>50	
<u>Willow Tit</u>	5	42	-43	-55	-29	>25	
<u>Kingfisher</u>	5	59	-42	-50	-30	>25	
<u>Grey Wagtail</u>	5	196	-41	-46	-29	>25	
<u>Greenfinch</u>	5	1903	-35	-38	-33	>25	
<u>Starling</u>	5	1638	-33	-36	-29	>25	
<u>Little Owl</u>	5	103	-32	-42	-22	>25	
<u>Lapwing</u>	5	671	-30	-34	-24	>25	
<u>Spotted Flycatcher</u>	5	124	-30	-41	-17	>25	
<u>Snipe</u>	5	130	-26	-34	-13	>25	
<u>Cuckoo</u>	5	530	-26	-31	-22	>25	
<u>Mistle Thrush</u>	5	1044	-25	-28	-22	>25	

13. Table of declines >25% for BBS Scotland 2006-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
<u>Stonechat</u>	5	56	-73	-78	-58	>50	
<u>Grey Wagtail</u>	5	40	-56	-69	-38	>50	
<u>Wren</u>	5	284	-46	-49	-39	>25	
<u>Goldcrest</u>	5	119	-43	-54	-28	>25	
<u>Kestrel</u>	5	44	-40	-60	-17	>25	
<u>Swift</u>	5	59	-38	-51	-11	>25	
<u>Mistle Thrush</u>	5	99	-37	-49	-20	>25	
<u>Lapwing</u>	5	91	-35	-41	-23	>25	
<u>Starling</u>	5	171	-35	-50	-16	>25	
<u>Greenfinch</u>	5	127	-29	-39	-18	>25	
<u>Song Thrush</u>	5	229	-25	-32	-16	>25	

14. Table of declines >25% for BBS Wales 2006-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
<u>Starling</u>	5	77	-40	-51	-29	>25	
<u>Greenfinch</u>	5	134	-39	-45	-31	>25	
<u>Stonechat</u>	5	44	-36	-50	-24	>25	
<u>Grey Heron</u>	5	43	-27	-40	-10	>25	Non-breeders included
<u>Robin</u>	5	216	-25	-29	-19	>25	

15. Table of declines >25% for BBS Northern Ireland 2006-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
<u>Greenfinch</u>	5	65	-52	-61	-43	>50	
<u>Meadow Pipit</u>	5	73	-51	-56	-43	>50	
<u>Goldcrest</u>	5	56	-51	-56	-38	>50	
<u>Wren</u>	5	111	-35	-37	-26	>25	
<u>Skylark</u>	5	33	-33	-42	-22	>25	
<u>Mistle Thrush</u>	5	70	-33	-43	-25	>25	
<u>Reed Bunting</u>	5	39	-28	-42	-12	>25	

16. Table of population increases for BBS UK 1995-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
<u>Jackdaw</u>	16	1665	51	40	65		
<u>Lesser Redpoll</u>	16	159	55	26	96		
<u>Canada Goose</u>	16	465	57	34	94		
<u>Buzzard</u>	16	921	80	62	96		
<u>Gadwall</u>	16	36	86	7	205		
<u>Nuthatch</u>	16	469	88	65	105		
<u>Siskin</u>	16	165	89	42	129		

Chiffchaff	Species	16 Period (yrs)	1453 Plots (n)	93 Change (%)	85 Lower limit	104 Upper limit	Alert	Comment
<u>Goldfinch</u>		16	1553	109	97	122		
<u>Tree Sparrow</u>		16	170	113	71	168		
<u>Crossbill</u>		16	56	130	46	231		
<u>Blackcap</u>		16	1545	133	124	151		
<u>Great Spotted Woodpecker</u>		16	1047	139	125	155		
<u>Greylag Goose</u>		16	178	179	58	379		
<u>Barn Owl</u>		16	46	279	180	497		
<u>Red Kite</u>		16	87	676	369	1265		
<u>Ring-necked Parakeet</u>		16	59	1057	458	3284		

17. Table of population increases for BBS England 1995-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
<u>Jackdaw</u>	16	1333	53	44	62		
<u>Tree Sparrow</u>	16	136	75	39	119		
<u>Gadwall</u>	16	34	82	17	189		
<u>Chiffchaff</u>	16	1221	90	82	101		
<u>Nuthatch</u>	16	397	92	72	113		
<u>Siskin</u>	16	58	96	5	350		
<u>Goldfinch</u>	16	1282	100	88	116		
<u>Blackcap</u>	16	1321	108	99	119		
<u>Great Spotted Woodpecker</u>	16	912	117	104	130		
<u>Buzzard</u>	16	607	167	136	210		
<u>Greylag Goose</u>	16	147	248	137	515		
<u>Barn Owl</u>	16	43	285	181	469		
<u>Ring-necked Parakeet</u>	16	59	1058	451	3122		
<u>Red Kite</u>	16	62	11260	4751	10585		

18. Table of population increases for BBS Scotland 1995-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
<u>Great Tit</u>	16	152	51	26	81		
<u>Bullfinch</u>	16	41	52	5	97		
<u>Lesser Redpoll</u>	16	46	55	12	129		
<u>Dunnock</u>	16	141	60	30	92		
<u>Siskin</u>	16	73	81	28	129		
<u>Tree Pipit</u>	16	33	84	22	150		
<u>Whitethroat</u>	16	79	112	63	196		
<u>House Martin</u>	16	62	114	45	216		
<u>Goldfinch</u>	16	88	164	84	237		
<u>Great Spotted Woodpecker</u>	16	48	329	223	465		
<u>Blackcap</u>	16	57	357	215	534		
<u>Chiffchaff</u>	16	47	414	257	716		

19. Table of population increases for BBS Wales 1995-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
<u>Great Tit</u>	16	170	56	38	78		
<u>Chiffchaff</u>	16	137	75	48	109		
<u>Goldfinch</u>	16	126	79	45	130		
<u>Stonechat</u>	16	36	82	18	179		
<u>House Sparrow</u>	16	122	99	69	137		
<u>Blackcap</u>	16	120	163	118	211		
<u>Great Spotted Woodpecker</u>	16	76	198	136	266		

20. Table of population increases for BBS Northern Ireland 1995-2011

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
<u>Dunnock</u>	16	70	62	1	104		
<u>House Martin</u>	16	42	69	2	171		
<u>Woodpigeon</u>	16	83	93	50	145		
<u>Coal Tit</u>	16	63	93	39	138		
<u>Jackdaw</u>	16	75	107	51	148		
<u>Willow Warbler</u>	16	79	114	55	152		
<u>Hooded Crow</u>	16	81	127	77	189		
<u>Pheasant</u>	16	41	144	41	231		
<u>Great Tit</u>	16	72	176	110	204		

Breeding performance

- 1. [Clutch size](#)
- 2. [Brood size](#)
- 3. [Egg-stage nest failure rate](#)
- 4. [Chick-stage nest failure rate](#)

1. Table of significant trends in Clutch size measured between 1968-2011

Species	Period (yrs)	Mean annual sample	Trend	Predicted in first year	Predicted in last year	Change	Comment
Magpie	43	43	Linear decline	5.77 eggs	4.83 eggs	-0.94 eggs	
Long-tailed Tit	43	42	Curvilinear	7.75 eggs	7.01 eggs	-0.74 eggs	
Peregrine	43	18	Curvilinear	3.89 eggs	3.27 eggs	-0.62 eggs	Small sample
Grey Heron	43	15	Linear decline	4.05 eggs	3.44 eggs	-0.61 eggs	Small sample
Great Tit	43	333	Linear decline	8.02 eggs	7.49 eggs	-0.53 eggs	
Greenfinch	43	89	Linear decline	4.76 eggs	4.56 eggs	-0.2 eggs	
Pied Wagtail	43	65	Linear decline	5.08 eggs	4.93 eggs	-0.15 eggs	
Golden Plover	43	12	Linear decline	3.98 eggs	3.84 eggs	-0.14 eggs	Small sample
Common Sandpiper	43	12	Curvilinear	3.98 eggs	3.85 eggs	-0.13 eggs	Small sample
Linnet	43	123	Linear decline	4.74 eggs	4.62 eggs	-0.12 eggs	
Nightjar	43	19	Linear decline	1.97 eggs	1.88 eggs	-0.09 eggs	Small sample
Grey Wagtail	43	38	Curvilinear	4.74 eggs	4.66 eggs	-0.08 eggs	
Collared Dove	43	43	Linear decline	1.95 eggs	1.88 eggs	-0.07 eggs	
Buzzard	43	35	Curvilinear	2.2 eggs	2.16 eggs	-0.04 eggs	
Stock Dove	43	116	Curvilinear	2.07 eggs	2.1 eggs	0.03 eggs	
Lapwing	43	195	Linear increase	3.71 eggs	3.8 eggs	0.09 eggs	
Carion Crow	43	31	Curvilinear	4.05 eggs	4.17 eggs	0.12 eggs	Includes Hooded Crow
Mistle Thrush	43	32	Linear increase	3.89 eggs	4.04 eggs	0.15 eggs	
Duncock	43	108	Linear increase	3.96 eggs	4.15 eggs	0.19 eggs	
Little Owl	43	22	Linear increase	3.38 eggs	3.65 eggs	0.27 eggs	Small sample
Skylark	43	35	Linear increase	3.37 eggs	3.7 eggs	0.33 eggs	
Redstart	43	50	Curvilinear	5.91 eggs	6.24 eggs	0.33 eggs	
Pied Flycatcher	43	357	Linear increase	6.55 eggs	6.89 eggs	0.34 eggs	
Tree Sparrow	43	278	Curvilinear	4.77 eggs	5.17 eggs	0.4 eggs	
Starling	43	77	Linear increase	4.44 eggs	4.97 eggs	0.53 eggs	
Barn Owl	43	44	Curvilinear	4.29 eggs	4.85 eggs	0.56 eggs	

2. Table of significant trends in Brood size measured between 1968-2011

Species	Period (yrs)	Mean annual sample	Trend	Predicted in first year	Predicted in last year	Change	Comment
Great Tit	43	700	Linear decline	7.19 chicks	6.31 chicks	-0.88 chicks	
Sand Martin	43	58	Curvilinear	3.22 chicks	2.35 chicks	-0.87 chicks	
Blue Tit	43	800	Linear decline	8.13 chicks	7.34 chicks	-0.79 chicks	
Magpie	43	83	Curvilinear	3.23 chicks	2.57 chicks	-0.66 chicks	
Yellow Wagtail	43	12	Linear decline	4.83 chicks	4.28 chicks	-0.55 chicks	Small sample
House Sparrow	43	164	Curvilinear	3.36 chicks	2.86 chicks	-0.5 chicks	
Coal Tit	43	75	Curvilinear	7.35 chicks	6.87 chicks	-0.48 chicks	
Carion Crow	43	80	Curvilinear	2.91 chicks	2.49 chicks	-0.42 chicks	Includes Hooded Crow
Grey Heron	43	90	Linear decline	2.84 chicks	2.43 chicks	-0.41 chicks	
Chiffchaff	43	44	Linear decline	5.1 chicks	4.7 chicks	-0.4 chicks	
Greenfinch	43	111	Linear decline	4.1 chicks	3.75 chicks	-0.35 chicks	
Bullfinch	43	37	Linear decline	4.18 chicks	3.88 chicks	-0.3 chicks	
Raven	43	70	Linear decline	3.15 chicks	2.91 chicks	-0.24 chicks	
Rook	43	79	Curvilinear	2.22 chicks	2.02 chicks	-0.2 chicks	
Pied Wagtail	43	136	Linear decline	4.5 chicks	4.32 chicks	-0.18 chicks	
Turtle Dove	43	15	Curvilinear	1.83 chicks	1.73 chicks	-0.1 chicks	Small sample

<u>Barn Owl</u>	43	360	Curvilinear	3.05 chicks	2.95 chicks	-0.1 chicks	
Species	Period (yrs)	Mean annual sample	Trend	Predicted in first year	Predicted in last year	Change	Comment
<u>Robin</u>	43	231	Curvilinear	4.4 chicks	4.31 chicks	-0.09 chicks	
<u>Blackbird</u>	43	255	Curvilinear	3.35 chicks	3.28 chicks	-0.07 chicks	
<u>Linnet</u>	43	141	Curvilinear	4.11 chicks	4.11 chicks	0 chicks	
<u>Stock Dove</u>	43	176	Curvilinear	1.82 chicks	1.83 chicks	0.01 chicks	
<u>Yellowhammer</u>	43	66	Curvilinear	2.98 chicks	3 chicks	0.02 chicks	
<u>Swallow</u>	43	820	Curvilinear	4.12 chicks	4.16 chicks	0.04 chicks	
<u>Duncock</u>	43	122	Curvilinear	3.41 chicks	3.45 chicks	0.04 chicks	
<u>Spotted Flycatcher</u>	43	128	Curvilinear	3.63 chicks	3.67 chicks	0.04 chicks	
<u>Grey Wagtail</u>	43	80	Curvilinear	4.03 chicks	4.08 chicks	0.05 chicks	
<u>Mute Swan</u>	43	67	Curvilinear	4.42 chicks	4.5 chicks	0.08 chicks	
<u>Buzzard</u>	43	109	Curvilinear	1.89 chicks	1.97 chicks	0.08 chicks	
<u>Great Spotted Woodpecker</u>	43	31	Curvilinear	3.7 chicks	3.82 chicks	0.12 chicks	
<u>Tree Pipit</u>	43	29	Curvilinear	4.3 chicks	4.49 chicks	0.19 chicks	Small sample
<u>Sparrowhawk</u>	43	69	Curvilinear	3.16 chicks	3.39 chicks	0.23 chicks	
<u>Peregrine</u>	43	46	Linear increase	2.35 chicks	2.6 chicks	0.25 chicks	
<u>Willow Warbler</u>	43	149	Linear increase	5.12 chicks	5.38 chicks	0.26 chicks	
<u>Skylark</u>	43	65	Linear increase	3.18 chicks	3.45 chicks	0.27 chicks	
<u>Corn Bunting</u>	43	15	Curvilinear	3.32 chicks	3.59 chicks	0.27 chicks	Small sample
<u>Little Owl</u>	43	47	Linear increase	2.52 chicks	2.84 chicks	0.32 chicks	
<u>Dipper</u>	43	150	Curvilinear	3.45 chicks	3.77 chicks	0.32 chicks	
<u>Martin</u>	43	55	Linear increase	3.51 chicks	3.83 chicks	0.32 chicks	
<u>Tree Sparrow</u>	43	372	Curvilinear	3.81 chicks	4.18 chicks	0.37 chicks	
<u>Redstart</u>	43	90	Curvilinear	5.15 chicks	5.62 chicks	0.47 chicks	
<u>Jay</u>	43	11	Linear increase	3.41 chicks	3.93 chicks	0.52 chicks	Small sample
<u>Wren</u>	43	127	Curvilinear	3.61 chicks	4.21 chicks	0.6 chicks	
<u>Nuthatch</u>	43	76	Linear increase	4.98 chicks	5.61 chicks	0.63 chicks	

3. Table of significant trends in Daily failure rate (eggs) measured between 1968-2011

Species	Period (yrs)	Mean annual sample	Trend	Predicted in first year	Predicted in last year	Change	Comment
<u>Woodlark</u>	43	23	Curvilinear	0.0531 nests/day	0.021 nests/day	-0.0321 nests/day	Small sample
<u>Magpie</u>	43	49	Linear decline	0.0274 nests/day	0.002 nests/day	-0.0254 nests/day	
<u>Dipper</u>	43	108	Curvilinear	0.028 nests/day	0.0036 nests/day	-0.0244 nests/day	
<u>Redshank</u>	43	30	Curvilinear	0.0446 nests/day	0.0206 nests/day	-0.024 nests/day	
<u>Long-tailed Tit</u>	43	59	Linear decline	0.0322 nests/day	0.0085 nests/day	-0.0237 nests/day	
<u>Sand Martin</u>	43	37	Curvilinear	0.0204 nests/day	0.0007 nests/day	-0.0197 nests/day	
<u>Yellowhammer</u>	43	63	Curvilinear	0.0496 nests/day	0.0304 nests/day	-0.0192 nests/day	
<u>Snipe</u>	43	14	Linear decline	0.0318 nests/day	0.0129 nests/day	-0.0189 nests/day	Small sample
<u>Woodpigeon</u>	43	97	Curvilinear	0.0448 nests/day	0.029 nests/day	-0.0158 nests/day	
<u>Carion Crow</u>	43	48	Linear decline	0.0165 nests/day	0.0015 nests/day	-0.015 nests/day	Includes Hooded Crow
<u>Pied Wagtail</u>	43	89	Linear decline	0.0188 nests/day	0.0064 nests/day	-0.0124 nests/day	
<u>Robin</u>	43	227	Curvilinear	0.0241 nests/day	0.0131 nests/day	-0.011 nests/day	
<u>Wood Warbler</u>	43	24	Linear decline	0.0181 nests/day	0.0075 nests/day	-0.0106 nests/day	Small sample
<u>Stock Dove</u>	43	109	Linear decline	0.0157 nests/day	0.0053 nests/day	-0.0104 nests/day	
<u>Starling</u>	43	125	Linear decline	0.0112 nests/day	0.0023 nests/day	-0.0089 nests/day	
<u>Tawny Owl</u>	43	68	Curvilinear	0.0109 nests/day	0.0023 nests/day	-0.0086 nests/day	Nocturnal species
<u>Redstart</u>	43	76	Curvilinear	0.0136 nests/day	0.0052 nests/day	-0.0084 nests/day	
<u>Grey Wagtail</u>	43	57	Linear decline	0.0179 nests/day	0.0096 nests/day	-0.0083 nests/day	
<u>Treecreeper</u>	43	22	Curvilinear	0.0234 nests/day	0.0153 nests/day	-0.0081 nests/day	Small sample
<u>Buzzard</u>	43	29	Linear decline	0.0085 nests/day	0.0005 nests/day	-0.008 nests/day	Small sample
<u>Wheatear</u>	43	16	Curvilinear	0.0079 nests/day	0.0001 nests/day	-0.0078 nests/day	Small sample
<u>Barn Owl</u>	43	30	Linear decline	0.0081 nests/day	0.0005 nests/day	-0.0076 nests/day	Small sample
<u>Nuthatch</u>	43	54	Linear decline	0.0094 nests/day	0.002 nests/day	-0.0074 nests/day	
<u>House Sparrow</u>	43	115	Linear decline	0.0109 nests/day	0.0038 nests/day	-0.0071 nests/day	
<u>Greenfinch</u>	43	123	Curvilinear	0.0261 nests/day	0.0192 nests/day	-0.0069 nests/day	
<u>Wren</u>	43	139	Linear decline	0.0187 nests/day	0.012 nests/day	-0.0067 nests/day	
<u>Marsh Tit</u>	43	21	Linear decline	0.0074 nests/day	0.0011 nests/day	-0.0063 nests/day	Small sample
<u>Kestrel</u>	43	41	Curvilinear	0.0074 nests/day	0.0012 nests/day	-0.0062 nests/day	
<u>Martin</u>	43	23	Linear decline	0.0072 nests/day	0.0017 nests/day	-0.0055 nests/day	Small sample
<u>Tree Sparrow</u>	43	367	Linear decline	0.0085 nests/day	0.0032 nests/day	-0.0053 nests/day	

Species	Period (yrs)	Mean annual sample	Trend	Predicted in first year	Predicted in last year	Change	Comment
Jackdaw	43	69	Linear decline	0.007 nests/day	0.0017 nests/day	-0.0053 nests/day	
Tree Pipit	43	14	Curvilinear	0.048 nests/day	0.0435 nests/day	-0.0045 nests/day	Small sample
Sparrowhawk	43	32	Linear decline	0.0047 nests/day	0.0006 nests/day	-0.0041 nests/day	
Reed Warbler	43	183	Linear decline	0.0168 nests/day	0.0128 nests/day	-0.004 nests/day	
Coal Tit	43	56	Linear decline	0.005 nests/day	0.0014 nests/day	-0.0036 nests/day	
Collared Dove	43	61	Curvilinear	0.0314 nests/day	0.0279 nests/day	-0.0035 nests/day	
Pied Flycatcher	43	436	Curvilinear	0.0057 nests/day	0.0026 nests/day	-0.0031 nests/day	
Great Tit	43	619	Linear decline	0.0053 nests/day	0.0023 nests/day	-0.003 nests/day	
Spotted Flycatcher	43	115	Curvilinear	0.0179 nests/day	0.0156 nests/day	-0.0023 nests/day	
Raven	43	22	Curvilinear	0.0026 nests/day	0.0003 nests/day	-0.0023 nests/day	Small sample
Duncock	43	154	Curvilinear	0.0251 nests/day	0.0231 nests/day	-0.002 nests/day	
Blue Tit	43	717	Linear decline	0.0036 nests/day	0.0019 nests/day	-0.0017 nests/day	
Peregrine	43	23	Curvilinear	0.0015 nests/day	0.0013 nests/day	-0.0002 nests/day	Small sample
Grey Heron	43	17	Curvilinear	0 nests/day	0.0002 nests/day	0.0002 nests/day	Non-breeders include
Hen Harrier	43	10	Curvilinear	0.0002 nests/day	0.0005 nests/day	0.0003 nests/day	Small sample
Curlew	43	22	Curvilinear	0.028 nests/day	0.0291 nests/day	0.0011 nests/day	Small sample
Sedge Warbler	43	41	Curvilinear	0.0147 nests/day	0.0167 nests/day	0.002 nests/day	
Linnet	43	172	Linear increase	0.0185 nests/day	0.0235 nests/day	0.005 nests/day	
Reed Bunting	43	51	Curvilinear	0.0063 nests/day	0.0119 nests/day	0.0056 nests/day	
Whitethroat	43	44	Curvilinear	0.0102 nests/day	0.0166 nests/day	0.0064 nests/day	
Lapwing	43	215	Curvilinear	0.0158 nests/day	0.0238 nests/day	0.008 nests/day	
Chaffinch	43	189	Curvilinear	0.0296 nests/day	0.0378 nests/day	0.0082 nests/day	
Willow Warbler	43	69	Linear increase	0.0094 nests/day	0.0176 nests/day	0.0082 nests/day	
Ringed Plover	43	126	Linear increase	0.0222 nests/day	0.0326 nests/day	0.0104 nests/day	
Moorhen	43	140	Linear increase	0.011 nests/day	0.0217 nests/day	0.0107 nests/day	
Blackbird	43	285	Curvilinear	0.026 nests/day	0.0378 nests/day	0.0118 nests/day	
Whinchat	43	16	Linear increase	0.0065 nests/day	0.0247 nests/day	0.0182 nests/day	Small sample
Nightjar	43	25	Linear increase	0.0128 nests/day	0.0379 nests/day	0.0251 nests/day	Small sample
Oystercatcher	43	160	Curvilinear	0.0146 nests/day	0.0441 nests/day	0.0295 nests/day	

4. Table of significant trends in Daily failure rate (chicks) measured between 1968-2011

Species	Period (yrs)	Mean annual sample	Trend	Predicted in first year	Predicted in last year	Change	Comment
Corn Bunting	43	15	Curvilinear	0.045 nests/day	0.0248 nests/day	-0.0202 nests/day	Small sample
Sand Martin	43	61	Linear decline	0.0175 nests/day	0.0004 nests/day	-0.0171 nests/day	
Skylark	43	53	Linear decline	0.0477 nests/day	0.0307 nests/day	-0.017 nests/day	
Magpie	43	49	Linear decline	0.0167 nests/day	0.001 nests/day	-0.0157 nests/day	
Grey Wagtail	43	56	Linear decline	0.0219 nests/day	0.0074 nests/day	-0.0145 nests/day	
Reed Warbler	43	142	Linear decline	0.0196 nests/day	0.0062 nests/day	-0.0134 nests/day	
Meadow Pipit	43	65	Curvilinear	0.0309 nests/day	0.0177 nests/day	-0.0132 nests/day	
Blackbird	43	233	Linear decline	0.0287 nests/day	0.0189 nests/day	-0.0098 nests/day	
Tree Sparrow	43	253	Linear decline	0.015 nests/day	0.0054 nests/day	-0.0096 nests/day	
Jackdaw	43	66	Linear decline	0.011 nests/day	0.0022 nests/day	-0.0088 nests/day	
Redstart	43	54	Linear decline	0.012 nests/day	0.0034 nests/day	-0.0086 nests/day	
House Sparrow	43	119	Curvilinear	0.0151 nests/day	0.0069 nests/day	-0.0082 nests/day	
Carriion Crow	43	42	Linear decline	0.0078 nests/day	0.0009 nests/day	-0.0069 nests/day	Includes Hooded Crow
Merlin	43	28	Linear decline	0.009 nests/day	0.0022 nests/day	-0.0068 nests/day	Small sample
Yellowhammer	43	50	Curvilinear	0.0434 nests/day	0.0383 nests/day	-0.0051 nests/day	
Stock Dove	43	72	Linear decline	0.0114 nests/day	0.0069 nests/day	-0.0045 nests/day	
Starling	43	146	Linear decline	0.0059 nests/day	0.0016 nests/day	-0.0043 nests/day	
Pied Wagtail	43	98	Linear decline	0.0126 nests/day	0.0086 nests/day	-0.004 nests/day	
Stonechat	43	62	Curvilinear	0.0157 nests/day	0.0125 nests/day	-0.0032 nests/day	
Nuthatch	43	62	Linear decline	0.0047 nests/day	0.0019 nests/day	-0.0028 nests/day	
Tawny Owl	43	103	Curvilinear	0.0033 nests/day	0.0009 nests/day	-0.0024 nests/day	Nocturnal species
Barn Owl	43	136	Linear decline	0.0021 nests/day	0.0002 nests/day	-0.0019 nests/day	
Kestrel	43	70	Linear decline	0.0021 nests/day	0.0008 nests/day	-0.0013 nests/day	
Dipper	43	82	Curvilinear	0.0066 nests/day	0.0054 nests/day	-0.0012 nests/day	
Duncock	43	124	Curvilinear	0.0246 nests/day	0.025 nests/day	0.0004 nests/day	
Great Tit	43	429	Curvilinear	0.0058 nests/day	0.0063 nests/day	0.0005 nests/day	
Blue Tit	43	510	Curvilinear	0.006 nests/day	0.0067 nests/day	0.0007 nests/day	

Swallow	43	499	Linear increase	0.0031 nests/day	0.0043 nests/day	0.0012 nests/day	Comment
Species	Period (yrs)	Mean annual sample	Trend	Predicted in first year	Predicted in last year	Change	
Moorhen	43	52	Linear increase	0.0004 nests/day	0.0018 nests/day	0.0014 nests/day	
Chaffinch	43	129	Curvilinear	0.03 nests/day	0.0321 nests/day	0.0021 nests/day	
Coal Tit	43	59	Linear increase	0.0018 nests/day	0.0042 nests/day	0.0024 nests/day	Small sample
Pied Flycatcher	43	358	Linear increase	0.0038 nests/day	0.0066 nests/day	0.0028 nests/day	
Bullfinch	43	34	Curvilinear	0.0305 nests/day	0.0348 nests/day	0.0043 nests/day	
Willow Warbler	43	128	Linear increase	0.0154 nests/day	0.0204 nests/day	0.005 nests/day	
Nightjar	43	22	Curvilinear	0.0015 nests/day	0.0086 nests/day	0.0071 nests/day	
Linnet	43	122	Linear increase	0.0153 nests/day	0.0227 nests/day	0.0074 nests/day	Small sample
Long-tailed Tit	43	40	Linear increase	0.0075 nests/day	0.0194 nests/day	0.0119 nests/day	
Wood Warbler	43	31	Curvilinear	0.0247 nests/day	0.0406 nests/day	0.0159 nests/day	
Garden Warbler	43	20	Linear increase	0.011 nests/day	0.0273 nests/day	0.0163 nests/day	Small sample
Tree Pipit	43	22	Curvilinear	0.033 nests/day	0.0531 nests/day	0.0201 nests/day	Small sample

Discussion

In this discussion we:

1. Review the latest population change measures and alerts for species that are on the Birds of Conservation Concern (BoCC3) red or amber lists for the UK for reasons of population decline (Eaton *et al.* 2009) ([here](#)).
2. Identify species not on the BoCC3 lists but which raise alerts on account of long-term declines and, conversely, currently listed species where recovery may be sufficient to downgrade their listing status in the future ([here](#)).
3. Briefly review declines along waterways and in scrub and wetland habitats as shown by the WBS/WBBS and CES schemes ([here](#)).
4. Review trends over the last 10 years in species that have shown long-term declines, to identify the extent of ongoing declines and check for any evidence of recovery ([here](#)).
5. Identify those species that have shown rapid long-term population increases ([here](#)).
6. Discuss patterns of changes in breeding performance and relationships between trends in abundance and breeding performance([here](#)).
7. Summarise the overall patterns found ([here](#)).

Except where otherwise indicated, our discussion is based on the best long-term trend that is available for each species. These are the trends presented as the main trend graph for each species. Details of estimating and comparing trends are given in the methods section. Full details of all trends available for each species are given on the species pages. Summary tables of all alerts raised by each scheme are presented in the summary tables.

It should be noted that a number of species included in the BoCC3 red and amber lists are not covered by this report, and that not every species listed amber is in UK decline. Thus tables relating to red or amber list status do not include every species so listed.

Latest long-term alerts

This report uses a standardised system for setting 'alerts' that has been agreed between the providers and users of population monitoring information in the UK. The system provides alerts to population declines of 25–50% and of >50% over short, medium and longer terms (5 years, 10 years and 25+ years respectively). These help to highlight the scale and timing of declines, and act as an aid to interpreting the trend graphs presented. Our main emphasis is on long-term declines measured over the longest period available (usually 44 years) and over 25 years, which is one of the periods used to determine red and amber listing (Eaton *et al.* 2009). Alerts triggered over the short term should be considered as early warnings, indicating that conservation issues may be developing for the species concerned. Some short-term declines might stem, however, from normal fluctuations in abundance, from which the population is able to recover without assistance. The steep decline of a suite of species of similar ecology should be considered as a stronger indication that potential problems may be developing. Details of the alerts and methodology used in this report are given in the methods section.

These alerts are therefore important for conservation practitioners who need to set priorities for conservation action, but we hope that they will also interest more-general readers of the report. Similar alerts for wetland birds are provided by the Wetland Bird Survey (Cook *et al.* 2013).

Where this section discusses conservation-listed species, it uses the current version of these lists, introduced in 2009 and abbreviated as BoCC3. The full paper (Eaton *et al.* 2009) details the criteria by which each listed species qualifies for its red or amber status. All UK breeding birds that are red listed will necessarily have met red-list criteria for UK decline, but amber-listed birds may be listed for reasons other than breeding decline (see Key to species texts).

Long-term trends of 'Birds of Conservation Concern' red-listed species

The species considered in this section are red-listed wholly or partly because of severe UK population declines revealed by annual census data, amounting to more than 50% over the 25-year period 1981–2006 or, in four cases ([Skylark](#), [Song Thrush](#), [Marsh Tit](#) and [Linnet](#)), over the 37-year period 1969–2006. The latest long-term population changes and alerts for these severely declining species are shown in Table A1, over the maximum period available (usually the 44 years 1967–2011) and over 25 years (1986–2011). This table thus updates the figures that were used to produce the current BoCC3 red list.

The 19 species in Table A1 are listed in descending order of longest-term percentage change. [Tree Sparrow](#) heads the table once again, with the strongest long-term decline of any species, despite significant increases in numbers recorded by BBS since 1995. The figures for [Lesser Spotted Woodpecker](#) are likely to be a very large underestimate of the current population change, because the species had by 1999 become too rare for further annual monitoring. Were recent data available, this species might easily surpass Tree Sparrow in the strength of its decline.

Table A1 Latest trends for red-listed species

Species	Period (yrs)	Source	Change (%)	Lower limit	Upper limit	Alert	Comment
Tree Sparrow	44	CBC/BBS England	-95	-98	-92	>50	
Tree Sparrow	25	CBC/BBS England	-60	-81	-33	>50	
Turtle Dove	44	CBC/BBS UK	-94	-97	-91	>50	
Turtle Dove	25	CBC/BBS UK	-91	-94	-87	>50	
Willow Tit	44	CBC/BBS UK	-92	-96	-86	>50	
Willow Tit	25	CBC/BBS UK	-90	-94	-84	>50	
Grey Partridge	44	CBC/BBS UK	-90	-93	-86	>50	
Grey Partridge	25	CBC/BBS UK	-76	-82	-71	>50	
Spotted Flycatcher	44	CBC/BBS UK	-89	-93	-85	>50	
Spotted Flycatcher	25	CBC/BBS UK	-81	-86	-75	>50	
Starling	44	CBC/BBS England	-88	-92	-84	>50	
Starling	25	CBC/BBS England	-80	-84	-77	>50	
Corn Bunting	44	CBC/BBS UK	-87	-94	-77	>50	
Corn Bunting	25	CBC/BBS UK	-65	-82	-46	>50	
Tree Pipit	44	CBC/BBS England	-86	-92	-74	>50	
Tree Pipit	25	CBC/BBS England	-86	-92	-77	>50	
Lesser Redpoll	44	CBC/BBS England	-83	-94	-53	>50	
Lesser Redpoll	25	CBC/BBS England	-88	-95	-78	>50	
Cuckoo	44	CBC/BBS England	-73	-81	-63	>50	
Cuckoo	25	CBC/BBS England	-72	-76	-67	>50	
Yellow Wagtail	44	CBC/BBS UK	-73	-88	-39	>50	
Yellow Wagtail	25	CBC/BBS UK	-65	-78	-51	>50	
Marsh Tit	44	CBC/BBS UK	-73	-81	-64	>50	
Marsh Tit	25	CBC/BBS UK	-41	-55	-22	>25	
Linnet	44	CBC/BBS England	-71	-79	-63	>50	
Linnet	25	CBC/BBS England	-5	-23	13		

House Sparrow	34	CBC/BBS England	-68	-77	-57	>50	
Species	Period	Source	Change	Lower	Upper	Alert	Comment
House Sparrow	25 (yrs)	CBC/BBS England	-43 (%)	-57limit	-23limit	>25	
<u>Skylark</u>	44	CBC/BBS England	-62	-70	-55	>50	
<u>Skylark</u>	25	CBC/BBS England	-32	-39	-22	>25	
<u>Lesser Spotted Woodpecker</u>	31	CBC to 1999	-60	-81	40		Small sample
<u>Lesser Spotted Woodpecker</u>	25	CBC to 1999	-73	-86	-31	>50	Small sample
<u>Song Thrush</u>	44	CBC/BBS UK	-57	-63	-49	>50	
<u>Song Thrush</u>	25	CBC/BBS UK	-4	-14	7		
<u>Yellowhammer</u>	44	CBC/BBS UK	-55	-65	-44	>50	
<u>Yellowhammer</u>	25	CBC/BBS UK	-50	-56	-43	>25	
<u>Lapwing</u>	44	CBC/BBS UK	-54	-71	-29	>50	
<u>Lapwing</u>	25	CBC/BBS UK	-60	-69	-47	>50	

For [Marsh Tit](#), [Linnet](#), [House Sparrow](#), [Skylark](#), [Song Thrush](#) and [Yellowhammer](#) the 25-year change is less than 50%, indicating that, while these species meet red-list criteria for long-term change, their recent rate of decline has been slower than for most other red-listed birds. For [Linnet](#) and [Song Thrush](#), the 25-year trend is effectively stable.

Long-term trends of declining amber-listed species

There are 40 amber-listed species that are included in this report, of which about half (19 species) are listed because of UK population declines over the periods 1981–2006 or 1969–2006. Long-term trends are available from annual census data for 13 of these species, which are listed in Table A2 in descending order of longest-term percentage change (normally over the 43 years 1967–2010). Where available the 25-year change (1985–2010) is also shown.

Table A2 Latest trends for declining amber-listed species

Species	Period (yrs)	Source	Change (%)	Lower limit	Upper limit	Alert	Comment
<u>House Martin</u>	44	CBC/BBS England	-65	-91	0		
<u>House Martin</u>	25	CBC/BBS England	-55	-83	40		
<u>Redshank</u>	36	WBS/WBBS waterways	-63	-89	-36	>50	
<u>Redshank</u>	25	WBS/WBBS waterways	-62	-81	-45	>50	
<u>Mistle Thrush</u>	44	CBC/BBS UK	-59	-65	-51	>50	
<u>Mistle Thrush</u>	25	CBC/BBS UK	-44	-50	-36	>25	
<u>Grey Wagtail</u>	36	WBS/WBBS waterways	-58	-67	-44	>50	
<u>Grey Wagtail</u>	25	WBS/WBBS waterways	-25	-36	-9	>25	
<u>Willow Warbler</u>	44	CBC/BBS England	-57	-70	-44	>50	
<u>Willow Warbler</u>	25	CBC/BBS England	-57	-66	-50	>50	
<u>Whitethroat</u>	44	CBC/BBS UK	-56	-68	-42	>50	
<u>Whitethroat</u>	25	CBC/BBS UK	133	101	166		
<u>Little Grebe</u>	36	WBS/WBBS waterways	-49	-73	10		
<u>Little Grebe</u>	25	WBS/WBBS waterways	-23	-53	43		
<u>Meadow Pipit</u>	44	CBC/BBS England	-48	-75	-22	>25	
<u>Meadow Pipit</u>	25	CBC/BBS England	-26	-44	-9	>25	
<u>Common Sandpiper</u>	36	WBS/WBBS waterways	-44	-58	-32	>25	
<u>Common Sandpiper</u>	25	WBS/WBBS waterways	-49	-60	-40	>25	
<u>Curlew</u>	44	CBC/BBS England	-39	-71	24		
<u>Curlew</u>	25	CBC/BBS England	-28	-57	4		
<u>Bullfinch</u>	44	CBC/BBS UK	-36	-48	-22	>25	
<u>Bullfinch</u>	25	CBC/BBS UK	0	-13	15		
<u>Dunnock</u>	44	CBC/BBS UK	-33	-42	-24	>25	
<u>Dunnock</u>	25	CBC/BBS UK	17	7	29		
<u>Reed Bunting</u>	44	CBC/BBS UK	-27	-42	-10	>25	
<u>Reed Bunting</u>	25	CBC/BBS UK	10	-9	32		

Species	Period (yrs)	Source	Change (%)	Lower limit	Upper limit	Alert	Comment
---------	--------------	--------	------------	-------------	-------------	-------	---------

Our best estimate of long-term change in the English [House Martin](#) population shows a decline of more than 50%, but statistically it is not significantly different from no change and therefore no alerts are raised formally for this species. It may be a candidate for red listing but for now is perhaps best regarded as 'data deficient'. BBS data indicate increases in Scotland and Northern Ireland, however.

Five species raise high alerts, having shown significant declines of greater than 50%, and are therefore potential red-list candidates. [Redshank](#) has declined steeply in lowland Britain, according to waterways surveys, raising high alerts; a major decline is also documented for its breeding sites on saltmarsh, and BBS data show that decline has occurred recently across a wide range of habitats. Accelerating decline for [Mistle Thrush](#) has taken its 44-year trend well past the 50% threshold for rapid decline. [Grey Wagtail](#) has also dropped steeply in recent seasons, raising alerts in the long-term and 25-year reporting periods. English [Willow Warblers](#) meet the red-list criterion for population decline, but there has been little change in Wales and overall increase in Scotland and Northern Ireland since 1995. [Whitethroat](#) shows substantial decline over the 44-year period, since this includes the extraordinary population crash that occurred between 1968 and 1969, but the 25-year period has seen a considerable reversal of this decrease.

Five species raise only the lower level of alert. [Meadow Pipit](#) and [Common Sandpiper](#) meet the 25% criterion (equivalent to amber listing) in both periods. Populations of [Bullfinch](#), [Dunnock](#) and [Reed Bunting](#) are recovering and show stable or increasing trends over the shorter, 25-year period. Data for [Little Grebe](#) and [Curlew](#) suggest a similar overall rate of decline but their trends should be treated with caution, as the confidence intervals are very wide. For [Little Grebe](#), BBS results show little change since 1995.

Long-term declines of species that are not currently red or amber listed (for declines)

This section of the report draws attention to declines which currently surpass red or amber criteria but which were not recognised in the 2009 listings (Table A3). These species may be candidates for conservation listing (for declines) at the next review.

Table A3 Long-term trends for declining species not on the red or amber list (for declines)

Species	Period (yrs)	Source	Change (%)	Lower limit	Upper limit	Alert	Comment
Snipe	36	WBS/WBBS waterways	-93	-99	-76	>50	Small sample
Snipe	25	WBS/WBBS waterways	-91	-98	-80	>50	Small sample
Woodcock	31	CBC to 1999	-74	-88	-49	>50	Small sample
Woodcock	25	CBC to 1999	-76	-88	-51	>50	Small sample
Little Owl	44	CBC/BBS UK	-64	-80	-40	>50	
Little Owl	25	CBC/BBS UK	-65	-75	-53	>50	
Dipper	36	WBS/WBBS waterways	-30	-47	-8	>25	
Dipper	25	WBS/WBBS waterways	-26	-40	-7	>25	
Tawny Owl	25	CBC/BBS UK	-27	-45	-9	>25	

The WBS/WBBS trend for [Snipe](#) is based now on a very small sample of plots, the species having deserted so many of its former riverside haunts. It is currently amber-listed solely because it is a Species of European Conservation Concern (SPEC category 3) through its moderate decline on the European scale (BiE04). There is ample evidence, however, that its breeding range has contracted sharply, especially in lowland England.

Similarly, [Woodcock](#) is currently amber-listed solely because it is a Species of European Conservation Concern (SPEC category 3) through its moderate decline on the European scale (BiE04). The only UK census data indicating a trend are from CBC, which recorded steep declines. Samples were small, however, and the CBC's mapping method was not well suited to monitoring this species: for these reasons, the CBC trend is no longer used to support the species' conservation listing.

[Little Owl](#) also meets red-list criteria for population decline but, as an introduced species, is not eligible for any conservation listing. [Tawny Owl](#) has passed the criteria for amber listing, with a decline >25% over the 25-year period. Although the trends are statistically significant, it should be borne in mind that neither CBC nor BBS field techniques cater well for nocturnal and crepuscular species like these.

Fluctuations in the UK [Dipper](#) population since 1974 appear to be underlain by decrease. The current estimate of long-term change clearly raises an alert and since the 2012 report the 25-year change has also passed this threshold.

Declines along linear waterways

The Waterways Bird Survey and Waterways Breeding Bird Survey supplement the results from CBC and BBS, which are more broadly-based surveys, by measuring trends in bird populations alongside rivers and canals. Joint WBS/WBBS trends allow trend assessment to be continuous since 1974 for up to 25 species that were covered by WBS. WBBS, ongoing since 1998, includes all bird species but waterways trends are presented here only for waterway-specialist species, for which joint WBS/WBBS trends are available. A full set of up-to-date WBS/WBBS trends can be obtained from the [Table generator](#).

For several species, such as [Canada Goose](#), [Goosander](#) and [Kingfisher](#), that are abundant in waterway habitats, the WBS/WBBS trend provides our headline information on population trends. For [Redshank](#), [Little Grebe](#), [Common Sandpiper](#), [Grey Wagtail](#), [Snipe](#) and [Dipper](#), which are also in this category and are in decline, latest trends appear in Tables A2 or A3, as appropriate. Even where WBS/WBBS is not the headline trend for a species, however, the waterways data provide valuable supplementary information from this sensitive habitat.

Table A4 lists all statistically significant declines of greater than 25% recorded from the full period of waterway monitoring (nominally 1975–2011). It does not include [Little Grebe](#), for which the decline is not statistically significant (Table A2).

Table A4 Population declines of greater than 25% recorded by the joint Waterways Bird Survey/Waterways Breeding Bird Survey (WBS/WBBS) between 1975 and 2011

Species	Period (yrs)	Source	Change (%)	Lower limit	Upper limit	Alert	Comment
Yellow Wagtail	36	WBS/WBBS waterways	-95	-99	-89	>50	Small sample
Snipe	36	WBS/WBBS waterways	-93	-99	-76	>50	
Pied Wagtail	36	WBS/WBBS waterways	-67	-75	-60	>50	
Redshank	36	WBS/WBBS waterways	-63	-89	-36	>50	
Reed Bunting	36	WBS/WBBS waterways	-59	-71	-41	>50	
Grey Wagtail	36	WBS/WBBS waterways	-58	-67	-44	>50	
Lapwing	31	WBS/WBBS waterways	-47	-72	-7	>25	
Common Sandpiper	36	WBS/WBBS waterways	-44	-58	-32	>25	
Sedge Warbler	36	WBS/WBBS waterways	-37	-57	-9	>25	
Moorhen	36	WBS/WBBS waterways	-30	-45	-4	>25	
Dipper	36	WBS/WBBS waterways	-30	-47	-8	>25	

Six species are included here for which the WBS/WBBS trend is not the headline one and so is not listed in Tables A2 or A3. These species are discussed briefly below. The trends for [Yellow Wagtail](#), [Reed Bunting](#) and [Lapwing](#) are consistent in direction with the 44-year trends reported from CBC/BBS, but in each case the declines on waterways have been more severe. The [Pied Wagtail](#) declines along waterways, which are significant in all the periods assessed, are intriguing because they contrast markedly with the fluctuating but generally upward trend as measured by CBC/BBS.

For [Sedge Warbler](#), the headline trend for the UK is a non-significant 44-year shallow decline, from CBC/BBS. Large fluctuations make trends difficult to determine in this species, but the WBS/WBBS data add firmer evidence for a long-term moderate decrease.

[Moorhen](#) numbers have dipped sharply by all measures in recent seasons, perhaps through extra mortality in cold winters, and its long-term change has tipped marginally over the alert threshold.

A full set of alerts raised by WBS/WBBS, and long-term increases detected by that index, are tabulated in WBS/WBBS alerts and population increases.

Declines on CES plots

The Constant Effort Sites Scheme provides trends from standardised ringing in scrub and wetland habitats. It is possibly our best scheme for monitoring some bird populations inhabiting reed beds but its main objective is to collect integrated data on relative abundance, productivity and survival for a suite of species. The longest trends currently available from the CES cover a period of 27 years (Table A5).

Table A5 Population declines of greater than 25% recorded by the Constant Effort Sites scheme between 1984 and 2011

Species	Period (yrs)	Source	Change (%)	Lower limit	Upper limit	Alert	Comment
Willow Warbler	27	CES adults	-66	-73	-57	>50	Small sample
Willow Warbler	25	CES adults	-62	-68	-52	>50	
Lesser Whitethroat	27	CES adults	-64	-82	-49	>50	
Lesser Whitethroat	25	CES adults	-67	-79	-55	>50	
Reed Bunting	27	CES adults	-62	-72	-48	>50	
Reed Bunting	25	CES adults	-59	-70	-44	>50	
Willow Tit	27	CES adults	-57	-85	-21	>50	
Willow Tit	25	CES adults	-63	-87	-38	>50	
Sedge Warbler	27	CES adults	-42	-59	-26	>25	
Sedge Warbler	25	CES adults	-50	-62	-38	>25	
Song Thrush	27	CES adults	-33	-47	-17	>25	
Reed Warbler	27	CES adults	-29	-45	-9	>25	

Most of the species that are declining on CES sites show broadly similar trends to those from CBC/BBS or WBS/WBBS data. [Willow Tit](#) and [Song Thrush](#) are red listed on the strength of their long-term CBC/BBS declines (Table A1). [Willow Warbler](#) and [Reed Bunting](#) are similarly amber listed.

For reasons unknown, CES trends for [Lesser Whitethroat](#), [Reed Bunting](#), [Sedge Warbler](#), [Song Thrush](#) and [Reed Warbler](#) are considerably more negative than those from census data. Both CBC/BBS and WBS/WBBS show strong increases for [Reed Warbler](#), in stark contrast to the CES data presented here.

A full set of alerts raised by CES, and long-term increases detected by that scheme, are tabulated in CES alerts and population increases.

Ten-year trends and evidence of species recovery

If the status of species that have shown long-term declines were now improving, we would expect to find trends to be more positive in recent years than in the earlier part of the time series. To examine this, we list in Table B1 the best change estimates over the most recent ten-year period for which we have data (2001–11 in all but two cases), for all of the declining species listed in Tables A1–A3 (previous section). For [Lesser Spotted Woodpecker](#) and for [Woodcock](#), both now too scarce for annual monitoring to continue, the ten-year period for which data are tabulated is 1989–99.

Table B1 also includes six further species which are listed red or amber in BoCC3 because of recent breeding decline, and for which we can report ten-year trends, but which lacked monitoring data before 1994. These are [Wood Warbler](#) and [Grasshopper Warbler](#) (both red listed), and [Whinchat](#), [Nightingale](#), [Swift](#) and [Red Grouse](#) (all amber listed).

Table B1 Ten-year trends for species that have shown long-term declines

Species	Period (yrs)	Source	Change (%)	Lower limit	Upper limit	Alert	Comment
Turtle Dove	10	CBC/BBS UK	-77	-81	-72	>50	Small sample
Willow Tit	10	CBC/BBS UK	-64	-74	-57	>50	
Lesser Spotted Woodpecker	10	CBC to 1999	-51	-75	-22	>50	
Little Owl	10	CBC/BBS UK	-50	-59	-38	>50	
Snipe	10	WBS/WBBS waterways	-50	-71	-20	>25	
Starling	10	CBC/BBS England	-46	-50	-43	>25	
Cuckoo	10	CBC/BBS England	-44	-49	-40	>25	
Whinchat	10	BBS UK	-44	-55	-29	>25	Small sample
Grey Wagtail	10	WBS/WBBS waterways	-41	-49	-33	>25	
Woodcock	10	CBC to 1999	-40	-62	-11	>25	
Wood Warbler	10	BBS UK	-40	-57	-18	>25	
Redshank	10	WBS/WBBS waterways	-39	-56	-20	>25	
Mistle Thrush	10	CBC/BBS UK	-36	-39	-31	>25	
Spotted Flycatcher	10	CBC/BBS UK	-35	-49	-21	>25	
Lapwing	10	CBC/BBS UK	-34	-39	-26	>25	
Nightingale	10	BBS England	-32	-47	7		
Grey Partridge	10	CBC/BBS UK	-31	-42	-23	>25	
Marsh Tit	10	CBC/BBS UK	-28	-39	-15	>25	
Swift	10	BBS UK	-28	-35	-19	>25	
Tree Pipit	10	CBC/BBS England	-28	-43	-9	>25	
Yellow Wagtail	10	CBC/BBS UK	-25	-34	-13	>25	
Curlew	10	CBC/BBS England	-23	-29	-17		
Common Sandpiper	10	WBS/WBBS waterways	-22	-32	-9		
Dipper	10	WBS/WBBS waterways	-18	-28	-2		
House Martin	10	CBC/BBS England	-18	-24	-11		
Tawny Owl	10	CBC/BBS UK	-14	-29	2		
Meadow Pipit	10	CBC/BBS England	-12	-20	-4		
Skylark	10	CBC/BBS England	-10	-13	-5		
Willow Warbler	10	CBC/BBS England	-10	-17	-3		
Corn Bunting	10	CBC/BBS UK	-9	-25	7		
Linnet	10	CBC/BBS England	-9	-13	-1		
Song Thrush	10	CBC/BBS UK	-9	-13	-6		
Red Grouse	10	BBS UK	-8	-22	7		
Little Grebe	10	WBS/WBBS waterways	-6	-35	55		
Yellowhammer	10	CBC/BBS UK	-4	-10	2		
House Sparrow	10	CBC/BBS England	2	-2	8		
Dunnock	10	CBC/BBS UK	8	4	12		
Reed Bunting	10	CBC/BBS UK	14	4	28		
Grasshopper Warbler	10	BBS UK	25	15	70		

Whitethroat	Species	10 ^{Period} (yrs)	CBC/BBS UKSource	25 ^{Change} (%)	22 ^{Lower} limit	32 ^{Upper} limit	Alert	Comment
	<u>Bullfinch</u>	10	CBC/BBS UK	34	26	44		
	<u>Lesser Redpoll</u>	10	CBC/BBS England	41	-16	106		
	<u>Tree Sparrow</u>	10	CBC/BBS England	62	37	102		

Species are listed in ascending order of population change. Thus the species with the steepest recent decline appear first. Towards the foot of the table are species that remain in long-term decline but have shown partial recovery of those losses during the recent ten-year period.

As indicated at the top of Table B1, there is high confidence that the populations of [Turtle Dove](#), [Willow Tit](#) and [Little Owl](#) have halved within just the last ten years, or even a shorter period. These are the only species in long-term decline that suffered a 50% fall during 2001–11, but [Lesser Spotted Woodpecker](#) also met this criterion during the most recent ten-year period for which data are available. A further 16 species also continue to raise alerts, having declined significantly by more than 25% (but less than 50%) in their most recent ten-year period. All these declines compound earlier losses for these species. The ongoing declines of so many of the species listed in Table B1 raise serious conservation concern.

The 25% threshold, which is used to define decreases over the 25-year period that are worthy of amber listing, is equivalent to a change of 11% (10.9%) over ten years, assuming a constant rate of change. Thus a decrease of 11% or greater listed in Table B1 indicates that these species (27 in all, including non-significant declines for [Nightingale](#) and [Tawny Owl](#)) are on course for red or amber listing. A smaller decrease, or an increase, indicates that the population decline may be easing off. Species that have declined in the longer term but with losses smaller than 11%, or with no significant population change, over the ten-year period are [Skylark](#), [Willow Warbler](#), [Corn Bunting](#), [Linnet](#), [Song Thrush](#), [Red Grouse](#), [Little Grebe](#), [Yellowhammer](#), [House Sparrow](#) and [Lesser Redpoll](#).

Six species at the foot of the table show significant gains in population over the last ten years. Despite its recent increase, the long-term decline of [Whitethroat](#) was recognised in 2009 by the move of the species from the green to the amber list. [Whitethroat](#) numbers have increased steadily since the mid 1980s but are still far below the level prior to their population crash in 1968/69. [Tree Sparrow](#) and [Grasshopper Warbler](#) remain on the red list, and [Dunnoch](#) and [Bullfinch](#) on the amber list, because their recent increases also represent only a small recovery from earlier losses. The strong increase in [Tree Sparrow](#) numbers is very welcome but the upturn is coming from such a low level that numbers remain far below those of the mid 1970s, with the population trend graph still showing little sign of a clear recovery. Because of its recent increase, however, [Reed Bunting](#) was moved in 2009 from the red to the amber list.

Increasing species

Population changes of species for which our best long-term trend estimate from CBC/BBS (usually over 44 years) or from WBS/WBBS (a maximum of 36 years) shows an increase of more than 50% are shown in Table C1. There are 30 species listed, exactly as last year. Twenty-one of the species have more than doubled their population size over the periods given.

Table C1 Long-term population increases of greater than 50% from CBC/BBS (1967-2011) or WBS/WBBS (1975-2011), using the best survey for each species

Species	Period (yrs)	Source	Change (%)	Lower limit	Upper limit	Alert	Comment
Buzzard	44	CBC/BBS England	712	433	1829		
Greylag Goose	18	WBS/WBBS waterways	486	158	1301		
Great Spotted Woodpecker	44	CBC/BBS UK	405	265	630		
Collared Dove	39	CBC/BBS UK	372	201	552		
Shelduck	31	CBC to 1999	300	94	787		Small sample
Blackcap	44	CBC/BBS UK	272	218	398		
Canada Goose	30	WBS/WBBS waterways	262	59	1136		
Nuthatch	44	CBC/BBS UK	251	162	365		
Mute Swan	44	CBC/BBS UK	240	47	642		
Green Woodpecker	44	CBC/BBS England	199	119	335		
Stock Dove	44	CBC/BBS England	187	102	307		
Coot	44	CBC/BBS UK	182	75	550		
Woodpigeon	44	CBC/BBS UK	173	55	490		
Mallard	44	CBC/BBS UK	167	106	249		
Sparrowhawk	36	CBC/BBS England	153	49	312		
Carrion Crow	44	CBC/BBS England	125	82	186		
Jackdaw	44	CBC/BBS UK	123	38	235		
Long-tailed Tit	44	CBC/BBS England	122	63	240		
Great Tit	44	CBC/BBS UK	110	82	140		
Reed Warbler	44	CBC/BBS UK	109	24	333		
Magpie	44	CBC/BBS UK	103	68	148		
Chiffchaff	44	CBC/BBS UK	91	56	140		
Goldfinch	44	CBC/BBS England	91	41	139		
Pheasant	44	CBC/BBS England	90	53	154		
Goosander	30	WBS/WBBS waterways	81	12	189		
Coal Tit	44	CBC/BBS UK	79	1	188		
Tufted Duck	36	WBS/WBBS waterways	71	-29	275		
Oystercatcher	36	WBS/WBBS waterways	62	26	158		
Wren	44	CBC/BBS UK	54	35	76		
Pied Wagtail	44	CBC/BBS UK	50	9	107		

Four of the fastest-increasing species in this report are actually not included in Table C1, because their monitoring data cover too short a period. The population of [Ring-necked Parakeet](#) is estimated to have risen by 1057% (more than an elevenfold increase) over the 16 years 1995–2011. Arguably, however, this is more a conservation problem than a success! Unmitigated successes are the growth during 1995–2011, estimated through BBS, of the reintroduced [Red Kite](#) (+676%) and of [Barn Owl](#) (+279%). Though the trajectory has been moderated considerably by recent cold-weather-related setbacks, attention should also be drawn to the rapid rise of [Cetti's Warbler](#), a newly established native species, which CES now estimates to have increased by 73% over the recent ten-year period.

Four groups stand out among the increasing species: corvids – [Carrion Crow](#), [Magpie](#) and [Jackdaw](#); doves – [Collared Dove](#), [Stock Dove](#) and [Woodpigeon](#); woodpeckers and other smaller species of woodland and gardens; and some waterbirds. Corvids appear to have benefited from gamebird management practices in recent years, and the larger doves from the increased acreage of brassica crops (particularly oilseed rape).

The majority of the third group are species primarily of woodland that are also common in gardens in some areas [Great Spotted Woodpecker](#), [Green Woodpecker](#), [Nuthatch](#), [Blackcap](#), [Great Tit](#), [Wren](#), [Long-tailed Tit](#) and [Coal Tit](#). The reasons for these increases are presently unclear. [Pied Wagtail](#) has increased in numbers by 50% on CBC/BBS plots over 44 years, but declined by 67% on WBS/WBBS plots over the past 36 years. The former index is likely to be more representative of the UK population as a whole. [Reed Warbler](#), also an insectivore, has been expanding its range northwards and westwards and might be benefiting from climate change.

A number of species associated with freshwater habitats are becoming more abundant, although differences between their ecological requirements make it unlikely that the major causal factors are common to all. For [Mallard](#), the CBC/BBS increase was matched by a WBS/WBBS increase of 214% over 36 years. The long-term increases recorded for [Mute Swan](#) on both CBC/BBS and WBS/WBBS plots are likely to be the result of banning the use of lead weights by anglers, which took effect in 1986. [Greylag Goose](#), [Shelduck](#), [Canada Goose](#), [Tufted Duck](#), [Coot](#) and [Goosander](#) are other wildfowl among this report's increasing species. [Oystercatchers](#) have increased by 72% on WBS/WBBS plots over the last 36 years. This finding is consistent with the results of the most recent survey of *Breeding Waders of Wet Meadows* which found that numbers of [Oystercatchers](#) using these habitats in England and Wales increased by 51% between 1982 and 2002 (Wilson *et al.* 2005).

Two widespread raptors have shown remarkable recoveries from low population levels after the banning of certain poisonous farmland pesticides in the early 1960s, assisted by lower levels of illegal predator control by shooting interests. [Buzzards](#) increased by a remarkable 712% between 1967 and 2011, with a rapid increase of 83% over the last ten years alone. [Sparrowhawks](#), too scarce for CBC to monitor until the mid 1970s, showed a 153% increase over the 36-year period from 1975 to 2011. However, their recovery appears to have been completed earlier than [Buzzard's](#), with the population currently stable or in shallow decline.

While [Pheasant](#) holds a place in this table, its increase in census data has been driven largely by the hugely increasing scale of releases of artificially reared poults for shooting, from which the corvids may also have benefited.

Changes in breeding performance

Changes in a range of aspects of breeding performance can be measured under the Nest Record Scheme (NRS) and the Constant Effort Sites (CES) scheme. The NRS provides information on components of breeding performance (clutch size, brood size and failure rates at the egg and nestling stages) that can be combined to give an overall estimate of productivity per nesting attempt (FPBA) – see NRS page for further information. The CES scheme provides an index of breeding performance accrued over all nesting attempts in a particular year. CES results also take into account any changes in the survival rates of fledglings in the first few months after leaving the nest, a period when losses of young can be high.

Breeding performance may be influenced by a variety of factors, including food availability, predation pressure and weather conditions. Variation in breeding performance may help to influence fluctuations in abundance and may even be the main demographic factor responsible for determining the size of the population. Conversely, the breeding performance of a population may be inversely related to its size, with productivity decreasing as the number of individuals increases, and vice versa. This relationship may be due to the action of density-dependent factors, such as competition for resources: as numbers increase, competition for resources is likely to increase, possibly resulting in poorer productivity. Alternatively, increases in abundance may be accompanied by range expansion into new, suboptimal habitats where breeding performance is poorer, thus reducing the average productivity of the population. The converse is also true, and where declines result from the loss of individuals from these suboptimal habitats, there may be a subsequent increase in average productivity.

Changes in Fledglings Per Breeding Attempt from Nest Record Scheme data

The NRS started collating nest histories of individual breeding attempts in 1939 and sufficient data are available for trends to be produced from the mid 1960s onwards. Previous reports have explored annual variation in clutch size, brood size and stage-specific nest failure rates, and these breeding parameters are included in the Summary tables. While detailed exploration of annual variation in productivity is essential if the impacts of environmental factors on breeding success are to be fully understood, the combined effects of concurrent changes in the number of offspring and failure rates can be difficult to interpret. These measures are therefore integrated into a single annual figure representing the mean number of young leaving each nest, termed Fledglings Per Breeding Attempt (FPBA; Siriwardena *et al.* 2000b, Crick *et al.* 2003).

All species displaying significant temporal trends in mean FPBA are included in Table D1. In total, 40 species exhibited significant trends in FPBA over the past 43 years, of which nine were negative, indicating that reproductive output has decreased over time. Birds exhibiting declines in productivity include three BoCC red-listed species ([Nightjar](#), [Tree Pipit](#) and [Linnet](#)), two amber-listed species ([Willow Warbler](#) and [Bullfinch](#)) and four green-listed species ([Moorhen](#), [Coal Tit](#), [Chaffinch](#) and [Greenfinch](#)). While productivity of [Moorhen](#), [Nightjar](#), [Willow Warbler](#) and [Linnet](#) has been falling consistently, trends for the other five species are curvilinear, increasing up to the mid 1980s and decreasing thereafter.

Table D1 Significant trends in fledglings per breeding attempt measured between 1968 and 2011

Species	Period (yrs)	Mean annual sample	Trend	Predicted in first year	Predicted in last year	Change	Comment
Nightjar	43	21	Linear decline	1.42 fledglings	0.71 fledglings	-0.71 fledglings	Small sample
Coal Tit	43	54	Curvilinear	6.7 fledglings	6.26 fledglings	-0.44 fledglings	
Willow Warbler	43	69	Linear decline	3.59 fledglings	3.16 fledglings	-0.43 fledglings	
Moorhen	43	52	Linear decline	2.51 fledglings	2.13 fledglings	-0.38 fledglings	
Linnet	43	122	Linear decline	2.69 fledglings	2.36 fledglings	-0.33 fledglings	
Tree Pipit	43	13	Curvilinear	1.54 fledglings	1.24 fledglings	-0.3 fledglings	Small sample
Chaffinch	43	129	Curvilinear	1.6 fledglings	1.38 fledglings	-0.22 fledglings	
Greenfinch	43	91	Curvilinear	2.14 fledglings	2.02 fledglings	-0.12 fledglings	
Bullfinch	43	33	Curvilinear	1.49 fledglings	1.37 fledglings	-0.12 fledglings	
Blackbird	43	233	Curvilinear	1.47 fledglings	1.54 fledglings	0.07 fledglings	
Duncock	43	118	Curvilinear	1.68 fledglings	1.76 fledglings	0.08 fledglings	
Collared Dove	43	54	Curvilinear	0.81 fledglings	0.91 fledglings	0.1 fledglings	
Trecreeper	43	20	Curvilinear	2.69 fledglings	2.81 fledglings	0.12 fledglings	Small sample
House Sparrow	43	103	Curvilinear	2.3 fledglings	2.47 fledglings	0.17 fledglings	
Woodpigeon	43	77	Curvilinear	0.55 fledglings	0.74 fledglings	0.19 fledglings	
Meadow Pipit	43	46	Curvilinear	2.07 fledglings	2.32 fledglings	0.25 fledglings	
Yellowhammer	43	49	Curvilinear	0.82 fledglings	1.16 fledglings	0.34 fledglings	
Stock Dove	43	72	Linear increase	0.98 fledglings	1.4 fledglings	0.42 fledglings	
Robin	43	207	Linear increase	2.42 fledglings	2.85 fledglings	0.43 fledglings	
Wren	43	96	Curvilinear	2.37 fledglings	2.86 fledglings	0.49 fledglings	
Peregrine	43	22	Linear increase	1.77 fledglings	2.27 fledglings	0.5 fledglings	Small sample
Reed Warbler	43	141	Linear increase	2.34 fledglings	2.84 fledglings	0.5 fledglings	
Sparrowhawk	43	32	Curvilinear	2.64 fledglings	3.15 fledglings	0.51 fledglings	
Skylark	43	42	Linear increase	1.06 fledglings	1.57 fledglings	0.51 fledglings	
Buzzard	43	28	Linear increase	1.51 fledglings	2.07 fledglings	0.56 fledglings	Small sample
Tawny Owl	43	68	Linear increase	1.39 fledglings	1.97 fledglings	0.58 fledglings	Nocturnal species
Pied Wagtail	43	88	Linear increase	2.99 fledglings	3.58 fledglings	0.59 fledglings	
Carion Crow	43	40	Curvilinear	1.73 fledglings	2.35 fledglings	0.62 fledglings	Includes Hooded Crow
Kestrel	43	41	Curvilinear	2.92 fledglings	3.6 fledglings	0.68 fledglings	
Starling	43	120	Linear increase	2.6 fledglings	3.32 fledglings	0.72 fledglings	
Grey Wagtail	43	54	Linear increase	2.6 fledglings	3.37 fledglings	0.77 fledglings	
Jackdaw	43	61	Linear increase	1.73 fledglings	2.53 fledglings	0.8 fledglings	
Barn Owl	43	30	Linear increase	2.33 fledglings	3.25 fledglings	0.92 fledglings	Small sample

Merlin	43	21	Linear increase	2.44 fledglings	3.38 fledglings	0.94 fledglings	Small sample
Dipper	43	82	Curvilinear	2.05 fledglings	3.06 fledglings	1.01 fledglings	
Tree Sparrow	43	253	Linear increase	2.74 fledglings	3.87 fledglings	1.13 fledglings	
Wheatear	43	16	Curvilinear	3.84 fledglings	4.99 fledglings	1.15 fledglings	Small sample
Magpie	43	43	Curvilinear	1.14 fledglings	2.36 fledglings	1.22 fledglings	
Redstart	43	54	Curvilinear	3.52 fledglings	5.01 fledglings	1.49 fledglings	
Nuthatch	43	54	Linear increase	3.71 fledglings	5.28 fledglings	1.57 fledglings	

See Key to species texts for help with interpretation

There is increasing evidence that organisms at lower trophic levels are responding to climatic change more rapidly than those towards the top of the food chain (Visser & Both 2005, Thackeray *et al.* 2010). Resulting mismatches in the timing of food availability and of offspring food demand, referred to as phenological disjunction, can have severe impacts on breeding success and ultimately on population trends of bird species (Both *et al.* 2009), although there is evidence that the magnitude of these impacts may vary with diet and breeding habitat (Dunn & Møller 2013) and that reduced productivity may be buffered by density dependent increase in survival in some species, including Reed *et al.* 2012, 2013). Long-distance migrants are thought to be particularly susceptible, due to their later arrival on the breeding grounds and the energetic demands of their journey northwards, which may constrain their ability to advance their laying dates (Rubolini *et al.* 2010, Ockendon *et al.* 2012, but see Goodenough *et al.* 2011); this mechanism could therefore contribute to the increasing nestling failure rates underpinning the productivity declines detected for Shortall *et al.* 2009), particularly moths (Conrad *et al.* 2006, Fox 2013), have been reported across the UK and these may also impact on the productivity of nesting attempts of all three species, increasing the incidence of whole brood failure due to starvation or desertion by under-nourished parents. Trans-Saharan migrants may also be experiencing negative impacts of climate change in their African wintering grounds, where reduced rainfall could lead to a fall in insect abundance and a subsequent loss of condition, resulting in a lower reproductive output in the following spring (Saino *et al.* 2004, 2011, Schaub *et al.* 2011, Ockendon *et al.* 2013).

Woodland passerines that depend on short-lived peaks in the availability of larval Lepidoptera to provide food for their nestlings may also suffer reduced productivity as a result of climate-induced changes in phenology. As springs have become warmer, oak leafing dates have advanced, a shift matched by caterpillars (Buse *et al.* 1999) but not by tits (Visser *et al.* 1998) or flycatchers (Both *et al.* 2009). A recent study in the Netherlands found that responses to disjunction may vary spatially, with the negative effects exacerbated in more seasonal habitats, where the window of prey availability is smaller (Both *et al.* 2010), and regional variation in breeding success at sites across the UK is currently being investigated. While the figures presented in this report indicate that [Blue Tit](#) and [Great Tit](#) brood sizes have fallen and that nestling stage failure rates of both these tit species and [Pied Flycatcher](#) have risen, as would be predicted under a mismatch scenario, FPBA trends are not significant due to a concurrent drop in egg-stage failure rates. However, FPBA of [Chaffinch](#), another woodland insectivore heavily reliant on moth larvae to provision its offspring, has decreased significantly, as has that of [Coal Tit](#); again, increasing nestling failure rates have contributed to these declines in productivity.

Increasing egg-stage failure rates are the main driver of the drop in Groom 1993, Stoate & Szczur 2001, 2006), previous studies have failed to find any evidence of a significant impact at a national scale (Gooch *et al.* 1991, Thomson *et al.* 1998, Chamberlain *et al.* 2009, Newson *et al.* 2009, Vögeli *et al.* 2011). However, several recent studies have suggested that predation pressure may increase in response to climatic warming. Cox *et al.* (2013) found that the incidence of nest predation by birds and snakes, but not mammals, increased with temperature in the USA, although the mechanism is unknown, while Auer & Martin (2013) demonstrated an increase in the proportion of predated nests across a range of species due to climate-induced shifts in plant–herbivore interactions.

Increased grazing pressure by deer, numbers of which are rising rapidly in many areas of the UK (Newson *et al.* 2012), has been identified as a possible driver of population declines in the UK (Fuller *et al.* 2005) and the USA (Martin *et al.* 2011), the removal of the herb and shrub layers potentially reducing the availability of both food and well-concealed nesting sites. This process may have contributed to the observed declines in productivity of both [Willow Warbler](#) and [Bullfinch](#). A recent study using BBS deer data indicated that declines in [Willow Warbler](#) were most pronounced in areas where Reeves's muntjac had increased at the fastest rate (Newson *et al.* 2012), in agreement with a previous study looking at regional variation in Morrison *et al.* 2010). While a similar negative relationship was identified for Siriwardena *et al.* 1998a, 1999, 2000b, 2001, Proffitt *et al.* 2004).

Declining food availability may also be an issue for farmland bird species displaying negative trends in FPBA. Reduced access to winter stubbles due to changes in farming practices have been linked to declines in survival rates of species such as Siriwardena *et al.* 1998b, Peach *et al.* 1999, Siriwardena *et al.* 2000b). If adults of stubble-feeding species are in poorer condition at the start of the breeding season, their investment in reproduction may also be reduced, and the granivorous diet of Siriwardena *et al.* 1999, 2000b).

Increasing human activity in the countryside, resulting from a growing population, could increase disturbance levels, which could in turn influence the rates of predation and desertion. An investigation of Langston *et al.* 2007) and a recent review of impacts of recreational disturbance found breeding success to be adversely affected by human activity levels in 28 out of 33 papers cited (Steven *et al.* 2011). Further research into the impacts of nest predators on population trajectories, at a variety of spatial scales, is urgently required.

The colonisation of urban habitats by [Greenfinch](#) may also have increased the proportion of data originating from gardens, which may represent a relatively resource-poor breeding environment when compared with their more traditional farmland habitats, resulting in the smaller broods and clutch sizes observed. Similar reductions in reproductive output across an urban gradient have been observed for tit species, although results from localised studies are conflicting (see Chamberlain *et al.* 2009 for review) and more research is needed to see whether these are representative at a national scale. The recent outbreak of trichomonosis, which has significantly and rapidly reduced the abundance of Robinson *et al.* 2010b), may provide a good test of the hypothesis that productivity declines over the last 50 years represent a density-dependent response. Greenwood & Baillie 2008). [Moorhen](#) populations were relatively stable until the recent run of cold winters, and the causes of the productivity decline in this poorly studied species are currently unclear. Failure rates at both the egg and chick stage have increased substantially, possibly due to increasing numbers of mammalian predators, such as American mink (*Neovison vison*) and Eurasian otter (*Lutra lutra*), or to the competitor species [Coot](#), which is known to destroy clutches of Moorhens nesting nearby.

FPBA has increased significantly over the last 43 years for 31 species, across a wide range of taxonomic groups (Table D1). Population trends are also upward for 16 of these species, including raptors ([Sparrowhawk](#), [Buzzard](#), [Merlin](#), [Peregrine](#), [Barn Owl](#)), pigeons ([Stock Dove](#), [Woodpigeon](#), [Collared Dove](#)), corvids ([Magpie](#), [Jackdaw](#), [Carion Crow](#)), and some small passerines ([Reed Warbler](#), [Nuthatch](#), [Wren](#), [Robin](#) and [Pied Wagtail](#)). It is therefore possible that increasing productivity has contributed to the population growth exhibited by these species over recent decades. Conversely, 13 species ([Kestrel](#), [Tawny Owl](#), [Skylark](#), [Starling](#), [Dipper](#), [Blackbird](#), [Wheatear](#), [Duncock](#), [House Sparrow](#), [Tree Sparrow](#), [Grey Wagtail](#), [Meadow Pipit](#) and [Yellowhammer](#)), have declined in number as FPBA has increased, suggesting that a density-dependent reduction in intraspecific competition may have enabled breeding success to rise.

Changes in productivity from Constant Effort Sites ringing data

The CES started monitoring populations in 1983, so the changes in productivity (Table D2) cover roughly half the period of the Nest Record Scheme results. The CES data set is unique in providing relative measures of adult abundance and productivity from the same set of sites in wetland and scrub habitats. While the NRS data set monitors the productivity of individual nesting attempts, the proportion of juveniles in the CES catch provides a relative measure of annual variation in productivity that

integrates the effects of the number of fledglings produced per attempt, number of nesting attempts and immediate post-fledging survival. Use of these two techniques in combination provides a powerful method of determining which factors are responsible for observed declines in recruitment of young birds into the breeding population.

Table D2 Changes in productivity indices (percentage juveniles) for CES, 1984-2011, calculated from smoothed trend

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Comment
Goldfinch	27	36	-68	-88	-21	
Garden Warbler	27	78	-62	-76	-36	
Sedge Warbler	27	72	-60	-75	-33	
Reed Bunting	27	62	-56	-76	-13	
Blue Tit	27	103	-55	-67	-38	
Blackcap	27	98	-43	-54	-22	
Blackbird	27	101	-37	-53	-20	
Great Tit	27	102	-34	-54	-8	
Song Thrush	27	90	-30	-48	-2	
Willow Warbler	27	99	-24	-43	-0	
Reed Warbler	27	62	29	1	99	
Chaffinch	27	84	93	3	229	

See Key to species texts for help with interpretation

Overall, ten species exhibit significant declines in the proportion of juveniles captured (Table D2). The apparent productivity of [Blue Tit](#), [Garden Warbler](#), [Sedge Warbler](#), [Goldfinch](#) and [Reed Bunting](#) has fallen by more than 50% over the last 25 years, while [Great Tit](#), [Willow Warbler](#), [Blackcap](#), [Blackbird](#) and [Song Thrush](#) show reductions in relative productivity of between 25% and 50%.

Although three of these species, [Peach](#) *et al.* 1991, 1995a, 1999, [Robinson](#) *et al.* 2004, 2010, [Baillie](#) *et al.* 2009). [Peach](#) *et al.* 1999). NRS data demonstrate a similar decline in [Morrison](#) *et al.* 2010).

For species such as [Blue Tit](#), [Great Tit](#), [Blackcap](#) and [Goldfinch](#), where population increase has occurred, reductions in productivity may be driven by density-dependent processes, whereby increased competition for resources in an expanding population reduces the mean breeding success per pair. [Garden Warbler](#) populations have fluctuated but remained stable over the period during which CES has operated.

Two species, [Chaffinch](#) and [Reed Warbler](#), have displayed a significant increase in productivity at CE sites. In the case of [Reed Warbler](#), the NRS dataset also indicates a significant increase in breeding success over this period. The marked difference between the [Chaffinch](#) CES trend and the decline in productivity identified by the NRS data set requires further investigation, but it may be that changes in post-juvenile survival over time are responsible.

Changes in average laying dates from Nest Record Scheme data

Over the past 25 years, many species have exhibited a trend towards progressively earlier clutch initiation ([Crick](#) *et al.* 1997) with laying dates showing curvilinear responses over the past 50 years as spring temperatures have cooled and then warmed ([Crick](#) & [Sparks](#) 1999). Table D3 confirms that, since the mid 1960s, the majority of species exhibiting significant trends show an advancement of laying dates rather than a delay. Thus 42 species are laying between one and 31 days earlier, on average, than they were 43 years ago. It is interesting to note that, while the results of previous studies predict laying-date advancement to be more constrained in long-distance migrants (Both *et al.* 2009, [Rubolini](#) *et al.* 2010), the magnitude of the laying-date shift in both [Pied Flycatcher](#) and [Redstart](#) (12 days and 14 days respectively), is greater than that displayed by many resident species. However, the mean laying date of the migrant species is still approximately a fortnight later than that of common residents such as [Blue Tit](#) and [Great Tit](#). No taxonomic or ecological associations are apparent and a wide range of species demonstrate trends of a similar magnitude ([Crick](#) *et al.* 1997).

Table D3 Significant trends in laying date measured between 1968 and 2011

Species	Period (yrs)	Mean annual sample	Trend	Predicted in first year	Predicted in last year	Change	Comment
Magpie	43	33	Linear decline	Apr 24	Mar 24	-31 days	
Greenfinch	43	91	Linear decline	May 25	May 8	-17 days	
Long-tailed Tit	43	53	Linear decline	Apr 21	Apr 5	-16 days	
Goldfinch	43	24	Curvilinear	Jun 5	May 21	-15 days	Small sample
Redstart	43	64	Curvilinear	May 21	May 7	-14 days	
Chiffchaff	43	58	Linear decline	May 16	May 2	-14 days	
Coal Tit	43	45	Linear decline	May 3	Apr 19	-14 days	
Corn Bunting	43	17	Linear decline	Jun 27	Jun 13	-14 days	Small sample
Blackcap	43	42	Curvilinear	May 20	May 7	-13 days	

Dipper	43	67	Linear decline	Apr 19	Apr 7	-12 days	
Pied Flycatcher	43	441	Linear decline	May 21	May 9	-12 days	
Species	Period (yrs)	Mean annual sample	Trend	Predicted in first year	Predicted in last year	Change	Comment
Nuthatch	43	31	Linear decline	May 2	Apr 20	-12 days	
Tree Creeper	43	13	Linear decline	May 7	Apr 25	-12 days	Small sample
Peregrine	43	10	Linear decline	Apr 14	Apr 3	-11 days	Small sample
Reed Warbler	43	208	Curvilinear	Jun 17	Jun 6	-11 days	
Marsh Tit	43	14	Linear decline	Apr 28	Apr 17	-11 days	Small sample
Chaffinch	43	122	Linear decline	May 12	May 1	-11 days	
Stonechat	43	42	Curvilinear	May 3	Apr 23	-10 days	
Whitethroat	43	20	Curvilinear	May 27	May 17	-10 days	Small sample
Great Tit	43	379	Curvilinear	May 1	Apr 21	-10 days	
Carion Crow	43	30	Linear decline	Apr 18	Apr 8	-10 days	Includes Hooded Crow
Kestrel	43	23	Linear decline	May 5	Apr 26	-9 days	Small sample
Sedge Warbler	43	47	Curvilinear	May 29	May 20	-9 days	
Garden Warbler	43	22	Linear decline	May 28	May 19	-9 days	Small sample
Willow Warbler	43	88	Curvilinear	May 19	May 10	-9 days	
Blue Tit	43	519	Curvilinear	Apr 30	Apr 21	-9 days	
House Sparrow	43	64	Linear decline	May 25	May 16	-9 days	
Swallow	43	207	Curvilinear	Jun 19	Jun 11	-8 days	
Tree Pipit	43	20	Curvilinear	May 27	May 19	-8 days	Small sample
Grey Wagtail	43	60	Curvilinear	May 5	Apr 27	-8 days	
Robin	43	149	Linear decline	Apr 28	Apr 20	-8 days	
Ring Ouzel	43	22	Linear decline	May 15	May 7	-8 days	Small sample
Jackdaw	43	30	Curvilinear	Apr 24	Apr 16	-8 days	
Wood Warbler	43	35	Curvilinear	May 24	May 17	-7 days	
Starling	43	86	Curvilinear	Apr 27	Apr 20	-7 days	
Moorhen	43	83	Linear decline	May 10	May 4	-6 days	
Wren	43	87	Linear decline	May 14	May 8	-6 days	
Oystercatcher	43	65	Curvilinear	May 18	May 13	-5 days	
Whinchat	43	29	Linear decline	May 30	May 25	-5 days	Small sample
Tree Sparrow	43	281	Linear decline	May 28	May 23	-5 days	
Pied Wagtail	43	88	Curvilinear	May 18	May 16	-2 days	
Meadow Pipit	43	39	Curvilinear	May 19	May 18	-1 days	
Blackbird	43	231	Curvilinear	Apr 23	Apr 25	2 days	
Skylark	43	19	Curvilinear	May 24	May 30	6 days	Small sample
Bullfinch	43	34	Linear increase	May 26	Jun 1	6 days	
Yellowhammer	43	26	Linear increase	May 31	Jun 7	7 days	Small sample
Woodpigeon	43	87	Linear increase	Jun 1	Jun 23	22 days	

See Key to species texts for help with interpretation

The significance of the changes in phenology for breeding performance is poorly understood but has stimulated a large number of scientific studies, including several ongoing projects at BTO. Earlier average laying may be beneficial for birds because earlier fledging is often related to improved survival to the following year – thus early-nesting parents have an increased chance of having their offspring recruited into the next generation (Visser *et al.* 1998). However, the timing of leaf emergence and the speed of caterpillar development is also changing under increased temperatures (Buse *et al.* 1999, Visser & Holleman 2001) and the results of several recent studies have suggested that some birds may be unable to advance their breeding sufficiently to match phenological changes in their food supply, such that later-nesting birds are suffering from poorer productivity. Both *et al.* (2006) demonstrated that mismatches between periods of food availability and chick demand can affect abundance in Dutch [Pied Flycatcher](#) populations, with those demonstrating the largest mismatches between arrival in spring and peak caterpillar abundance exhibiting the greatest declines. As a consequence of climate change there may be an increasing mismatch between predator activities and the availability of their food supplies at different trophic levels within ecosystems (Both *et al.* 2009). Recent studies in the Netherlands have suggested that the magnitude of disjunction may be mediated by habitat type, with species in more seasonal habitats at greatest risk of negative impacts on productivity (Both *et al.* 2010). However, while Reed *et al.* 2012, 2013). Whether such compensation will persist as the climate warms further remains to be seen and the population level significance of trophic mismatches remains an active research area with potentially important policy implications for conservation.

Only five species exhibit significant trends towards later laying, all of which produce multiple broods per season. A recent collaboration between BTO and Aberdeen University, using NRS data, identified an increase in the frequency of repeat brooding in Cornulier *et al.* 2009) which, as mean laying dates are calculated across all broods, would result in the observed shift. Increased production of repeat broods could be stimulated by climatic amelioration, with later nests being more productive in warmer conditions, or by movement of birds away from farmland and into habitats where they are released from constraints on multiple brooding. Previous research into multiple brooding in Chamberlain & Siriwardena 2000), but this species may also increasingly have moved to alternative habitats. A recent study using data from North America and Europe identified a positive temporal trend in the breeding season length of multi-brooded, but not single-brooded, bird species, consistent with the hypothesis that climate change is extending the window of opportunity for nesting for species less reliant on seasonal resource peaks (Dunn & Møller 2013).

It is likely that the laying dates of the majority of those species that do not show a significant trend in timing of breeding are also related to weather, but that their weather-mediated cues do not show any trend over time (Crick & Sparks 1999).

Conclusion

This report is designed to be useful as a ready source of information for conservation practitioners, and as a source of information for those involved in more strategic conservation policy-making, as well as to the general student of bird populations. It provides a relatively simple and concise overview of the way in which populations are changing, suggesting areas where further research is required or where conservation action needs to be taken. The information presented here is a summary of a very extensive and much more detailed data set held by the BTO.

Alerts are raised as a result of declines in the population sizes of a considerable number of species. These alerts will help conservation organisations to prioritise future conservation action, alongside the Birds of Conservation Concern list (Eaton *et al.* 2009) and other information.

The demographic information contained in this report will also help conservation organisations to target their resources more effectively. For declining species of conservation importance, declines in breeding performance may indicate that conservation action should be targeted towards the breeding season; such responses may sometimes be masked, however, by density-dependent improvements in breeding success as the population declines (Green 1999). The lack of a decline in breeding performance may suggest that factors other than nesting success, such as loss of habitat or changes in survival rates are more likely to be influencing the observed population declines. A report of this kind can provide only an initial summary of such information, and a full assessment of the population dynamics of a declining species will generally require more detailed investigations (e.g. Peach *et al.* 1999, Freeman & Crick 2003, Robinson *et al.* 2004).

Finally, we hope that users of this report will provide feedback on how it can be improved. We would welcome comments on any aspect of this report, as they will help us to produce a better and more useful next edition.

Email your comments to: john.marchant@bto.org

Utilities

The tables of population change that appear on the species pages are species-based selections from a single unified table, with data newly calculated for this edition of the report. A number of additional selections from this table, by scheme and time-period, are presented in the Summary tables section. Using the [table generator](#), you can interrogate the master table by data source or time-period, for all species or for your own selection of species, and choose how your extract will be sorted.

This edition of the Bird Trends report is the latest in an annual series that began in 1997. Citations for previous editions are listed under Previous reports. Links are given to the full text of previous reports, where these are still available online.

Previous reports

Previous reports in this series are listed, from the most recent to the earliest. The first two (Cricket *et al.* 1997, 1998) were produced as paper reports, but all subsequent editions are purely web-based and url addresses must be included in their citations.

Note that www.bto.org/about-birds/birdtrends will always link to the home page of the most recent version of this report. Web addresses including a year (e.g. [.../birdtrends/2012/...](http://www.bto.org/about-birds/birdtrends/2012/)) may lead you to earlier reports in the series, now superseded.

BirdTrends 2012: trends in numbers, breeding success and survival for UK breeding birds

Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2013) *BirdTrends 2012: trends in numbers, breeding success and survival for UK breeding birds* Research Report 644. BTO, Thetford. www.bto.org/about-birds/birdtrends/2012

BirdTrends 2011: trends in numbers and demography for UK breeding birds

Baillie, S.R., Marchant, J.H., Leech, D.I., Renwick, A.R., Eglington, S.M., Joys, A.C., Noble, D.G., Barimore, C., Conway, G.J., Downie, I.S., Risely, K. & Robinson, R.A. (2012) *BirdTrends 2011: trends in numbers and demography for UK breeding birds* Research Report 609. BTO, Thetford. (www.bto.org/about-birds/birdtrends/2011)

Breeding Birds in the Wider Countryside: their conservation status 2010

Baillie, S.R., Marchant, J.H., Leech, D.I., Renwick, A.R., Joys, A.C., Noble, D.G., Barimore, C., Conway, G.J., Downie, I.S., Risely, K. & Robinson, R.A. (2010) *Breeding Birds in the Wider Countryside: their conservation status 2010*. Research Report 565. BTO, Thetford. (www.bto.org/birdtrends2010/index.htm)

Breeding Birds in the Wider Countryside: their conservation status 2009

Baillie, S.R., Marchant, J.H., Leech, D.I., Joys, A.C., Noble, D.G., Barimore, C., Downie, I.S., Grantham, M.J., Risely, K. & Robinson, R.A. (2010) *Breeding Birds in the Wider Countryside: their conservation status 2009*. Research Report 541. BTO, Thetford. (www.bto.org/birdtrends2009/index.htm)

Breeding Birds in the Wider Countryside: their conservation status 2008

Baillie, S.R., Marchant, J.H., Leech, D.I., Joys, A.C., Noble, D.G., Barimore, C., Downie, I.S., Grantham, M.J., Risely, K. & Robinson, R.A. (2009) *Breeding Birds in the Wider Countryside: their conservation status 2008*. Research Report 516. BTO, Thetford. (www.bto.org/birdtrends2008/index.htm)

Breeding Birds in the Wider Countryside: their conservation status 2007

Baillie, S.R., Marchant, J.H., Crick, H.Q.P., Noble, D.G., Balmer, D.E., Barimore, C., Coombes, R.H., Downie, I.S., Freeman, S.N., Joys, A.C., Leech, D.I., Raven, M.J., Robinson, R.A. & Thewlis, R.M. (2007) *Breeding Birds in the Wider Countryside: their conservation status 2007*. Research Report 487. BTO, Thetford. (www.bto.org/birdtrends2007/index.htm)

Breeding Birds in the Wider Countryside: their conservation status 2006

Baillie, S.R., Marchant, J.H., Crick, H.Q.P., Noble, D.G., Balmer, D.E., Barimore, C., Coombes, R.H., Downie, I.S., Freeman, S.N., Joys, A.C., Leech, D.I., Raven, M.J., Robinson, R.A. & Thewlis, R.M. (2007) *Breeding Birds in the Wider Countryside: their conservation status 2006* Research Report 470. BTO, Thetford. (www.bto.org/birdtrends2006/index.htm)

Breeding Birds in the Wider Countryside: their conservation status 2005

Baillie, S.R., Marchant, J.H., Crick, H.Q.P., Noble, D.G., Balmer, D.E., Coombes, R.H., Downie, I.S., Freeman, S.N., Joys, A.C., Leech, D.I., Raven, M.J., Robinson, R.A. & Thewlis, R.M. (2006) *Breeding Birds in the Wider Countryside: their conservation status 2005*. Research Report 435. BTO, Thetford. (www.bto.org/birdtrends2005/index.htm)

Breeding Birds in the Wider Countryside: their conservation status 2004

Baillie, S.R., Marchant, J.H., Crick, H.Q.P., Noble, D.G., Balmer, D.E., Beaven, L.P., Coombes, R.H., Downie, I.S., Freeman, S.N., Joys, A.C., Leech, D.I., Raven, M.J., Robinson, R.A. & Thewlis, R.M. (2005) *Breeding Birds in the Wider Countryside: their conservation status 2004* Research Report 385. BTO, Thetford. (www.bto.org/birdtrends2004/index.htm)

Breeding Birds in the Wider Countryside: their conservation status 2003

Crick, H.Q.P., Marchant, J.H., Noble, D.G., Baillie, S.R., Balmer, D.E., Beaven, L.P., Coombes, R.H., Downie, I.S., Freeman, S.N., Joys, A.C., Leech, D.I., Raven, M.J., Robinson, R.A. & Thewlis, R.M. (2004) *Breeding Birds in the Wider Countryside: their conservation status 2003* Research Report 353. BTO, Thetford. (www.bto.org/birdtrends2003/index.htm)

Breeding Birds in the Wider Countryside: their conservation status 2001

Baillie, S.R., Crick, H.Q.P., Balmer, D.E., Beaven, L.P., Downie, I.S., Freeman, S.N., Leech, D.I., Marchant, J.H., Noble, D.G., Raven, M.J., Simpkin, A.P., Thewlis, R.M. & Wernham, C.V. (2002) *Breeding Birds in the Wider Countryside: their conservation status 2001* Research Report 278. BTO, Thetford. (www.bto.org/birdtrends2001/index.htm)

Breeding Birds in the Wider Countryside: their conservation status 2000

Baillie, S.R., Crick, H.Q.P., Balmer, D.E., Bashford, R.I., Beaven, L.P., Freeman, S.N., Marchant, J.H., Noble, D.G., Raven, M.J., Siriwardena, G.M., Thewlis, R. & Wernham, C.V. (2001) *Breeding Birds in the Wider Countryside: their conservation status 2000*. Research Report 252. BTO, Thetford. (www.bto.org/birdtrends2000/index.htm)

Breeding Birds in the Wider Countryside: their conservation status 1972–1996

Crick, H.Q.P., Baillie, S.R., Balmer, D.E., Bashford, R.I., Beaven, L.P., Dudley, C., Glue, D.E., Gregory, R.D., Marchant, J.H., Peach, W.J. & Wilson, A.M. (1998) *Breeding Birds in the Wider Countryside: their conservation status (1972–1996)*. Research Report 198. BTO, Thetford.

Breeding Birds in the Wider Countryside: their conservation status 1971–1995

Crick, H.Q.P., Baillie, S.R., Balmer, D.E., Bashford, R.I., Dudley, C., Glue, D.E., Gregory, R.D., Marchant, J.H., Peach, W.J. & Wilson, A.M. (1997) *Breeding Birds in the Wider Countryside: their conservation status (1971–1995)*. Research Report 187. BTO, Thetford.

References

Clicking on reference links within the text of this report will bring you to its full details in this section: the reference sought will be at the very top of your view.

In some cases, we provide an onward link either to an abstract or, where it is freely available, to the full text of the listed publication. Alternatively, your own web search will often take you to the summary of an article and the opportunity to purchase the text in full. The [doi](#) (digital object identifier), where given, is a useful key to copy to a search engine.

Most of the listed publications are available in printed form to BTO members and other bona fide researchers through the Chris Mead Library at BTO headquarters in Thetford.

Aebischer, N.J. (1999) Multi-way comparisons and generalised linear models of nest success: extensions of the Mayfield method *Bird Study* 46: S22–S31.

Aebischer, N.J. & Ewald, J.A. (2004) Managing the UK Grey Partridge *Perdix perdix* recovery: population change, reproduction, habitat and shooting. *Ibis* 146 (S2): 181–191.

Aebischer, N.J. & Ewald, J. (2010) Grey Partridge *Perdix perdix* in the UK: recovery status, set-aside and shooting. *Ibis* 152: 530–542.

Aebischer, N.J. & Potts, G.R. (1998) Spatial changes in Grey Partridge (*Perdix perdix*) distribution in relation to 25 years of changing agriculture in Sussex, UK. *Gibier Faune Sauvage* 15: 293–308.

Aebischer, N.J., Evans, A.D., Grice, P.V. & Vickery, J.A. (2000) *The Ecology and Conservation of Lowland Farmland Birds*. British Ornithologists' Union, Tring.

Ahola, M.P., Laaksonen, T., Eeva, T. & Lehikoinen, E. (2009) Great Tits lay increasingly smaller clutches than selected for: a study of climate- and density-related changes in reproductive traits. *Journal of Animal Ecology* 78: 1298–1306.

Amar, A. & Redpath, S. (2002) Determining the cause of the Hen Harrier decline on the Orkney Islands: an experimental test of two hypotheses *Animal Conservation* 5: 21–28.

Amar, A. & Redpath, S.M. (2005) Habitat use by Hen Harriers *Circus cyaneus* on Orkney: implications of land-use change for this declining population. *Ibis* 147: 37–47.

Amar, A., Arroyo, B. & Redpath, S. (2002) *Analysis of breeding success of Orkney Hen Harriers in relation to habitat*. Unpublished contract report to SNH: BAT/PA02e/01/02/100.

Amar, A., Redpath, S. & Thirgood, S. (2003) Evidence for food limitation in the declining hen harrier population on the Orkney Islands, Scotland *Biological Conservation* 111: 377–384.

Amar, A., Picozzi, N., Meek, E.R., Redpath, S.M. & Lambin, X. (2005) Decline of the Orkney Hen Harrier *Circus cyaneus* population: do changes to demographic parameters and mating system fit a declining food hypothesis? *Bird Study* 52: 18–24.

Amar, A., Hewson, C.M., Thewlis, R.M., Smith, K.W., Fuller, R.J., Lindsell, J.A., Conway, G., Butler, S. & MacDonald, M.A. (2006) *What's Happening to Our Woodland Birds? Long-term changes in the populations of woodland birds*. BTO Research Report 169 & RSPB Research Report 19. BTO, Thetford and RSPB, Sandy. [Full text](#)

Amar, A., Arroyo, B., Meek, E., Redpath, S. & Riley, H. (2008a) Influence of habitat on breeding performance of Hen Harriers *Circus cyaneus* in Orkney. *Ibis* 150: 400–404.

Amar, A., Thirgood, S., Pearce-Higgins, J. & Redpath, S. (2008b) The impact of raptors on the abundance of upland passerines and waders *Oikos* 117: 1143–1152.

Amar, A., Smith, K.W., Butler, S., Lindsell, J.A., Hewson, C.M., Fuller, R.J. & Charman, E.C. (2010) Recent patterns of change in vegetation structure and tree composition of British broadleaved woodland: evidence from large-scale surveys. *Forestry* 83: 345–356.

Amar, A., Court, I.R., Davison, M., Downing, S., Grimshaw, T., Pickford, T. & Raw, D. (2012) Linking nest histories, remotely sensed land use data and wildlife crime records to explore the impact of grouse moor management on peregrine falcon populations. *Biological Conservation* 145: 86–94. doi:[10.1016/j.biocon.2011.10.014](#)

Andrén, H. (1992) Corvid density and nest predation in relation to forest fragmentation: a landscape perspective. *Ecology* 73: 794–804.

Anganuzzi, A.A. (1993) A comparison of tests for detecting trends in abundance indices of dolphins. *Fishery Bulletin* 91: 183–194.

Angelstam, P., Breuss, M., Mikusinski, G., Stenström, M., Stighäll, K. & Thorell, D. (2002) Effects of forest structure on the presence of woodpeckers with different specialisation in a landscape history gradient in NE Poland. *Avian Landscape Ecology: pure and applied issues in the large-scale ecology of birds: Proceedings of the 11th annual IALE(UK) conference, 10–13 September 2002* (eds Chamberlain, D. & Wilson, A.), 25–38. University of East Anglia.

Anonymous (1995) *Biodiversity: the UK Steering Group report. Vol. 1: Meeting the Rio Challenge. Vol. 2: Action Plans*. HMSO, London.

Auer, S.K. & Martin, T.E. (2013) Climate change has indirect effects on resource use and overlap among coexisting bird species with negative consequences for their reproductive success. *Global Change Biology* 19: 411–419. doi:[10.1111/gcb.12062](#)

Austin, G.E. & Houston, D.C. (1997) The breeding performance of the Buzzard *Buteo buteo* in Argyll, Scotland and a comparison with other areas in Britain. *Bird Study* 44: 146–154.

Austin, G.E., Rehfish, M.M., Allan, J.R. & Holloway, S.J. (2007) Population size and differential population growth of introduced Greater Canada Geese *Branta canadensis* and re-established Greylag Geese *Anser anser* across habitats in Great Britain in the year 2000. *Bird Study* 54: 343–352. [Abstract](#)

Austin, G.E., Collier, M.P., Calbrade, N.A., Hall, C. & Musgrove, A.J. (2008) *Waterbirds in the UK 2006/07: The Wetland Bird Survey*. BTO/WWT/RSPB/JNCC, Thetford.

Full text

Austin, G.E., Read, W.J., Calbrade, N.A., Mellan, H.J., Musgrove, A.J., Skellorn, W., Hearn, R.D., Stroud, D.A., Wotton, S.R. & Holt, C.A., (2014) *Waterbirds in the UK 2011/12: the Wetland Bird Survey*. BTO, RSPB and JNCC in association with WWT, BTO, Thetford. [Full text](#), [interactive report](#).

Baillie, S.R. (1990) Integrated population monitoring of breeding birds in Britain and Ireland. *Ibis* 132: 151–166.

Baillie, S.R. (1991) Monitoring terrestrial breeding bird populations. In Goldsmith, F.B. (ed.) *Monitoring for Conservation and Ecology*: 112–132. Chapman & Hall, London.

Baillie, S.R. & Peach, W.J. (1992) Population limitation in Palaearctic–African migrant passerines. *Ibis* 134 Suppl. 1: 120–132.

Baillie, S.R. & Rehfisch, M.M. (eds) (2006) *National and site-based alert systems for UK birds*. Research Report 226. BTO, Thetford.

[Full text](#) (PDF, 1.28 MB)

Baillie, S.R., Crick, H.Q.P., Balmer, D.E., Beaven, L.P., Downie, I.S., Freeman, S.N., Leech, D.I., Marchant, J.H., Noble, D.G., Raven, M.J., Simpkin, A.P., Thewlis, R.M. & Wernham, C.V. (2002) *Breeding Birds in the Wider Countryside: their conservation status 2001* Research Report 278. BTO, Thetford. (www.bto.org/birdtrends2001)

Baillie, S.R., Brooks, S.P., King, R. & Thomas, L. (2009) Using a state-space model of the British Song Thrush *Turdus philomelos* population to diagnose the causes of a population decline. in *Modeling Demographic Processes in Marked Populations* (eds Thomson, D.L., Cooch, E.G. & Conroy, M.J.), pp 541–561. Springer, New York. [Contents](#)

Baines, D. (1990) The roles of predation, food and agricultural practice in determining the breeding success of the Lapwing *Vanellus vanellus* on upland grasslands. *Journal of Animal Ecology* 59: 915–929.

Baines, D. & Richardson, M. (2013) Hen harriers on a Scottish grouse moor: multiple factors predict breeding density and productivity. *Journal of Applied Ecology* 50: 1397–1405.

Baker, H., Stroud, D.A., Aebischer, N.J., Cranswick, P.A., Gregory, R.D., McSorley, C.A., Noble, D.G. & Rehfisch, M.M. (2006) Population estimates of birds in Great Britain and the United Kingdom. *British Birds* 99: 25–44. (APEP06)

Balmer, D.E., Adams, S.Y. & Crick H.Q.P. (2000) *Report on Barn Owl Release Scheme: Monitoring Project Phase II*. Research Report 250. BTO, Thetford.

[Full text](#) (PDF, 118.85 KB)

Balmer, D.E., Gillings, S., Caffrey, B.J., Swann, R.L., Downie, I.S. & Fuller, R.J. (eds) (2013) *Bird Atlas 2007–11: the breeding and wintering birds of Britain and Ireland*. BTO Books, Thetford.

Banks, A.N., Coombes, R.H. & Crick, H.Q.P. (2003) *The Peregrine Falcon breeding population of the UK & Isle of Man in 2002*. Research Report 330. BTO, Thetford.

[Full text](#) (PDF, 176.59 KB)

Banks, A.N., Crick, H.Q.P., Coombes, R., Benn, S., Ratcliffe, D.A. & Humphreys, E.M. (2010) The breeding status of Peregrine Falcons *Falco peregrinus* in the UK and Isle of Man in 2002. *Bird Study* 57: 421–436. doi: [10.1080/00063657.2010.511148](https://doi.org/10.1080/00063657.2010.511148)

Battaglia, A., Ghidini, S., Campanini, G. & Spaggiari, R. (2005) Heavy metal contamination in Little Owl (*Athene noctua*) and Common Buzzard (*Buteo buteo*) from northern Italy. *Ecotoxicology and Environmental Safety* 60: 61–66.

Bauer, Z., Trnka, M., Baueroová, J., Mozný, M., Stepánek, P., Bartosová, L., & Zalud, Z. (2010) Changing climate and the phenological response of Great Tit and Collared Flycatcher populations in floodplain forest ecosystems in Central Europe. *International Journal of Biometeorology* 54: 99–111.

Beale, C.M., Burfield, I.J., Sim, I.M.W., Rebecca, G.W., Pearce-Higgins, J.W. & Grant, M.C. (2006) Climate change may account for the decline in British ring ouzels *Turdus torquatus*. *Journal of Animal Ecology* 75: 826–835.

Bell, C.P., Baker, S.W., Parkes, N.G., Brooke, M. de L. & Chamberlain, D.E. (2010) The role of the Eurasian Sparrowhawk (*Accipiter nisus*) in the decline of the House Sparrow (*Passer domesticus*) in Britain. *Auk* 127: 411–420.

Bellebaum, J. & Bock, C. (2009) Influence of ground predators and water levels on Lapwing *Vanellus vanellus* breeding success in two continental wetlands. *Journal of Ornithology* 150: 221–230.

Benton, T.G., Bryant, D.M., Cole, L. & Crick, H.Q.P. (2002) Linking agricultural practice to insect and bird populations: a historical study over three decades. *Journal of Applied Ecology* 39: 673–687.

Besbeas, P., Freeman, S.N., Morgan, B.J.T. & Catchpole, E.A. (2002) Integrating mark–recapture–recovery and census data to estimate animal abundance and demographic parameters. *Biometrics* 58: 540–547.

Bibby, C.J. (1989) A survey of breeding Wood Warblers *Phylloscopus sibilatrix*, in Britain 1984–85. *Bird Study* 36: 56–72.

Bibby, C.J. & Etheridge, B. (1993) Status of the Hen Harrier *Circus cyaneus* in Scotland in 1988–89. *Bird Study* 40: 1–11.

BirdLife International (2004) *Birds in Europe: population estimates, trends and conservation status*. BirdLife Conservation Series no 12. BirdLife International, Cambridge. (BiE04)

BirdLife International (2013) *IUCN Red List for Birds*. URL: www.birdlife.org [31 July 2013].

Birkhead, M. & Perrins, C. (1985) The breeding biology of the Mute Swan *Cygnus olor* on the River Thames with special reference to lead poisoning. *Biological Conservation* 32: 1–11.

Birkhead, M., Bacon, P.J. & Walter, P. (1983) Factors affecting the breeding success of the Mute Swan *Cygnus olor*. *Journal of Animal Ecology* 52: 727–741.

- Blus, L.J. (1994) A review of lead poisoning in swans. *Comparative Biochemistry and Physiology Part C Pharmacology, Toxicology and Endocrinology* 108: 259–267.
- Boatman, N.D., Brickle, N.W., Hart, J.D., Milsom, T.P., Morris, A.J., Murray, A.W.A., Murray, K.A. & Robertson, P.A. (2004) Evidence for the indirect effects of pesticides on farmland birds. *Ibis* 146: 131–143.
- Bodey, T.W., McDonald, R.A., Sheldon, R.D. & Bearhop, S. (2011) Absence of effects of predator control on nesting success of Northern Lapwings *Vanellus vanellus*: implications for conservation. *Ibis* 153: 543–555.
- Bolton, M., Tyler, G., Smith, K. & Bamford, R. (2007) The impact of predator control on lapwing *Vanellus vanellus* breeding success on wet grassland nature reserves. *Journal of Applied Ecology* 44: 534–544.
- Bonham, P.F. & Robertson, J.C.M. (1975) The spread of Cetti's Warbler in north-west Europe. *British Birds* 68: 393–408.
- Both, C. (2002) Nemen Bonte Vliegenvangers *Ficedula hypoleuca* af door klimaatsverandering? [Decrease of European Pied Flycatchers due to climate change?] *Limosa* 75: 73–78.
- Both, C. (2010) Flexibility of timing of avian migration to climate change masked by environmental constraints en route. *Current Biology* 20: 243–248.
- Both, C., Bouwhuis, S., Lessells, C.M. & Visser, M.E. (2006) Climate change and population declines in a long-distance migratory bird. *Nature* 441 (4): 81–83. [Full text](#)
- Both, C., van Asch, M., Bijlsma, R.G., van den Burg, A.B. & Visser, M.E. (2009) Climate change and unequal phenological changes across four trophic levels: constraints or adaptations? *Journal of Animal Ecology* 78: 73–83.
- Both, C., Van Turnhout, C.A.M., Bijlsma, R.G., Siepel, H., Van Strien, A.J. & Foppen, R.P.B. (2010) Avian population consequences of climate change are most severe for long-distance migrants in seasonal habitats. *Proceedings of the Royal Society of London Series B* 277: 1259–1266. [Full text](#)
- Bradbury, R.B. & Bradter, U. (2004) Habitat associations of Yellow Wagtails *Motacilla flava flavissima* on lowland wet grassland. *Ibis* 146: 241–246.
- Bradbury, R. & Stoate, C. (2000) The ecology of Yellowhammers *Emberiza citrinella* on lowland farmland. In *Ecology and Conservation of Lowland Farmland Birds* (eds Aebischer, N.J., Evans, A.D., Grice, P.V. & Vickery, J.A.), 165–172. British Ornithologists' Union, Tring.
- Brickle, N.W. (1999) *The effect of agricultural intensification on the decline of the Corn Bunting, Miliaria calandra*. DPhil thesis, University of Sussex.
- Brickle, N.W. & Harper, D.G.C. (1999) Diet of nestling Corn Buntings *Miliaria calandra* in southern England examined by compositional analysis of faeces. *Bird Study* 46: 319–329.
- Brickle, N.W. & Harper, D.G.C. (2000) Habitat use by Corn Buntings *Miliaria calandra* in winter and summer. In *Ecology and Conservation of Lowland Farmland Birds* (eds Aebischer, N.J., Evans, A.D., Grice, P.V. & Vickery, J.A.), 156–164. British Ornithologists' Union, Tring.
- Brickle, N.W. & Harper, D.G. (2002) Agricultural intensification and the timing of breeding of Corn Buntings *Miliaria calandra*. *Bird Study* 49: 219–228.
- Brickle, N.W., Harper, D.G.C., Aebischer, N.J. & Cockayne, S.H. (2000) Effects of agricultural intensification on the breeding success of Corn Buntings *Miliaria calandra*. *Journal of Applied Ecology* 37: 742–755.
- Brindley, E., Norris, K., Cook, T., Babbs, S., Forster-Browne, C. & Yaxley, R. (1998) The abundance and conservation status of redshank (*Tringa totanus*) nesting on saltmarshes in Great Britain. *Biological Conservation* 86: 289–297.
- Brinkhof, M.W.G. & Cavé, A.J. (1997) Food supply and seasonal variation in breeding success: an experiment in the European Coot. *Proceedings of the Royal Society B Biological Sciences* 264: 291–296.
- Bro, E., Sarrazin, F., Clobert, J. & Reitz, F. (2000) Demography and the decline of the Grey Partridge *Perdix perdix* in France. *Journal of Applied Ecology* 37: 432–448.
- Brooke, M. de L. & Davies, N.B. (1987) Recent changes in host usage by cuckoos *Cuculus canorus* in Britain. *Journal of Animal Ecology* 56: 873–883.
- Broughton, R., Hinsley, S., Bellamy, P., Hill, R. & Rothery, P. (2006) Marsh Tit *Poecile palustris* territories in a British broad-leaved wood. *Ibis* 148: 744–752.
- Brown, A.F., Crick, H.Q.P. & Stillman, R.A. (1995) The distribution, numbers and breeding ecology of Twite *Acanthis flavirostris* in the south Pennines of England. *Bird Study* 42: 107–121.
- Brown, A.W. & Brown, L.M. (1984) The status of the Mute Swan in the Lothians. *Scottish Birds* 13: 8–15.
- Browne, S. & Aebischer, N. (2001) *The role of agricultural intensification in the decline of the Turtle Dove Streptopelia turtur*. English Nature, Peterborough.
- Browne, S. & Aebischer, N. (2002) Temporal changes in the breeding and feeding ecology of Turtle Doves (*Streptopelia turtur*) in the UK: an overview. *Zeitschrift für Jagdwissenschaft* 48: 215–221.
- Browne, S.J. & Aebischer, N.J. (2003) Temporal changes in the migration phenology of Turtle Doves *Streptopelia turtur* in Britain, based on sightings from coastal bird observatories. *Journal of Avian Biology* 34: 65–71.
- Browne, S.J. & Aebischer, N.J. (2004) Temporal changes in the breeding ecology of European Turtle Doves *Streptopelia turtur* in Britain, and implications for conservation. *Ibis* 146: 125–137.
- Browne, S. & Aebischer, N. (2005) Studies of West Palearctic birds: Turtle Dove. *British Birds* 98: 58–72.
- Browne, S., Vickery, J.A. & Chamberlain, D.E. (2000) Densities and population estimates of breeding Skylarks *Alauda arvensis* in Britain in 1997. *Bird Study* 47: 52–65. doi: [10.1080/00063650009461160](https://doi.org/10.1080/00063650009461160)

- Browne, S., Aebischer, N., Yfantis, G. & Marchant, J.H. (2004) Habitat availability and use by Turtle Doves *Streptopelia turtur* between 1965 and 1995: an analysis of Common Birds Census data. *Bird Study* 51: 1–11.
- Browne, S.J., Aebischer, N.J. & Crick, H.Q.P. (2005) The breeding ecology of Turtle Doves *Streptopelia turtur* in Britain during the period 1941 to 2000: an analysis of BTO Nest Record Cards. *Bird Study* 52: 1–9. doi: [10.1080/00063650509461368](https://doi.org/10.1080/00063650509461368)
- Browne, S.J., Aebischer, N.J., Moreby, S.J. & Teague, L. (2006) The diet and disease susceptibility of Grey Partridges *Perdix perdix* on arable farmland in East Anglia, England. *Wildlife Biology* 12: 3–10.
- Buchanan, G.M., Pearce-Higgins, J.W., Wotton, S.R., Grant, M.C. & Whitfield, D.P. (2003) Correlates of the change in Ring Ouzel *Turdus torquatus* abundance in Scotland from 1988–91 to 1999. *Bird Study* 50: 97–105. doi: [10.1080/00063650309461300](https://doi.org/10.1080/00063650309461300)
- Buckland, S.T., Cattanaach, K.L. & Anganuzzi, A.A. (1992) Estimating trends in abundance of dolphins associated with tuna in the eastern tropical Pacific Ocean, using sightings data collected on commercial tuna vessels. *Fishery Bulletin* 90: 1–20.
- Burfield, I.J. & Brooke, M. de L. (2005) The decline of the Ring Ouzel *Turdus torquatus* in Britain: evidence from bird observatory data. *Ringing & Migration* 22: 199–204.
- Burton, N.H.K. (2007) Influences of restock age and habitat patchiness on Tree Pipits *Anthus trivialis* breeding in Breckland pine plantations. *Ibis* 149 (suppl. 2): 193–204. doi: [10.1111/j.1474-919X.2007.00737.x](https://doi.org/10.1111/j.1474-919X.2007.00737.x)
- Burton, N.H.K. (2009) Reproductive success of Tree Pipits *Anthus trivialis* in relation to habitat selection in conifer plantations. *Ibis* 151: 361–372.
- Burton, N.H.K. & Conway, G.J. (2008) *Assessing population change of breeding Ringed Plovers in the UK between 1984 & 2007*. Research Report 503. BTO, Thetford.
- Buse, A., Dury, S.J., Woodburn, R.J.W., Perrins, C.M. & Good, J.E.G. (1999) Effects of elevated temperature on multi-species interactions: the case of pedunculate oak, winter moth and tits. *Functional Ecology* 13 (suppl.): 74–82.
- Butler, C. (2002) Breeding parrots in Britain. *British Birds* 95: 345–348.
- Butler, C.J. (2003) Population biology of the introduced Rose-ringed Parakeet *Psittacula krameri* in the UK. PhD thesis, University of Oxford.
- Butler, C.J., Cresswell, W., Gosler, A. & Perrins, C. (2013) The breeding biology of Rose-ringed Parakeets *Psittacula krameri* in England during a period of rapid population expansion. *Bird Study* 60: 527–532.
- Calbrade, N.A., Holt, C.A., Austin, G.E., Mellan, H.J., Hearn, R.D., Stroud, D.A., Wotton, S.R. & Musgrove, A.J. (2010) *Waterbirds in the UK 2008/09: the Wetland Bird Survey*. BTO/RSPB/JNCC in association with WWT, Thetford.
- Calladine, J. & Bray, J. (2012) The importance of altitude and aspect for breeding Whinchats *Saxicola rubetra* in the uplands: limitations of the uplands as a refuge for a declining, formerly widespread species? *Bird Study* 59: 43–51. doi: [10.1080/00063657.2011.623767](https://doi.org/10.1080/00063657.2011.623767)
- Campbell, L.H., Avery, M.I., Donald, P., Evans, A.D., Green, R.E. & Wilson, J.D. (1997) *A review of the indirect effects of pesticides on birds*. JNCC, Peterborough.
- Carpenter, J. (2008) *An investigation of causes of population decline in the Marsh Tit Poecile palustris in Britain*. DPhil Thesis, University of Oxford.
- Carpenter, J.E., Charman, E.C., Smart, J., Amar, A., Gruar, D., Bierman, S. & Grice, P. (2009) *Habitat associations of woodland birds II: completing the picture for woodland indicator species*. RSPB, Sandy.
- Carpenter, J., Smart, J., Amar, A., Gosler, A., Hinsley, S. & Charman, E. (2010) National-scale analyses of habitat associations of Marsh Tits *Poecile palustris* and Blue Tits *Cyanistes caeruleus*: two species with opposing population trends in Britain. *Bird Study* 57: 31–43. doi: [10.1080/0006365090302610](https://doi.org/10.1080/0006365090302610)
- Carter, I. (2001) *The Red Kite*. Arlequin Press, Chelmsford.
- Catchpole, E.A., Morgan, B.J.T., Freeman, S.N. & Peach, W.J. (1999) Modelling the survival of British Lapwings *Vanellus vanellus* using ring-recovery data and weather covariates. *Bird Study* 46 (suppl.): 5–13.
- Chamberlain, D.E. & Crick, H.Q.P. (1999) Population declines and reproductive performance of skylarks *Alauda arvensis* in different regions and habitats of Great Britain. *Ibis* 141: 38–51.
- Chamberlain, D.E. & Crick, H.Q.P. (2003) Temporal and spatial associations in aspects of reproductive performance of Lapwings *Vanellus vanellus* in the United Kingdom, 1962–99. *Ardea* 91: 183–196. [Abstract](#)
- Chamberlain, D. & Fuller, R. (2000) Local extinctions and changes in species richness of lowland farmland birds in England and Wales in relation to recent changes in agricultural land-use. *Agriculture, Ecosystems & Environment* 78: 1–17.
- Chamberlain, D.E. & Fuller, R.J. (2001) Contrasting patterns of change in the distribution and abundance of farmland birds in relation to farming system in lowland Britain. *Global Ecology and Biogeography* 10: 399–409.
- Chamberlain, D.E. & Siriwardena, G.M. (2000) The effects of agricultural intensification on Skylarks *Alauda arvensis*: evidence from monitoring studies in Great Britain. *Environmental Reviews* 8: 95–113.
- Chamberlain, D., Wilson, A., Browne, S. & Vickery, J. (1999) Effects of habitat type and management on the abundance of Skylarks in the breeding season. *Journal of Applied Ecology* 36: 856–870.
- Chamberlain, D.E., Vickery, J.A. & Gough, S. (2000a) Spatial and temporal distribution of breeding Skylarks *Alauda arvensis* in relation to crop type in periods of population increase and decrease. *Ardea* 88: 61–73.

- Chamberlain, D.E., Fuller, R.J., Bunce, R.G.H., Duckworth, J.C. & Shrubbs, M. (2000b) Changes in the abundance of farmland birds in relation to the timing of agricultural intensification in England and Wales. *Journal of Applied Ecology* 37: 771–788.
- Chamberlain, D.E., Toms, M.P., Cleary-McHarg, R. & Banks, A.N. (2007) House Sparrow (*Passer domesticus*) habitat use in urbanized landscapes. *Journal of Ornithology* 148: 453–462. doi: [10.1007/s10336-007-0165-x](https://doi.org/10.1007/s10336-007-0165-x)
- Chamberlain, D.E., Glue, D.E. & Toms, M.P. (2009) Sparrowhawk *Accipiter nisus* presence and winter bird abundance. *Journal of Ornithology* 150: 247–254. doi: [10.1007/s10336-008-0344-4](https://doi.org/10.1007/s10336-008-0344-4)
- Chamberlain, D.E., Austin, G.E., Green, R.E., Hulme, M.F. & Burton, N.H.K. (2013) Improved estimates of population trends of Great Cormorants *Phalacrocorax carbo* in England and Wales for effective management of a protected species at the centre of a human–wildlife conflict. *Bird Study* 60: 335–344.
- Charman, E., Carpenter, J. & Gruar, D. (2009) *Understanding the causes of decline in breeding bird numbers in England: a review of the evidence base for declining species in the woodland indicator for England*. Research Report 37. RSPB, Sandy. [Full text](#)
- Charman, E.C., Smith, K.W., Gruar, D.J., Dodd, S. & Grice, P.V. (2010) Characteristics of woods used recently and historically by Lesser Spotted Woodpeckers *Dendrocopos minor* in England. *Ibis* 152: 543–555. doi: [10.1111/j.1474-919X.2010.01020.x](https://doi.org/10.1111/j.1474-919X.2010.01020.x)
- Charman, E.C., Smith, K.W., Dillon, I.A., Dodd, S., Gruar, D.J., Cristinacce, A., Grice, P.V. & Gregory, R.D. (2012) Drivers of low breeding success in the Lesser Spotted Woodpecker *Dendrocopos minor* in England: testing hypotheses for the decline. *Bird Study* 59: 255–265.
- Chernetsov, N. & Huettmann, F. (2005) Linking global climate grid surfaces with local long-term migration monitoring data: spatial computations for the Pied Flycatcher to assess climate-related population dynamics on a continental scale. *Lecture Notes in Computer Science* 3482: 133–142. [Full text](#)
- Clements, R. (2001) The Hobby in Britain: a new population estimate. *British Birds* 94: 402–408.
- Clements, R. (2002) The Common Buzzard in Britain: a new population estimate. *British Birds* 95: 377–383.
- Clements, R. (2008) The Common Kestrel population in Britain. *British Birds* 101: 228–234.
- Clements, R.J. & Everett, C.M. (2011) Densities and dispersion of breeding Eurasian Hobbies *Falco subbuteo* in southeast England. *Bird Study* 59: 74–82.
- Coleman, A.E., Coleman, J.T., Coleman, P.A. & Minton, C.D.T. (2001) A 39-year study of a Mute Swan *Cygnus olor* population in the English Midlands. *Ardea* 89: 123–133.
- Conrad, K.F., Warren, M.S., Fox, R., Parsons, M.S. & Woiwod, I.P. (2006) Rapid declines of common, widespread British moths provide evidence of an insect biodiversity crisis. *Biological Conservation* 132: 279–291.
- Conway, G. & Burton, N. (2009) Changing fortunes for breeding plovers. *BTO News* 280: 10–11.
- Conway, G., Wotton, S., Henderson, I., Langston, R., Drewitt, A. & Currie, F. (2007) Status and distribution of European Nightjars *Caprimulgus europaeus* in the UK in 2004. *Bird Study* 54: 98–111. doi: [10.1080/00063650709461461](https://doi.org/10.1080/00063650709461461)
- Conway, G.J., Burton, N.H.K., Handschuh, M. & Austin, G.E. (2008) *UK population estimates from the 2007 Breeding Little Ringed Plover and Ringed Plover Surveys* Research Report 510. BTO, Thetford.
- Conway, G., Wotton, S., Henderson, I., Eaton, M., Drewitt, A. & Spencer, J. (2009) The status of breeding Woodlarks *Lullula arborea* in Britain in 2006. *Bird Study* 56: 310–325.
- Cook, A.S.C.P., Barimore, C., Holt, C.A., Read, W.J. & Austin, G.E. (2013) *Wetland Bird Survey Alerts 2009/2010: changes in numbers of wintering waterbirds in the Constituent Countries of the United Kingdom, Special Protection Areas (SPAs) and Sites of Special Scientific Interest (SSSIs)*. Research Report 641. BTO, Thetford. www.bto.org/webs-alerts
- Cornulier, T., Elston, D.A., Arcese, P., Benton, T.G., Douglas, D.J.T., Lambin, X., Reid, J., Robinson, R.A. & Sutherland, W.J. (2009) Estimating the annual number of breeding attempts from breeding dates using mixture models. *Ecology Letters* 12: 1184–1193. [Abstract](#)
- Cowley E. & Siriwardena, G.M. (2005) Long-term variation in survival rates of Sand Martins *Riparia riparia*: dependence on breeding and wintering ground weather, age and sex, and their population consequences. *Bird Study* 52: 237–251. doi: [10.1080/00063650509461397](https://doi.org/10.1080/00063650509461397)
- Cox, W.A., Thompson III, F.R. & Reidy, J.L. (2013) The effects of temperature on nest predation by mammals, birds, and snakes. *Auk* 130: 784–790.
- Cramp, S. & Perrins, C.M. (1994) *Handbook of the Birds of Europe, the Middle East and North Africa: the birds of the Western Palearctic*. Volume 8. Oxford University Press, Oxford.
- Crawley, M.J. (1993) *GLIM for Ecologists*. Blackwell Science, Oxford, UK.
- Crick, H.Q.P. (1992) *Trends in the breeding performance of Golden Plover in Britain*. Research Report 76. BTO, Thetford. [Full text](#)
- Crick, H.Q.P. (1993) Trends in breeding success of Merlins (*Falco columbarius*) in Britain from 1937–1989. In *Biology and Conservation of Small Falcons* (eds Nicholls, M.K. & Clarke, R.), pp 30–38. Hawk & Owl Trust, London.
- Crick, H.Q.P. (1997) Long-term trends in Corn Bunting *Miliaria calandra* productivity in Britain. In Donald, P.F. & Aebischer, N.J. (eds.) *The Ecology and Conservation of Corn Buntings* *Miliaria calandra*: 52–64. UK Nature Conservation No. 13. JNCC, Peterborough.
- Crick, H.Q.P. (1998) Decline in clutch size of Hen Harriers: reply. *BTO News* 218: 23.

- Crick, H.Q.P. & Ratcliffe, D.A. (1995) The Peregrine *Falco peregrinus* population of the United Kingdom in 1991. *Bird Study* 42: 1–19.
- Crick, H.Q.P. & Siriwardena, G.M. (2002) National trends in the breeding performance of House Sparrows. In *Investigation into the causes of the decline of Starlings and House Sparrows in Great Britain* (eds Crick, H.Q.P., Robinson, R.A., Appleton, G.F., Clark, N.A. & Rickard, A.D.), 163–191. Research Report 290. BTO, Thetford.
- Crick, H.Q.P. & Sparks, T.H. (1999) Climate change related to egg-laying trends. *Nature* 399: 423–424.
- Crick, H.Q.P., Dudley, C. & Glue, D.E. (1993) Breeding birds in 1991. *BTO News* 185: 15–18.
- Crick, H.Q.P., Dudley, C., Evans, A.D. & Smith, K.W. (1994) Causes of nest failure among buntings in the UK. *Bird Study* 41: 88–94.
- Crick, H.Q.P., Dudley, C., Glue, D.E. & Thomson, D.L. (1997) UK birds are laying eggs earlier. *Nature* 388: 526.
- Crick, H.Q.P., Baillie, S.R., Balmer, D.E., Bashford, R.I., Beaven, L.P., Dudley, C., Glue, D.E., Gregory, R.D., Marchant, J.H., Peach, W.J. & Wilson, A.M. (1998) *Breeding birds in the wider countryside: their conservation status (1972–1996)* Research Report 198. BTO, Thetford.
- Crick, H.Q.P., Robinson, R.A., Appleton, G.F., Clark, N.A. & Rickard, A.D. (eds) (2002) *Investigations into the causes of the decline of Starlings and House Sparrows in Great Britain*. BTO Research Report 290. 305 pp. DEFRA, Bristol.
- Crick, H.Q.P., Baillie, S.R. & Leech, D.I. (2003) The UK Nest Record Scheme: its value for science and conservation. *Bird Study* 50: 254–270.
- Cross, T. (2002) Common Raven (Raven) *Corvus corax*. In *The Migration Atlas: movements of the birds of Britain and Ireland* (eds Wernham, C.V., Toms, M.P., Marchant, J.H., Clark, J.A., Siriwardena, G.M. & Baillie, S.R.), pp 626–628. T. & A.D. Poyser, London.
- Crowe, O. (2012) *CBS trend 1998–2010*. BirdWatch Ireland unpublished report.
- Crowe, O., Coombes, R.H., Lysaght, L., O'Brien, C., Choudhury, K.R., Walsh, A.J., Wilson, J.H. & O'Halloran, J. (2010) Population trends of widespread breeding birds in the Republic of Ireland 1998–2008. *Bird Study* 57: 267–280. doi: [10.1080/00063651003615147](https://doi.org/10.1080/00063651003615147)
- Croxton, P.J., Sparks, T.H., Cade, M. & Loxton, R.G. (2006) Trends and temperature effects in the arrival of spring migrants in Portland (United Kingdom) 1959–2005. *Acta Ornithologica* 41: 103–111.
- Dadam, D., Barimore, C.J. & Leech, D.I. (2011) *The BTO Barn Owl Monitoring Programme: final report 2000–2009*. Research Report 577. BTO, Thetford.
- [Full text](#) (PDF, 7.50 MB)
- Delany, S., Greenwood, J.J.D. & Kirby, J. (1992) *National Mute Swan Survey 1990*. Report to the Joint Nature Conservation Committee. Wildfowl & Wetlands Trust, Slimbridge.
- De Leo, G., Focardi, S., Gatto, M. & Cattadori, I. (2004) The decline of the Grey Partridge in Europe: comparing demographics in traditional and modern agricultural landscapes. *Ecological Modelling* 177: 313–335.
- Dillon, I.A., Smith, T.D., Williams, S.J., Haysom, S. & Eaton, M.A. (2009) Status of Red-throated Divers *Gavia stellata* in Britain in 2006. *Bird Study* 56: 147–157.
- Dixon, A., Richards, C., Haffield, P., Roberts, G., Thomas, M. & Lowe, A. (2008) The National Peregrine Survey 2002: how accurate are the published results for Wales? *Welsh Birds* 5: 276–283.
- Dobson, A.P. & Hudson, P.J. (1992) Regulation and stability of a free-living host–parasite system *Trichostrongylus tenuis* in Red Grouse. II Population models. *Journal of Animal Ecology* 61: 487–500.
- Dolton, C.S. & Brooke M. de L. (1999) Changes in the biomass of birds breeding in Great Britain, 1968–88. *Bird Study* 46: 274–278.
- Donald, P.F. (1997) The Corn Bunting *Miliaria calandra* in Britain: a review of current status, patterns of decline and possible causes. In Donald, P.F. & Aebischer, N.J. (eds.) *The Ecology and Conservation of Corn Buntings* *Miliaria calandra*: 11–26. UK Nature Conservation no 13. Joint Nature Conservation Committee, Peterborough.
- Donald, P.F. & Evans, A.D. (1994) Habitat selection by Corn Buntings *Miliaria calandra* in winter. *Bird Study* 41: 199–210.
- Donald, P.F. & Evans, A.D. (1995) Habitat selection and population size of Corn Buntings *Miliaria calandra* breeding in Britain in 1993. *Bird Study* 42: 190–204.
- Donald, P.F. & Forrest, C. (1995) The effects of agricultural change on population size of Corn Buntings *Miliaria calandra* on individual farms. *Bird Study* 42: 205–215.
- Donald, P.F. & Morris, T.J. (2005) Saving the Sky Lark: new solutions for a declining farmland bird. *British Birds* 98: 570–578.
- Donald, P.F. & Vickery, J.A. (2000) The importance of cereal fields to breeding and wintering skylarks *Alauda arvensis* in the UK. In *Proceedings of the 1999 BOU Spring Conference: Ecology and Conservation of Lowland Farmland Birds* (eds Aebischer, N.J., Evans, A.D., Grice, P.V. & Vickery, J.A.), pp 140–150. British Ornithologists' Union, Tring.
- Donald, P.F. & Vickery, J.A. (2001) *The ecology and conservation of Skylarks*. RSPB, Sandy.
- Donald, P., Muirhead, L., Buckingham, D., Evans, A., Kirby, W. & Gruar, D. (2001) Body condition, growth rates and diet of Skylark *Alauda arvensis* nestlings on lowland farmland. *Ibis* 143: 658–669.
- Dougall, T.W., Holland, P.K. & Yalden, D.W. (2004) A revised estimate of the breeding population of Common Sandpipers *Actitis hypoleucos* in Great Britain and Ireland. *Wader Study Group Bulletin* 105: 42–49.
- Dougall, T.W., Holland, P.K. & Yalden, D.W. (2010) The population biology of Common Sandpipers in Britain. *British Birds* 103: 100–114.

- Douglas, D.J.T., Benton, T.G. & Vickery, J.A. (2010a) Contrasting patch selection of breeding Yellowhammers *Emberiza citrinella* in set-aside and cereal crops. *Bird Study* 57: 69–74.
- Douglas, D.J.T., Newson, S.E., Leech, D.I., Noble, D.G. & Robinson, R.A. (2010b) How important are climate-induced changes in host availability for population processes in an obligate brood parasite, the European cuckoo? *Oikos* 119: 1834–1840. doi: [10.1111/j.1600-0706.2010.18388.x](https://doi.org/10.1111/j.1600-0706.2010.18388.x)
- Driver, J. (2006) Raven *Corvus corax* population census of northwest Wales, 1998 to 2005. *Welsh Birds* 4: 442–453.
- Dunn, J.C. & Morris, A.J. (2012) Which features of UK farmland are important in retaining territories of the rapidly declining Turtle Dove *Streptopelia turtur*? *Bird Study* 59: 394–402.
- Dunn, P.O. & Møller, A.P. (2013) Changes in breeding phenology and population size of birds. *Journal of Animal Ecology*. doi: [10.1111/1365-2656.12162](https://doi.org/10.1111/1365-2656.12162)
- Durant, D., Tichit, M., Fritz, H. & Kerneis, E. (2008) Field occupancy by breeding Lapwings *Vanellus vanellus* and Redshanks *Tringa totanus* in agricultural wet grasslands. *Agriculture, Ecosystems & Environment* 128: 146–150.
- Eaton, M.A., Brown, A.F., Noble, D.G., Musgrove, A.J., Hearn, R.D., Aebischer, N.J., Gibbons, D.W., Evans, A. & Gregory, R.D. (2009) Birds of Conservation Concern 3: the population status of birds in the United Kingdom, Channel Islands and Isle of Man. *British Birds* 102: 296–341. (BoCC3) [Leaflet](#), [full text](#).
- Elliott, G.D. & Avery, M.I. (1991) A review of reports of buzzard persecution 1975–1989. *Bird Study* 38: 52–56.
- Eglington, S., Bolton, M., Smart, M., Sutherland, W., Watkinson, A. & Gill, J. (2010) Managing water levels on wet grasslands to improve foraging conditions for breeding Northern Lapwing *Vanellus vanellus*. *Journal of Applied Ecology* 47: 451–458.
- Enoksson, B. (1990) Autumn territories and population regulation in the Nuthatch *Sitta europaea*: an experimental study. *Journal of Animal Ecology* 59: 1047–1062.
- Enoksson, B. & Nilsson, S.G. (1983) Territory size and population density in relation to food supply in the Nuthatch *Sitta europaea* (Aves). *Journal of Animal Ecology* 52: 927–935.
- Eraud, C., Boutin, J.-M., Riviere, M., Brun, J., Barbraud, C. & Lormee, H. (2009) Survival of Turtle Doves *Streptopelia turtur* in relation to western Africa environmental conditions. *Ibis* 151: 186–190.
- Esselink, H. & Beekman, J.H. (1991) Between year variation and causes of mortality in the non-breeding population of the Mute Swan *Cygnus olor* in the Netherlands, with special reference to hunting. *Wildfowl* 42: 110–119.
- Etheridge, B., Summers, R.W. & Green, R.E. (1997) The effects of illegal killing and destruction of nests by humans on the population dynamics of the hen harrier *Circus cyaneus* in Scotland. *Journal of Applied Ecology* 34: 1081–1105.
- Evans, D.M., Redpath, S.M., Evans, S.A., Elston, D.A., Gardner, C.J., Dennis, P. & Pakeman, R.J. (2006) Low intensity, mixed livestock grazing improves the breeding abundance of a common insectivorous passerine. *Biology Letters* 2: 636–638. doi: [10.1098/rsbl.2006.0543](https://doi.org/10.1098/rsbl.2006.0543) [Full text](#)
- Evans, I.M. & Pienkowski, M.W. (1991) World status of the red kite: a background to the experimental reintroduction to England and Scotland. *British Birds* 84: 171–187.
- Evans, J., Wilson, J. & Browne, S. (1995) *The effect of organic farming regimes on breeding and winter bird populations – Part III. Habitat selection and breeding success of Skylarks (Alauda arvensis) on organic and conventional farmland*. Research Report 154. BTO, Thetford.
- Evans, K. (2004) The potential for interactions between predation and habitat change to cause population declines of farmland birds. *Ibis* 146: 1–13.
- Evans, K.L. & Robinson, R.A. (2004) Barn Swallows and agriculture. *British Birds* 97: 218–230.
- Ewald, J.A., Aebischer, N.J., Brickle, N.W., Moreby, S.J., Potts, G.R. & Wakeham-Dawson, A. (2002) Spatial variation in densities of farmland birds in relation to pesticide use and avian food resources. *Avian Landscape Ecology: pure and applied issues in the large-scale ecology of birds: Proceedings of the 11th annual IALE(UK) conference, 10–13 September 2002* (eds Chamberlain, D. & Wilson, A.), 305–312. University of East Anglia.
- Ewing, S.R., Rebecca, G.W., Heavisides, A., Court, I., Lindley, P., Ruddock, M., Cohen, S. & Eaton, M.A. (2011) Breeding status of the Merlin *Falco columbarius* in the UK in 2008. *Bird Study* 58: 379–389.
- Fewster, R.M., Buckland, S.T., Siriwardena, G.M., Baillie, S.R. & Wilson, J.D. (2000) Analysis of population trends for farmland birds using generalized additive models. *Ecology* 81: 1970–1984.
- Field, R.H. & Anderson, G.Q.A. (2004) Habitat use by breeding Tree Sparrows *Passer montanus*. *Ibis* 146: 60–68.
- Fielding, A., Haworth, P., Whitfield, P., McLeod, D. & Riley, H. (2011) *A conservation framework for Hen Harriers in the United Kingdom*. JNCC report 441. JNCC, Peterborough. [Full text](#)
- Fletcher, K., Howarth, D., Kirby, A., Dunn, R. & Smith, A. (2013) Effect of climate change on breeding phenology, clutch size and chick survival of an upland bird. *Ibis* 155: 456–463.
- Flyckt, G. (1999) Breeding biology of the Yellow Wagtail *Motacilla f. flava* in shore meadows in north-eastern Scania. *Ornis Svecica* 9: 217–223.
- Fox, R. (2013) The decline of moths in Great Britain: a review of possible causes. *Insect Conservation and Diversity* 6: 5–19. doi: [10.1111/j.1752-4598.2012.00186.x](https://doi.org/10.1111/j.1752-4598.2012.00186.x)
- Fox, T. & Heldbjerg, H. (2008) Which regional features of Danish agriculture favour the Corn Bunting in the contemporary farming landscape? *Agriculture, Ecosystems & Environment* 126: 261–269.
- Freeman, S.N. & Crick, H.Q.P. (2002) Population dynamics of House Sparrows *Passer domesticus* breeding in Britain: an integrated analysis. In *Investigation into the*

causes of the decline of starlings and house sparrows in Great Britain (eds Crick, H.Q.P., Robinson, R.A., Appleton, G.F., Clark, N.A. & Rickard, A.D.), pp 193–211. Research Report 290. BTO, Thetford.

Freeman, S.N. & Crick, H.Q.P. (2003) The decline of the Spotted Flycatcher *Muscicapa striata* in the UK: an integrated population model. *Ibis* 145: 400–412.

Freeman, S.N., Wernham, C.V. & Balmer, D.E. (2001) Long-term changes in the productivity of common songbirds in Britain and Ireland from constant effort ringing. Unpublished report. BTO, Thetford.

Freeman, S.N., Robinson, R.A., Clark, J.A., Griffin, B.M. & Adams, S.Y. (2002) Population dynamics of Starling *Sturnus vulgaris* breeding in Britain: an integrated analysis. In *Investigation into the causes of the decline of starlings and house sparrows in Great Britain* (eds Crick, H.Q.P., Robinson, R.A., Appleton, G.F., Clark, N.A. & Rickard, A.D.), pp 121–139. Research Report 290. BTO, Thetford.

Freeman, S.N., Noble, D.G., Newson, S.E. & Baillie, S.R. (2003) Modelling bird population changes using data from the Common Birds Census and the Breeding Bird Survey. Research Report 303. BTO, Thetford.

[Full text](#) (PDF, 526.15 KB)

Freeman, S.N., Noble, D.G., Newson, S.E. & Baillie, S.R. (2007a) Modelling bird population changes using data from different surveys: the Common Birds Census and the Breeding Bird Survey. *Bird Study* 54: 61–72. doi: [10.1080/00063650709461457](https://doi.org/10.1080/00063650709461457)

Freeman, S.N., Robinson, R.A., Clark, J.A., Griffin, B.M. & Adams, S.Y. (2007b) Changing demography and population decline in the Common Starling *Sturnus vulgaris*: a multisite approach to Integrated Population Monitoring. *Ibis* 149: 587–596. doi: [10.1111/j.1474-919X.2007.00684.x](https://doi.org/10.1111/j.1474-919X.2007.00684.x)

Freeman, S.N., Balmer, D.E. & Crick, H.Q.P. (2007c) Tawny Owl Survey 2005. *BTO News* 268: 6–7.

Fuller, R.J. (1995) *Bird life of woodland and forest*. Cambridge University Press, Cambridge.

Fuller, R.J. (1996) *Relationships between grazing and birds with particular reference to sheep in the British uplands*. Research Report 164. BTO, Thetford.

Fuller, R. (2000) Relationships between recent changes in lowland British agriculture and farmland bird populations: an overview. In *Ecology and Conservation of Lowland Farmland Birds* (eds Aebischer, N.J., Evans, A.D., Grice, P.V. & Vickery, J.A.), 5–16. British Ornithologists' Union, Tring.

Fuller, R. & Ausden, M. (2008) Birds and habitat change in Britain. Part 1: a review of losses and gains in the twentieth century *British Birds* 101: 644–675.

Fuller, R.J. & Gough, S.J. (1999) Changes in sheep numbers in Britain: implications for bird populations. *Biological Conservation* 91: 73–89.

Fuller, R. & Moreton, B. (1987) Breeding bird populations of Kentish sweet chestnut (*Castanea sativa*) coppice in relation to age and structure of the coppice. *Journal of Applied Ecology* 24: 13–27.

Fuller, R.J., Marchant, J.H. & Morgan, R.A. (1985) How representative of agricultural practice in Britain are Common Birds Census farmland plots? *Bird Study* 32: 56–70.

Fuller, R.J., Gregory, R.D., Gibbons, D.W., Marchant, J.H., Wilson, J.D., Baillie, S.R. & Carter, N. (1995) Population declines and range contractions among lowland farmland birds in Britain. *Conservation Biology* 9: 1425–1441.

Fuller, R.J., Chamberlain, D.E., Burton, N.H.K. & Gough, S.J. (2001) Distributions of birds in lowland agricultural landscapes of England and Wales: how distinctive are bird communities of hedgerows and woodland? *Agriculture, Ecosystems & Environment* 84: 79–92.

Fuller, R.J., Noble, D.G., Smith, K.W. & Vanhinsbergh, D. (2005) Recent declines in populations of woodland birds in Britain: a review of possible causes *British Birds* 98: 116–143.

Fuller, R., Atkinson, P., Garnett, M., Conway, G., Bibby, C. & Johnstone, I. (2006) Breeding bird communities in the upland margins (ffridd) of Wales in the mid-1980s. *Bird Study* 53: 177–186.

Furness, R.W. & Greenwood, J.J.D. (eds.) (1993) *Birds as Monitors of Environmental Change*. Chapman & Hall, London.

Galbraith, H. (1988) Effects of agriculture on the breeding ecology of lapwings *Vanellus vanellus*. *Journal of Applied Ecology* 25: 487–503.

Gibbons, D.W., Reid, J.B. & Chapman, R.A. (1993) *The New Atlas of Breeding Birds in Britain and Ireland: 1988–1991*. T. & A.D. Poyser, London.

Gibbons, D., Gates, S., Green, R.E., Fuller, R.J. & Fuller, R.M. (1995) Buzzards *Buteo buteo* and Ravens *Corvus corax* in the uplands of Britain: limits to distribution and abundance. *Ibis* 137: 75–84.

Gibbons, D.W., Avery, M.I., Baillie, S.R., Gregory, R.D., Kirby, J., Porter, R.F., Tucker, G.M. & Williams, G. (1996) Bird species of conservation concern in the United Kingdom, Channel Islands and Isle of Man: revising the Red Data List. *RSPB Conservation Review* 10: 7–18.

Gibbons, D.W., Bainbridge, I.P., Mudge, G.P., Tharme, A.P. & Ellis, P.M. (1997) The status and distribution of the Red-throated Diver *Gavia stellata* in Britain in 1994. *Bird Study* 44: 194–205.

Gilbert, G. (2012) Grasshopper Warbler *Locustella naevia* breeding habitat in Britain. *Bird Study* 59: 303–314.

Gill, R.M.A. & Beardall, V. (2001) The impact of deer on woodlands: the effects of browsing and seed dispersal on vegetation structure and composition *Forestry* 74: 209–218.

Gill, R.M.A. & Fuller, R.J. (2007) The effects of deer browsing on woodland structure and songbirds in lowland Britain *Ibis* 149 suppl. 2: 119–127. doi: [10.1111/j.1474-919X.2007.00731.x](https://doi.org/10.1111/j.1474-919X.2007.00731.x)

Gillings, S., Newson, S.E., Noble, D.G. & Vickery, J.A. (2005) Winter availability of cereal stubbles attracts declining farmland birds and positively influences breeding

population trends. *Proceedings of the Royal Society B* 272: 733–739. [Full text](#)

Gillings, S., Newson, S. & Sellers, R.M. (2007) Breeding population estimates for Northern Wheatear in Britain. *British Birds* 100: 179–181.

Gilroy, J., Anderson, G., Grice, P., Vickery, J., Bray, I., Watts, P.N. & Sutherland, W. (2008) Could soil degradation contribute to farmland bird declines? Links between soil penetrability and the abundance of Yellow Wagtails *Motacilla flava* in arable fields. *Biological Conservation* 14: 3116–3126.

Gilroy, J.J., Anderson, G.Q.A., Grice, P.V., Vickery, J.A., Watts, P.N. & Sutherland, W.J. (2009) Foraging habitat selection, diet and nestling condition in Yellow Wagtails *Motacilla flava* breeding on arable farmland. *Bird Study* 56: 221–232.

Gilroy, J.J., Anderson, G.Q.A., Grice, P.V., Vickery, J.A. & Sutherland, W.J. (2010) Mid-season shifts in the habitat associations of Yellow Wagtails *Motacilla flava* breeding in arable farmland. *Ibis* 152: 90–104.

Glue, D.E. (1990) Breeding biology of the Grasshopper Warbler in Britain. *British Birds* 83: 131–145.

Glue, D. (1993) Report on Garden Bird Feeding Survey. *BTO News* 188: 24.

Glue, D. (1995) Report on Garden Bird Feeding Survey. *BTO News* 200: 5.

Glue, D. (1997) Report on Garden Bird Feeding Survey 1996/1997. *BTO News* 212: 6–7.

Glue, D.E. (2006) Cuckoos in crisis? *BTO News* 263: 22–23.

Glue, D. & Boswell, T. (1994) Comparative nesting ecology of the three British breeding woodpeckers. *British Birds* 87: 253–269.

Gooch, S., Baillie, S.R. & Birkhead, T.R. (1991) Magpie *Pica pica* and songbird populations – retrospective investigation of trends in population density and breeding success. *Journal of Applied Ecology* 28: 1068–1086.

Goodenough, A.E., Elliot, S.L. & Hart, A.G. (2009) The challenges of conservation for declining migrants: are reserve-based initiatives during the breeding season appropriate for the Pied Flycatcher *Ficedula hypoleuca*? *Ibis* 151: 429–439.

Goodenough, A.E., Hart, A.G. & Elliot, S.L. (2011) What prevents phenological adjustment to climate change in migrant bird species? Evidence against the “arrival constraint” hypothesis. *International Journal of Biometeorology* 55: 97–102.

Gosler, A.G. (1990) The Birds of Wytham – an historical survey. *Fritillary* 1: 29–74.

Goss-Custard, J.C. (1993) The effect of migration and scale on the study of bird populations: 1991 Witherby Lecture. *Bird Study* 40: 81–96.

Goszczynski, J. (1997) Density and productivity of Common Buzzard *Buteo buteo* and Goshawk *Accipiter gentilis* populations in Rogow, central Poland. *Acta Ornithologica* 32: 149–154.

Goszczynski, J. (2001) The breeding performance of the Common Buzzard *Buteo buteo* and Goshawk *Accipiter gentilis* in central Poland. *Acta Ornithologica* 36: 105–110.

Graham, I.M., Redpath, S.M. & Thirgood, S.J. (1995) The diet and breeding density of Common Buzzards *Buteo buteo* in relation to indices of prey abundance. *Bird Study* 42: 165–173.

Green, M., Kaleta, R. & Keirle, I. (2012) Habitat associations and winter distribution of Ring Ouzels in the Atlas Mountains, Morocco. *British Birds* 105: 674–682.

Green, R.E. (1999) Applications of large-scale studies of demographic rates to bird conservation. *Bird Study* 46 (Suppl.): S279–S288.

Greenwood, J.J.D. (2000) How BTO's monitoring of birds contributes to conservation. In *European Monitoring for Nature Conservation, Bonn-Bad Godesberg (Bundesamt für Naturschutz)*, ed. by C. Bischoff & R. Droschmeister. *Schriftenreihe für Landschaftspflege und Naturschutz* 62: 105–117.

Greenwood, J.J.D. & Baillie, S.R. (1991) Effects of density-dependence and weather on population changes of English passerines using a non-experimental paradigm. *Ibis* 133: 121–133. doi:[10.1111/j.1474-919X.1991.tb07675.x](https://doi.org/10.1111/j.1474-919X.1991.tb07675.x)

Gregory, R.D. & Marchant, J.H. (1996) Population trends of Jays, Magpies, Jackdaws and Carrion Crows in the United Kingdom. *Bird Study* 43: 28–37.

Gregory, R.D., Carter, S.P. & Baillie, S.R. (1997) Abundance, distribution and habitat use of breeding Goosanders *Mergus merganser* and Red-breasted Mergansers *Mergus serrator* on British rivers. *Bird Study* 44: 1–12.

Gregory, R.D., Wilkinson, N.I., Noble, D.G., Robinson, J.A., Brown, A.F., Hughes, J., Procter, D., Gibbons, D.W. & Galbraith, C.A. (2002) The population status of birds in the United Kingdom, Channel Islands and Isle of Man: an analysis of conservation concern 2002–2007. *British Birds* 95: 410–448.

Gregory, R.D., Noble, D.G. & Custance, J. (2004) The state of play of farmland birds: population trends and conservation status of lowland farmland birds in the United Kingdom. *Ibis* 146 (suppl. 2): 1–13.

Groom, D.W. (1993) Magpie *Pica pica* predation on Blackbird *Turdus merula* nests in urban areas. *Bird Study* 40: 55–62.

Gruar, D., Barritt, D. & Peach, W.J. (2006) Summer utilization of Oilseed Rape by Reed Buntings *Emberiza schoeniclus* and other farmland birds. *Bird Study* 53: 47–54.

Guillemain, M., Elmberg, J., Gauthier-Clerc, M., Massez, G., Hearn, R., Champagnon, J. & Simon, G. (2010) Wintering French Mallard and Teal are heavier and in better body condition than 30 years ago: effects of a changing environment? *Ambio* 39: 170–180.

Halley, D.J. (1993) Population changes and territorial distribution of Common Buzzards *Buteo buteo* in the Central Highlands, Scotland. *Bird Study* 40: 24–30.

- Halupka, L., Dyrce, A. & Borowiec, M. (2008) Climate change affects breeding of Reed Warblers *Acrocephalus scirpaceus*. *Journal of Avian Biology* 39: 95–100.
- Hancock, M.H. & Wilson, J.D. (2003) Winter habitat associations of seed-eating passerines on Scottish farmland. *Bird Study* 50: 116–130.
- Hardey, J., Crick, H.Q.P., Wernham, C.V., Riley, H.T., Etheridge, B. & Thompson, D.B.A. (eds) (2009) *Raptors: a field guide to survey and monitoring*. Second edition. The Stationery Office, Edinburgh.
- Harris, S., Morris, P., Wray, S. & Yalden, D. (1995) *A review of British mammals: population estimates and conservation status of British mammals other than cetaceans*. JNCC, Peterborough.
- Hart, J., Milsom, T., Fisher, G., Wilkins, V., Moreby, S., Murray, A. & Robertson, P. (2006) The relationship between Yellowhammer breeding performance, arthropod abundance and insecticide applications on arable farmland. *Journal of Applied Ecology* 43: 81–91.
- Hastie, T.J. & Tibshirani, R.J. (1990) *Generalized additive models*. Chapman & Hall, London.
- Hayhow, D.B., Eaton, M.A., Bladwell, S., Etheridge, B., Ewing, S.R., Ruddock, M., Saunders, R., Sharpe, C., Sim, I.M.W. & Stevenson, A. (2013) The status of the Hen Harrier, *Circus cyaneus*, in the UK and Isle of Man in 2010. *Bird Study* 60: 446–458.
- Heldbjerg, H. & Fox, T. (2008) Long-term population declines in Danish trans-Saharan migrant birds. *Bird Study* 55: 267–279.
- Henderson, I.G. & Hart, P.J.B. (1993) Provisioning, parental investment and reproductive success in Jackdaws *Corvus monedula*. *Ornis Scandinavica* 24: 142–148.
- Henderson, I.G., Wilson, A.M., Steele, D. & Vickery, J.A. (2002) Population estimates, trends and habitat associations of breeding Lapwing *Vanellus vanellus*, Curlew *Numenius arquata* and Snipe *Gallinago gallinago* in Northern Ireland in 1999. *Bird Study* 49: 17–25. doi: [10.1080/00063650209461240](https://doi.org/10.1080/00063650209461240)
- Henderson, I.G., Fuller, R.J., Conway, G.J. & Gough, S.J. (2004) Evidence for declines in populations of grassland-associated birds in marginal upland areas of Britain. *Bird Study* 51: 12–19.
- Henderson, I., Holt, C. & Vickery, J. (2007) National and regional patterns of habitat association with foraging Barn Swallows *Hirundo rustica* in the UK. *Bird Study* 54: 371–377. doi: [10.1080/00063650709461497](https://doi.org/10.1080/00063650709461497)
- Heubeck, M. (2013) Censusing and monitoring breeding seabirds in Britain and Ireland. *British Birds* 106: 306–324.
- Hewson, C. & Noble, D. (2009) Population trends of breeding birds in British woodlands over a 32-year period: relationships with food, habitat use and migratory behaviour. *Ibis* 151: 464–486.
- Hewson, C.M., Amar, A., Lindsell, J.A., Thewlis, R.M., Butler, S., Smith, K. & Fuller, R.J. (2007) Recent changes in bird populations in British broadleaved woodland. *Ibis* 149 (suppl. 2): 14–28. doi: [10.1111/j.1474-919X.2007.00745.x](https://doi.org/10.1111/j.1474-919X.2007.00745.x)
- Hill, D.A. (1984) Population regulation in the Mallard (*Anas platyrhynchos*). *Journal of Animal Ecology* 53: 191–202.
- Hines, J.E., Kendall, W.L., Nichols, J.D. & Thompson, F.R. III (2003) On the use of the robust design with transient capture–recapture models. *Auk* 120: 1151–1158. doi: [10.1642/0004-8038\(2003\)120\[1151:OTUOTR\]2.0.CO;2](https://doi.org/10.1642/0004-8038(2003)120[1151:OTUOTR]2.0.CO;2)
- Hinsley, S.A., Bellamy, P.E., Newton, I. & Sparks, T.H. (1995) Habitat and landscape factors influencing the presence of individual breeding bird species in woodland fragments. *Journal of Avian Biology* 26: 94–104.
- Hinsley, S.A., Carpenter, J.E., Broughton, R.K., Bellamy, P.E., Rothery, P., Amar, A., Hewson, C.M. & Gosler, A.G. (2007) Habitat selection by Marsh Tits *Poecile palustris* in the UK. *Ibis* 149 (suppl. 2): 224–233.
- Hole, D.G. (2001) *The population ecology and ecological genetics of the house sparrow Passer domesticus on farmland in Oxfordshire*. PhD thesis, University of Oxford.
- Hole, D.G., Whittingham, M.J., Bradbury, R.B., Anderson, G.Q.A., Lee, P.L.M., Wilson, J.D. & Krebs, J.R. (2002) Agriculture: widespread local House Sparrow extinctions. *Nature* 418: 931–932.
- Holland, P.K. & Yalden, D.W. (2002) Population dynamics of Common Sandpipers *Actitis hypoleucos* in the Peak District of Derbyshire – a different decade: a report of the failure of a population to recover from a catastrophic snow storm. *Bird Study* 49: 131–138.
- Holling, M. & the Rare Breeding Birds Panel (2007a) Rare breeding birds in the United Kingdom in 2003 and 2004. *British Birds* 100: 321–367. [Full text RBBP reports online](#)
- Holling, M. & the Rare Breeding Birds Panel (2007b) Non-native breeding birds in the United Kingdom in 2003, 2004 and 2005. *British Birds* 100: 638–649. [Full text](#)
- Holling, M. & the Rare Breeding Birds Panel (2008) Rare breeding birds in the United Kingdom in 2005. *British Birds* 101: 276–316. [Full text RBBP reports online](#)
- Holling, M. & the Rare Breeding Birds Panel (2009) Rare breeding birds in the United Kingdom in 2006. *British Birds* 102: 158–202. [Full text RBBP reports online](#)
- Holling, M. & the Rare Breeding Birds Panel (2010a) Rare breeding birds in the United Kingdom in 2007. *British Birds* 103: 2–52. [Full text RBBP reports online](#)
- Holling, M. & the Rare Breeding Birds Panel (2010b) Rare breeding birds in the United Kingdom in 2008. *British Birds* 103: 482–538. [Full text RBBP reports online](#)
- Holling, M. & the Rare Breeding Birds Panel (2011a) Non-native breeding birds in the United Kingdom in 2006, 2007 and 2008. *British Birds* 104: 114–138. [Full text](#)
- Holling, M. & the Rare Breeding Birds Panel (2011b) Rare breeding birds in the United Kingdom in 2009. *British Birds* 104: 476–537. [Full text RBBP reports online](#)
- Holling, M. & the Rare Breeding Birds Panel (2012) Rare breeding birds in the United Kingdom in 2010. *British Birds* 105: 352–416. [Full text RBBP reports online](#)

- Holling, M. & the Rare Breeding Birds Panel (2013) Rare breeding birds in the United Kingdom in 2011. *British Birds* 106: 496–554.
- Holloway, S. (1996) *The Historical Atlas of Breeding Birds in Britain and Ireland 1875–1900*. T. & A.D. Poyser, London.
- Holt, C.A., Fuller, R.J. & Dolman, P.M. (2010) Experimental evidence that deer browsing reduces habitat suitability for breeding Common Nightingales *Luscinia megarhynchos*. *Ibis* 152: 335–346.
- Holt, C.A., Austin, G.E., Calbrade, N.A., Mellan, H.J., Hearn, R.D., Stroud, D.A., Wotton, S.R. & Musgrove, A.J. (2012a) *Waterbirds in the UK 2010/11: the Wetland Bird Survey*. BTO/RSPB/JNCC, Thetford. [Full text](#)
- Holt, C.A., Hewson, C.M. & Fuller, R.J. (2012b) The Nightingale in Britain: status, ecology and conservation needs *British Birds* 105: 172–187.
- Holt, C.A., Fraser, K.H., Bull, A.J. & Dolman, P.M. (2012c) Habitat use by Nightingales in a scrub–woodland mosaic in central England *Bird Study* 59: 416–425. doi: [10.1080/00063657.2012.722191](https://doi.org/10.1080/00063657.2012.722191)
- Holt, C.A., Fuller, R.J. & Dolman, P.M. (2012d) Deer reduce habitat quality for a woodland songbird: evidence from settlement patterns, demographic parameters and body condition. *Auk* 130: 13–20. [Full text](#)
- Hoodless, A.N., Lang, D., Aebischer, N.J., Fuller, R.J. & Ewald, J.A. (2009) Densities and population estimates of breeding Eurasian Woodcock *Scolopax rusticola* in Britain in 2003. *Bird Study* 56: 15–25.
- Hopkins, J.J. & Kirby, K.J. (2007) Ecological change in British broadleaved woodland since 1947. *Ibis* 149: 29–40.
- Hórak, P. & Lebreton, J.-D. (1998) Survival of adult Great Tits *Parus major* in relation to sex and habitat: a comparison of urban and rural populations. *Ibis* 140: 205–209.
- Hötter, H. (1991) Waders breeding on wet grasslands in the countries of the European Community – a brief summary of current knowledge on population sizes and population trends. *Wader Study Group Bulletin* 61 (suppl.): 50–55.
- Hudson, P.J. (1992) *Grouse in space and time*. Game Conservancy Trust, Fordingbridge.
- Hudson, R. (1972) Collared Doves in Britain and Ireland during 1965–70. *British Birds* 65: 139–155.
- Hudson, R., Tucker, G.M. & Fuller, R.J. (1994) Lapwing *Vanellus vanellus* populations in relation to agricultural changes: a review. In Tucker, G.M., Davies, S.M. & Fuller, R.J. (eds) *The Ecology and Conservation of Lapwings* *Vanellus vanellus*: 1–33. UK Nature Conservation no 9. JNCC, Peterborough.
- Hughes, S.W.M., Bacon, P., & Flegg, J.J.M. (1979) The 1975 census of the Great Crested Grebe in Britain *Bird Study* 26: 213–226.
- Hutchinson, C.D. (1989) *Birds in Ireland*. T. & A.D. Poyser, Calton.
- Inglis, I.R., Isaacson, A.J., Smith, G.C., Haynes, P.J. & Thearle, R.J.P. (1997) The effect on the Woodpigeon *Columba palumbus* of the introduction of oilseed rape into Britain. *Agriculture, Ecosystems & Environment* 61: 113–121.
- Jackson, D.B. (2007) Factors affecting the abundance of introduced hedgehogs (*Erinaceus europaeus*) to the Hebridean island of South Uist in the absence of natural predators and implications for nesting birds. *Journal of Zoology* 271: 210–217.
- Jackson, D. & Green, R. (2000) The importance of the introduced hedgehog (*Erinaceus europaeus*) as a predator of the eggs of waders (Charadrii) on machair in South Uist, Scotland. *Biological Conservation* 93: 333–348.
- Jackson, D.B., Fuller, R.J. & Campbell, S.T. (2004) Long-term population changes among breeding shorebirds in the Outer Hebrides, Scotland, in relation to introduced hedgehogs (*Erinaceus europaeus*). *Biological Conservation* 117: 151–166.
- Jenny, M. (1990) Populationsdynamik der Feldlerche *Alauda arvensis* in einer intensiv genutzten Agrarlandschaft der schweizerischen Mittellandes. *Ornithologische Beobachter* 8: 153–163.
- JNCC (1996) *Birds of Conservation Importance*. Press release (31 May 1996). Joint Nature Conservation Committee, Peterborough.
- JNCC (2012) *Seabird population trends and causes of change: 2012 report*. Joint Nature Conservation Committee. [Full text](#)
- Johnson, D.H. (1979) Estimating nest success: The Mayfield method and an alternative. *Auk* 96: 651–661.
- Johnstone, I., Dyda, J. & Lindley, P. (2007) The population status and hatching success of Curlews *Numenius arquata* in Wales in 2006. *Welsh Birds* 5: 78–87.
- Johnstone, I., Dyda, J. & Lindley, P. (2008) The population status of breeding Golden Plover and Dunlin in Wales in 2007 *Welsh Birds* 5: 300–310.
- Joys, A.C., Noble, D.G. & Baillie, S.R. (2003) *Evaluation of species coverage and precision using the BBS indexing method*. Research Report 317. BTO, Thetford.
- [Full text](#) (PDF, 330.42 KB)
- Julliard, R. (2004) Estimating the contribution of survival and recruitment to large scale population dynamics. *Animal Biodiversity and Conservation* 27: 417–426.
- Kallander, H. (1997) The Nuthatch *Sitta europaea* population of Dalby Soderskog during 15 years: trend and fluctuations. *Ornis Svecica* 7: 143–148.
- King, R., Brooks, S.P., Mazzetta, C., Freeman, S.N. & Morgan, B.J.T. (2008) Identifying and diagnosing population declines: a Bayesian assessment of Lapwings in the UK. *Applied Statistics* 57: 609–632.
- Kirby, J., Delany, S. & Quinn, J. (1994) Mute Swans in Great Britain: a review, current status and long-term trends *Hydrobiologia* 279–280: 467–482.

Kirby, W.B., Anderson, G.Q.A., Grice, P.V., Soanes, L., Thompson, C. & Peach, W.J. (2012) Breeding ecology of Yellow Wagtails *Motacilla flava* in an arable landscape dominated by autumn-sown crops. *Bird Study* 59: 383–393.

Kostrzewa, A. & Kostrzewa, R. (1990) The relationship of spring and summer weather with density and breeding performance of the Buzzard *Buteo buteo*, Goshawk *Accipiter gentilis* and Kestrel *Falco tinnunculus*. *Ibis* 132: 550–559.

Kostrzewa, R. & Kostrzewa, A. (1991) Winter weather, spring and summer density, and subsequent breeding success of Eurasian Kestrels, Common Buzzards, and Northern Goshawks. *Auk* 108: 342–347.

Krebs, J.R., Wilson, J.D., Bradbury R.B. & Siriwardena, G.M. (1999) The second silent spring? *Nature* 400: 611–612.

Kuijper, D.P.J., Oosterveld, E. & Wymenga, E. (2009) Decline and potential recovery of the European Grey Partridge (*Perdix perdix*) population – a review. *European Journal of Wildlife Research* 55: 455–463. [Full text](#)

Kyrkos, A. (1997) *Behavioural and demographic responses of yellowhammers to variation in agricultural practices* DPhil thesis, University of Oxford.

Kyrkos, A., Wilson, J. & Fuller, R. (1998) Farmland habitat change and abundance of Yellowhammers *Emberiza citrinella*: an analysis of Common Birds Census data. *Bird Study* 45: 232–246.

Lampila, S., Orell, M., Belda, E. & Koivula, K. (2006) Importance of adult survival, local recruitment and immigration in a declining boreal forest passerine, the Willow Tit *Parus montanus*. *Oecologia* 148: 405–413.

Langston, R.H.W., Liley, D., Murison, G., Woodfield, E. & Clarke, R.T. (2007) What effects do walkers and dogs have on the distribution and productivity of breeding European Nightjar *Caprimulgus europaeus*? *Ibis* 149, supplement 1: 27–36.

Lawson, B., Lachish, S., Colvile, K.M., Durrant, C., Peck, K.M., Toms, M.P., Sheldon, B.C. & Cunningham, A.A. (2012a) Emergence of a novel avian pox disease in British tit species. *PLoS ONE* 7(11): e40176. doi:[10.1371/journal.pone.0040176](https://doi.org/10.1371/journal.pone.0040176)

Lawson, B., Robinson, R.A., Colvile, K.M., Peck, K.M., Chantrey, J., Pennycott, T.W., Simpson, V.R., Toms, M.P. & Cunningham, A.A. (2012b) The emergence and spread of finch trichomonosis in the British Isles. *Philosophical Transactions of the Royal Society B* 367: 2852–2863. doi:[10.1098/rstb.2012.0130](https://doi.org/10.1098/rstb.2012.0130)

Lebreton, J.-D., Burnham, K.P., Clobert, J. & Anderson, D.R. (1992) Modeling survival and testing biological hypotheses using marked animals: a unified approach with case studies. *Ecological Monographs* 62: 67–118. [Full text](#)

Leech, D. & Barimore, C. (2008) Is avian breeding success weathering the storms? *BTO News* 279: 19–20.

Leech, D. & Crick, H. (2005) Nest Record Scheme breeding trends – latest results. *BTO News* 261: 18–19.

Leech, D.I., Crick, H.Q.P. & Shawyer, C.R. (2005) *The BTO Barn Owl Monitoring Programme: fourth year 2003*. Research Report 411. BTO, Thetford.

[Abstract](#) (PDF, 362.61 KB)

Leech, D., Crick, H. & Shawyer, C. (2006a) Barn Owls and winter weather. *BTO News* 262: 8–9.

Leech, D., Barimore, C. & Crick, H. (2006b) NRS Concern List – five new species added. *BTO News* 267: 4–5.

Leech, D., Barimore, C. & Crick, H. (2007) Volunteer boom. *BTO News* 273: 18–19.

Le Gouar, P.J., Schekkerman, H., van der Jeugd, H.P., Boele, A., van Harxen, R., Fuchs, P., Stroeken, P. & van Noordwijk, A.J. (2011) Long-term trends in survival of a declining population: the case of the Little Owl (*Athene noctua*) in the Netherlands. *Oecologia* 166: 369–379.

Lehikoinen, A., Green, M., Husby, M., Kålås, J.A. & Lindström, Å. (2014) Common montane birds are declining in northern Europe. *Journal of Avian Biology* 45: 3–14. doi: [10.1111/j.1600-048X.2013.00177.x](https://doi.org/10.1111/j.1600-048X.2013.00177.x)

Letty, J., Génot, J.-C. & Sarrazin, F. (2001) Analysis of population viability of Little Owl (*Athene noctua*) in the Northern Vosges natural park (north-eastern France). *Alauda* 69: 359–372.

Lewis, A.J.G., Amar, A., Cordi-Piec, D. & Thewlis, R.M. (2007) Factors influencing Willow Tit *Poecile montanus* site occupancy: a comparison of abandoned and occupied woods. *Ibis* 149 (suppl. 2): 205–213. doi: [10.1111/j.1474-919X.2007.00733.x](https://doi.org/10.1111/j.1474-919X.2007.00733.x)

Lewis, A.J.G., Amar, A., Daniells, L., Charman, E.C., Grice, P. & Smith, K. (2009a) Factors influencing patch occupancy and within-patch habitat use in an apparently stable population of Willow Tits *Poecile montanus kleinschmidtii* in Britain. *Bird Study* 56: 326–337.

Lewis, A.J.G., Amar, A., Charman, E.C. & Stewart, F.R.P. (2009b) The decline of the Willow Tit in Britain. *British Birds* 102: 386–393.

Liker, A. & Szekely, T. (1997) The impact of grazing and road use on hatching success of Lapwings (*Vanellus vanellus*). *Acta Zoologica Academiae Scientiarum Hungaricae* 43: 85–92.

Liley, D. & Sutherland, W.J. (2007) Predicting the population consequences of human disturbance for Ringed Plovers *Charadrius hiaticula*: a game theory approach. *Ibis* 149, supplement 1: 82–94.

Little, B., Davison, M. & Jardine, D. (1995) Merlins *Falco columbarius* in Kielder Forest: influences of habitat on breeding performance. *Forest Ecology and Management* 79: 147–152. [Abstract](#)

MacDonald, M.A. & Bolton, M. (2008a) Predation on wader nests in Europe. *Ibis* 150 (suppl 1): 54–73.

MacDonald, M.A. & Bolton, M. (2008b) Predation of Lapwing *Vanellus vanellus* nests on lowland wet grassland in England and Wales: effects of nest density, habitat and

predator abundance. *Journal of Ornithology* 149: 555–563.

MacLeod, R., Clark, J. & Cresswell, W. (2008) The starvation–predation risk trade-off, body mass and population status in the Common Starling *Sturnus vulgaris*. *Ibis* 150 (suppl. 1): 199–208.

Madders, M. (2000) Habitat selection and foraging success of Hen Harriers *Circus cyaneus* in west Scotland. *Bird Study* 47: 32–40.

Mallord, J.W., Dolman, P.M., Brown, A.F. & Sutherland, W.J. (2007) Linking recreational disturbance to population size in a ground-nesting passerine *Journal of Applied Ecology* 44: 185–195. doi: [10.1111/j.1365-2664.2006.01242.x](https://doi.org/10.1111/j.1365-2664.2006.01242.x)

Mallord, J.W., Orsman, C.J., Cristinacce, A., Butcher, N., Stowe, T.J. & Charman, E.C. (2012a) Mortality of Wood Warbler *Phylloscopus sibilatrix* nests in Welsh oakwoods: predation rates and the identification of nest predators using miniature nest cameras. *Bird Study* 59: 286–295.

Mallord, J.W., Charman, E.C., Cristinacce, A. & Orsman, C.J. (2012b) Habitat associations of Wood Warblers *Phylloscopus sibilatrix* breeding in Welsh oakwoods. *Bird Study* 59: 403–415.

Malpas, L.R., Smart, J., Drewitt, A., Sharps, E. & Garbutt, A. (2013) Continued declines of Redshank *Tringa totanus* breeding on saltmarsh in Great Britain: is there a solution to this conservation problem? *Bird Study* 60: 370–383.

Manly, B.F.J. (1991) *Randomisation and Monte Carlo Methods in Biology*. Chapman & Hall, London.

Marchant, J.H. & Gregory, R.D. (1999) Numbers of nesting Rooks *Corvus frugilegus* in the United Kingdom in 1996. *Bird Study* 46: 258–273. [Abstract](#)

Marchant, J.H., Hudson, R., Carter, S.P. & Whittington, P.A. (1990) *Population Trends in British Breeding Birds* BTO, Tring.

Marchant, J.H., Freeman, S.N., Crick, H.Q.P. & Beaven, L.P. (2004) The BTO Heronries Census of England and Wales 1928–2000: new indices and a comparison of analytical methods. *Ibis* 146: 323–334. doi: [10.1111/j.1474-919X.2004.00272.x](https://doi.org/10.1111/j.1474-919X.2004.00272.x)

Marquiss, M. (2007) Seasonal pattern in hawk predation on Common Bullfinches *Pyrrhula pyrrhula*: evidence of an interaction with habitat affecting food availability. *Bird Study* 54: 1–11.

Marquiss, M., Newton, I. & Ratcliffe, D.A. (1978) The decline of the Raven *Corvus corax* in relation to afforestation in southern Scotland and northern England. *Journal of Applied Ecology* 15: 129–144.

Martin, T.G., Arcese, P. & Scheerder, N. (2011) Browsing down our natural heritage: deer impacts on vegetation structure and songbird assemblages across an island archipelago. *Biological Conservation* 144: 459–469.

Martinez, J.A. & Zuberogitia, I. (2004) Effects of habitat loss on perceived and actual abundance of the Little Owl *Athene noctua* in eastern Spain. *Ardeola* 51: 215–219.

Martínez-Padilla, J., Redpath, S.M., Zeineddine, M. & Mougeot, F. (2014) Insights into population ecology from long-term studies of red grouse *Lagopus lagopus scoticus*. *Journal of Animal Ecology* 83: 85–98. doi: [10.1111/1365-2656.12098](https://doi.org/10.1111/1365-2656.12098)

Mason, C. & Macdonald, S. (1999) Habitat use by Lapwings and Golden Plovers in a largely arable landscape. *Bird Study* 46: 89–99.

Mason, C. & Macdonald, S. (2000) Influence of landscape and land-use on the distribution of breeding birds in farmland in eastern England. *Journal of Zoology* 251: 339–348.

Mason, C.F. & Macdonald, S.M. (2006) Recent marked decline in Corn Bunting numbers in northeast Essex. *British Birds* 99: 206–214.

Mason, W.L. (2007) Changes in the management of British forests between 1945 and 2000 and possible future trends. *Ibis* 149: 41–52.

Massimino, D., Johnston, A., Pearce-Higgins, J. & Baillie, S. (2013) *Maps of population density and trends*. www.bto.org/volunteer-surveys/bbs/latest-results/maps-population-density-and-trends

Matthysen, E. (1989) Nuthatch *Sitta europaea* demography, beech mast, and territoriality. *Ornis Scandinavica* 20: 278–282.

Mavor, R.A., Parsons, M., Heubeck, M., Pickerell, G. & Schmitt, S. (2003) *Seabirds numbers and breeding success in Britain and Ireland, 2002*. UK Nature Conservation no. 27. JNCC, Peterborough. [Abstract/Full text](#)

Mavor, R.A., Parsons, M., Heubeck, M. & Schmitt, S. (2004) *Seabird numbers and breeding success in Britain and Ireland, 2003*. UK Nature Conservation no. 28. JNCC, Peterborough. [Abstract/Full text](#)

Mavor, R.A., Parsons, M., Heubeck, M. & Schmitt, S. (2006) *Seabird numbers and breeding success in Britain and Ireland, 2005*. UK Nature Conservation no. 30. JNCC, Peterborough. [Abstract/Full text](#)

Mavor, R.A., Heubeck, M., Schmitt, S. & Parsons, M. (2008) *Seabird numbers and breeding success in Britain and Ireland, 2006*. UK Nature Conservation no. 31. JNCC, Peterborough. [Abstract/Full text](#)

Maxwell, J. (2002) Nest-site competition with Blue Tits and Great Tits as a possible cause of declines in Willow Tit numbers: observations in the Clyde area. *Glasgow Naturalist* 24: 47–50.

Maxwell, J. (2003) Scottish Willow Tits hanging on. *BTO News* 244: 24.

Mayfield, H. (1961) Nesting success calculated from exposure. *Wilson Bulletin* 73: 255–261.

Mayfield, H. (1975) Suggestions for calculating nest success. *Wilson Bulletin* 87: 456–466.

- McShea, W.J. & Rappole, J.H. (2000) Managing the abundance and diversity of breeding bird populations through manipulation of deer densities. *Conservation Biology* 14: 1161–1170.
- Mearns, R. (1983) The status of the Raven in southern Scotland and Northumbria. *Scottish Birds* 12: 211–218.
- Meek, W.R., Burman, P.J., Nowakowski, M., Sparks, T.H. & Burman, N.J. (2003) Barn owl release in lowland southern England – a twenty-one year study. *Biological Conservation* 109: 271–282.
- Messenger, A. & Roome, M. (2007) The breeding population of the Hobby in Derbyshire. *British Birds* 100: 594–608.
- Milsom, T.P. (2005) Decline of Northern Lapwing *Vanellus vanellus* breeding on arable farmland in relation to loss of spring tillage. *Bird Study* 52: 297–306.
- Mitchell, C., Patterson, D., Boyer, P., Cunningham, P., McDonald, R., Meek, E., Okill, J.D. & Symonds, F. (2000) The summer status and distribution of Greylag Geese in north and west Scotland. *Scottish Birds* 21: 69–77.
- Mitchell, C., King, R. & Cook, T. (2002) Mallard *Anas platyrhynchos*. In *The Migration Atlas: movements of the birds of Britain and Ireland* (eds Wernham, C.V., Toms, M.P., Marchant, J.H., Clark, J.A., Siriwardena, G.M. & Baillie, S.R.), pp 193–195. T. & A.D. Poyser, London.
- Mitchell, C., Griffin, L., Trinder, M., Newth, J. & Urquhart, C. (2011) The status and distribution of summering Greylag Geese *Anser anser* in Scotland, 2008–09. *Bird Study* 58: 338–348.
- Mitchell, C., Hearn, R. & Stroud, D. (2012) The merging of populations of Greylag Geese breeding in Britain. *British Birds* 105: 498–505.
- Mitchell, P.I., Newton, S.F., Ratcliffe, N. & Dunn, T.E. (2004) *Seabird Populations of Britain and Ireland*. T. & A.D. Poyser, London. [Summary](#)
- Møller, A.P., Saino, N., Adamík, P., Ambrosini, R., Antonov, A., Campobello, D., Stokke, B.G., Fossøy, F., Lehikoinen, E., Martin-Vivaldi, M., Moksnes, A., Moskat, C., Røskoft, E., Rubolini, D., Schulze-Hagen, K., Soler, M. & Shykoff, J.A. (2011) Rapid change in host use of the Common Cuckoo *Cuculus canorus* linked to climate change. *Proceedings of the Royal Society B Biological Sciences* 278: 733–738.
- Monaghan, P., Uttley, J.D., Burns, M.D., Thaine, C. & Blackwood, J. (1989) The relationship between food supply, reproductive effort and breeding success in Arctic Terns *Sterna paradisaea*. *Journal of Animal Ecology* 58: 261–274.
- Monaghan, P., Uttley, J.D. & Burns, M.D. (1992) Effect of changes in food availability on reproductive effort in Arctic terns *Sterna paradisaea*. *Ardea* 80: 71–81.
- Moorcroft, D. & Wilson, J.D. (2000) The ecology of Linnets *Carduelis cannabina* on lowland farmland. In *Ecology and Conservation of Lowland Farmland Birds* (eds Aebischer, N.J., Evans, A.D., Grice, P.V. & Vickery, J.A.), pp 173–181. British Ornithologists' Union, Tring.
- Moorcroft, D., Bradbury, R.B. & Wilson, J.D. (1997) The diet of nestling Linnets *Carduelis cannabina* before and after agricultural intensification. *1997 Brighton Crop Protection Conference – Weeds, Conference Proceedings* 3: 923–928. British Crop Protection Council, Farnham.
- Moorcroft, D., Wilson, J.D. & Bradbury, R.B. (2006) The diet of nestling Linnets *Carduelis cannabina* on lowland farmland before and after agricultural intensification. *Bird Study* 53: 156–162.
- Morris, A.J. & Gilroy, J.J. (2008) Close to the edge: predation risks for two declining farmland passerines. *Ibis* 150: 168–177.
- Morris, A., Burges, D., Fuller, R.J., Evans, A.D. & Smith, K.W. (1994) The status and distribution of Nightjars *Caprimulgus europaeus* in Britain in 1992. *Bird Study* 41: 181–191.
- Morris, A., Wilson, J., Whittingham, M. & Bradbury, R. (2005) Indirect effects of pesticides on breeding Yellowhammer *Emberiza citrinella*. *Agriculture, Ecosystems & Environment* 106: 1–16.
- Morrison, C.A., Robinson, R.A., Clark, J.A. & Gill, J.A. (2010) Spatial and temporal variation in population trends in a long-distance migratory bird. *Diversity & Distributions* 16: 620–627.
- Morrison, C.A., Robinson, R.A., Clark, J.A., Marca, A.D., Newton, J. & Gill, J.A. (2013) Using stable isotopes to link breeding population trends to winter ecology in Willow Warblers, *Phylloscopus trochilus*. *Bird Study* 60: 211–220.
- Morrison, C.A., Robinson, R.A., Leech, D.I., Dadam, D. & Toms, M. (2014) Using citizen science to investigate the role of productivity in House Sparrow *Passer domesticus* population trends. *Bird Study* 61: 91–100. doi: [10.1080/00063657.2013.874975](https://doi.org/10.1080/00063657.2013.874975)
- Moss, D. & Moss, G.M. (1993) Breeding biology of the Little Grebe *Tachybaptus ruficollis* in Britain and Ireland. *Bird Study* 40: 107–114.
- Mountford, M.D. (1982) Estimation of population fluctuations with application to the Common Birds Census. *Applied Statistics* 31: 135–143.
- Mountford, M.D. (1985) An index of population change with an application to the Common Birds Census. In Morgan, B.J.T. & North, P.M. (eds) *Statistics in Ornithology*. 121–132. Springer-Verlag, Berlin.
- Musgrove, A.J., Austin, G.E., Hearn, R.D., Holt, C.A., Stroud, D.A. & Wotton, S.R. (2011) Overwinter population estimates of British waterbirds. *British Birds* 104: 364–397.
- Musgrove, A.J., Aebischer, N.J., Eaton, M.A., Hearn, R.D., Newson, S.E., Noble, D.G., Parsons, M., Risely, K. & Stroud, D.A. (2013) Population estimates of birds in Great Britain and the United Kingdom. *British Birds* 106: 64–100.
- Nelson, S.H., Court, I., Vickery, J.A., Watts, P.N. & Bradbury, R.B. (2003) The status and ecology of the Yellow Wagtail in Britain. *British Wildlife* 14: 270–274.
- Newson, S.E., Woodburn, R.J.W., Noble, D.G., Baillie, S.R. & Gregory, R.D. (2005) Evaluating the Breeding Bird Survey for producing national population size and

density estimates. *Bird Study* 52: 42–54. doi: [10.1080/00063650509461373](https://doi.org/10.1080/00063650509461373)

Newson, S.E., Ekins, G.R., Marchant, J.H., Rehfisch, M.M. & Sellers, R.M. (2006) *The status of inland and coastal breeding Great Cormorants* *Phalacrocorax carbo* in England. Research Report 433. BTO, Thetford.

[Abstract](#) (PDF, 405.10 KB)

Newson, S.E., Marchant, J.H., Ekins, G.R. & Sellers, R.M. (2007) The status of inland-breeding Great Cormorants in England. *British Birds* 100: 289–299.

Newson, S.E., Evans, K.L., Noble, D.G., Greenwood, J.J.D. & Gaston, K.J. (2008) Use of distance sampling to improve estimates of national population sizes for common and widespread breeding birds in the UK. *Journal of Applied Ecology* 45: 1330–1338. [Full text](#)

Newson, S.E., Ockendon, N., Joys, A., Noble, D.G. & Baillie, S.R. (2009) Comparison of habitat-specific trends in the abundance of breeding birds in the UK. *Bird Study* 56: 233–243.

Newson, S.E., Leech D.I., Hewson, C.M., Crick, H.Q.P. & Grice, P.V. (2010) Potential impact of grey squirrels *Sciurus carolinensis* on woodland bird populations in England. *Journal of Ornithology* 151: 211–218.

Newson, S.E., Rexstad, E.A., Baillie, S.R., Buckland, S.T. & Aebischer, N.J. (2010b) Population change of avian predators and grey squirrels in England: is there evidence for an impact on avian prey populations? *Journal of Applied Ecology* 47: 244–252.

Newson, S.E., Johnston, A., Parrott, D. & Leech, D.I. (2011) Evaluating the population-level impact of an invasive species, Ring-necked Parakeet *Psittacula krameri*, on native avifauna. *Ibis* 153: 509–516. [Abstract](#)

Newson, S.E., Marchant, J.H., Sellers, R.M., Ekins, G.R., Hearn, R.D. & Burton, N.H.K. (2013) Colonisation and range expansion of inland-breeding Cormorants in England. *British Birds* 106: 737–743.

Newton, I. (1986) *The Sparrowhawk*. T. & A.D. Poyser, Calton.

Newton, I. (1988) A key factor analysis of a sparrowhawk population. *Oecologia* 76: 588–596.

Newton, I. (2004) The recent declines of farmland bird populations in Britain: an appraisal of causal factors and conservation actions. *Ibis* 146: 579–600.

Newton, I. (2006) Movement patterns of Common Crossbills *Loxia curvirostra* in Europe. *Ibis* 148: 782–788.

Newton, I. (2013) Organochlorine pesticides and birds. *British Birds* 106: 189–205.

Newton, I. & Marquiss, M. (1986) Population regulation in Sparrowhawks. *Journal of Animal Ecology* 55: 463–480.

Newton, I. & Wyllie, I. (1992) Recovery of a Sparrowhawk population in relation to declining pesticide contamination. *Journal of Applied Ecology* 29: 476–484.

Nilsson, A.L.K., Knudsen, E., Jerstad, K., Røstad, O.W., Walseng, B., Slagsvold, T. & Stenseth, N.C. (2011) Climate effects on population fluctuations of the white-throated dipper *Cinclus cinclus*. *Journal of Animal Ecology* 80: 235–243.

Nilsson, S.G. (1982) Seasonal variation in the survival rate of adult Nuthatches *Sitta europaea* in Sweden. *Ibis* 124: 96–100.

Nilsson, S.G. (1987) Limitation and regulation of population density in the Nuthatch *Sitta europaea* (Aves) breeding in natural cavities. *Journal of Animal Ecology* 56: 921–937.

Nilsson, S.G., Olsson, O., Svensson, S. & Wiktander, U. (1992) Population trends and fluctuations in Swedish woodpeckers. *Ornis Svecica* 2: 13–21.

Norman, D. & Peach, W.J. (2013) Density-dependent survival and recruitment in a long-distance Palaearctic migrant, the Sand Martin *Riparia riparia*. *Ibis* 155: 284–296.

Norris, K., Brindley, E., Cook, T., Babbs, S., Forster-Brown, C. & Yaxley, R. (1998) Is the density of Redshank *Tringa totanus* nesting on saltmarshes in Great Britain declining due to changes in grazing management? *Journal of Applied Ecology* 35: 621–634.

O'Brien, M. (2005) Estimating the number of farmland waders breeding in the United Kingdom. *International Wader Studies* 14: 135–139.

O'Brien, M. & Wilson, J.D. (2011) Population changes of breeding waders on farmland in relation to agri-environment management. *Bird Study* 58: 399–408.

Ockendon, N., Hewson, C.M., Johnston, A. & Atkinson, P.W. (2012) Declines in British-breeding populations of Afro-Palaearctic migrant birds are linked to bioclimatic wintering zone in Africa, possibly via constraints on arrival time advancement. *Bird Study* 59: 111–125. doi: [10.1080/00063657.2011.645798](https://doi.org/10.1080/00063657.2011.645798)

Ockendon, N., Leech, D. & Pearce-Higgins, J.W. (2013) Climatic effects on breeding grounds are more important drivers of breeding phenology in migrant birds than carry-over effects from wintering grounds. *Biology Letters* 9. doi: [10.1098/rsbl.2013.0669](https://doi.org/10.1098/rsbl.2013.0669)

O'Connor, R.J. & Marchant, J.H. (1981) *A field validation of some Common Birds Census techniques*. Research Report 4. BTO, Tring.

O'Connor, R.J. & Mead, C.J. (1984) The Stock Dove in Britain, 1930–1980. *British Birds* 77: 181–201.

O'Connor, R.J. & Shrubb, M. (1986) *Farming and Birds*. Cambridge University Press, Cambridge.

Orell, M. (1989) Population fluctuations and survival of Great Tits *Parus major* dependent on food supplied by man in winter. *Ibis* 131: 112–127.

Ormerod, S.J. & Tyler, S.J. (1989) Long-term change in the suitability of Welsh streams for Dippers *Cinclus cinclus* as a result of acidification and recovery: a modelling study. *Environmental Pollution* 62: 171–182.

- Ormerod, S.J. & Tyler, S.J. (1990) Environmental pollutants in the eggs of Welsh Dippers *Cinclus cinclus*: a potential monitor of organochlorine and mercury contamination in upland rivers. *Bird Study* 37: 171–176.
- Ormerod, S.J., O'Halloran, J., Gribbin, S.D. & Tyler, S.J. (1991) The ecology of Dippers *Cinclus cinclus* in relation to stream acidity in upland Wales: breeding performance, calcium physiology and nestling growth. *Journal of Applied Ecology* 28: 419–433.
- Owen, M., Atkinson-Willes, G.L. & Salmon, D.G. (1986) *Wildfowl in Great Britain*. Cambridge University Press, Cambridge.
- PACEC (2006) *The Economic and Environmental Impact of Sporting Shooting*. Public and Corporate Economic Consultants, Cambridge. [Full text](#)
- Panek, M. (2005) Demography of Grey Partridges *Perdix perdix* in Poland in the years 1991–2004: reasons of population decline. *European Journal of Wildlife Research* 51: 14–18.
- Pannekoek, J. & van Strien, A. (1996) *TRIM (TRends and Indices for Monitoring data)*. Research paper 9634. Statistics Netherlands, Voorburg.
- Parkin, D.T., Collinson, M., Helbig, A.J., Knox, A.G. & Sangster, G. (2003) The taxonomic status of Carrion and Hooded Crows. *British Birds* 96: 274–290.
- Parr, S.J. (1994) Changes in the population size and nest sites of Merlins *Falco columbarius* in Wales between 1970 and 1991. *Bird Study* 41: 42–47.
- Payevsky, V. (2006) Mechanisms of population dynamics in trans-Saharan bird migrants: a review. *Zoologicheskyy Zhurnal* 85: 368–381.
- Peach, W.J. (1993) Combining mark recapture data sets for small passerines. In *Marked Individuals in the Study of Bird Populations* (eds J.-D. Lebreton & P.M. North), pp. 107–122. Birkhauser Verlag, Basel, Switzerland.
- Peach, W.J. & Baillie, S.R. (1994) Implementation of the Mountford indexing method for the Common Birds Census. In Hagemeyer, W. & Verstraël, T. (eds.) *Bird Numbers 1992. Distribution, Monitoring and Ecological Aspects*: 653–662. Proc. 12th Int. Conf. International Bird Census Council and European Ornithological Atlas Committee. SOVON, Beek-Ubbergen.
- Peach, W.J., Baillie, S.R. & Underhill, L. (1991) Survival of British Sedge Warblers *Acrocephalus schoenobaenus* in relation to west African rainfall. *Ibis* 133: 300–305.
- Peach, W.J., Thompson, P.S. & Coulson, J.C. (1994) Annual and long-term variation in the survival rates of British lapwings *Vanellus vanellus*. *Journal of Animal Ecology* 63: 60–70.
- Peach, W.J., Crick, H.Q.P. & Marchant, J.H. (1995a) The demography of the decline in the British Willow Warbler population. *Journal of Applied Statistics* 22: 905–922.
- Peach, W.J., du Feu, C. & McMeeking, J. (1995b) Site tenacity and survival rates of Wrens *Troglodytes troglodytes* and Treecreepers *Certhia familiaris* in a Nottinghamshire wood. *Ibis* 137: 497–507.
- Peach, W.J., Buckland, S.T. & Baillie, S.R. (1996) The use of constant effort mist-netting to measure between-year changes in the abundance and productivity of common passerines. *Bird Study* 43: 142–156.
- Peach, W.J., Baillie, S.R. & Balmer, D.E. (1998) Long-term changes in the abundance of passerines in Britain and Ireland as measured by constant effort mist-netting. *Bird Study* 45: 257–275.
- Peach, W.J., Siriwardena, G.M. & Gregory, R.D. (1999) Long-term changes in over-winter survival rates explain the decline of Reed Buntings *Emberiza schoeniclus* in Britain. *Journal of Applied Ecology* 36: 798–811. [Abstract](#)
- Peach, W.J., Robinson, R.A. & Murray, K.A. (2004) Demographic and environmental causes of the decline of rural Song Thrushes *Turdus philomelos* in lowland Britain. *Ibis* 146 (Suppl. 2): 50–59. doi: [10.1111/j.1474-919X.2004.00362.x](https://doi.org/10.1111/j.1474-919X.2004.00362.x)
- Peach, W.J., Vincent, K.E., Fowler, J.A. & Grice, P.V. (2008) Reproductive success of house sparrows along an urban gradient. *Animal Conservation* 11: 493–503. doi: [10.1111/j.1469-1795.2008.00209.x](https://doi.org/10.1111/j.1469-1795.2008.00209.x)
- Pearce-Higgins, J.W., Yalden, D.W. & Whittingham, M.J. (2005) Warmer springs advance the breeding phenology of golden plovers *Pluvialis apricaria* and their prey (Tipulidae). *Oecologia* 143: 470–476.
- PECBMS (2007) *The State of Europe's Common Birds 2007*. CSO/RSPB, Prague. [Full text](#)
- PECBMS (2009) *The State of Europe's Common Birds 2008*. CSO/RSPB, Prague. [Full text](#)
- PECBMS (2010) *Trends of common birds in Europe, 2010 update*. European Bird Census Council, Prague. www.ebcc.info/index.php?ID=387
- PECBMS (2011a) *Population Trends of Common European Breeding Birds 2011*. CSO, Prague. www.ebcc.info/index.php?ID=469 (leaflet), www.ebcc.info/index.php?ID=457 (full report, including graphs and methods)
- PECBMS (2011b) European wild bird indicators, 2011 update. www.ebcc.info/index.php?ID=459
- PECBMS (2012a) *Population trends of common European breeding birds 2012*. CSO, Prague. [Leaflet](#), [full text](#) (including graphs and methods)
- PECBMS (2012b) *European wild bird indicators, 2012 update*. [Full text](#)
- PECBMS (2013a) *Population trends of common European breeding birds 2013*. CSO, Prague. [Leaflet](#), [full text](#) (including graphs and methods)
- PECBMS (2013b) *European wild bird indicators, 2013 update*. [Full text](#)
- Percival, S.M. (1990) *Population trends in British Barn Owls, Tyto alba, and Tawny Owls, Strix aluco, in relation to environmental change*. Research Report 57. BTO,

Tring.

Perdeck, A.C., Visser, M.E. & Van Balen, J.H. (2000) Great Tit *Parus major* survival and the beech-crop cycle. *Ardea* 88: 99–108.

Perkins, A., Whittingham, M., Bradbury, R., Wilson, J., Morris, A. & Barnett, P. (2000) Habitat characteristics affecting use of lowland agricultural grassland by birds in winter. *Biological Conservation* 95: 279–294.

Perkins, A.J., Maggs, H.E., Wilson, J.D., Watson, A. & Smout, C. (2008) Targeted management intervention reduces rate of population decline of Corn Buntings *Emberiza calandra* in eastern Scotland. *Bird Study* 55: 52–58.

Perkins, A.J., Maggs, H.E., Watson, A. & Wilson, J.D. (2011) Adaptive management and targeting of agri-environment schemes does benefit biodiversity: a case study of the Corn Bunting *Emberiza calandra*. *Journal of Applied Ecology* 48: 514–522.

Perkins, A.J., Watson, A., Maggs, H.E. & Wilson, J.D. (2012) Conservation insights from changing associations between habitat, territory distribution and mating system of Corn Buntings *Emberiza calandra* over a 20-year population decline. *Ibis* 154: 601–615. doi: [10.1111/j.1474-919X.2012.01246.x](https://doi.org/10.1111/j.1474-919X.2012.01246.x)

Perrins, C. (2003) The status of Marsh and Willow Tits in the UK. *British Birds* 96: 418–426.

Perrins, C.M. & Martin, P. (1999) *The impact of lost and discarded fishing line and tackle on Mute Swans*. Phase 1, R & D Technical Report W200. Environment Agency, Bristol.

Perrins, C.M. & Overall, R. (2001) Effect of increasing numbers of deer on bird populations in Wytham Woods, central England. *Forestry* 74: 299–309.

Perrins, C.M. & Sears, J. (1991) Collisions with overhead wires as a cause of mortality in Mute Swans *Cygnus olor*. *Wildfowl* 42: 5–11.

Pienkowski, M.W. (1991) Using long-term ornithological studies in setting targets for conservation in Britain. *Ibis* 133 (Suppl. 1): 62–75.

Potts, G. (1980) The effects of modern agriculture, nest predation and game management on the population ecology of partridges *Perdix perdix* and *Alectoris rufa*. *Advances in Ecological Research* 11: 2–79.

Potts, G.R. (2012) *Partridges*. Collins, London.

Potts, G. & Aebischer, N. (1995) Population dynamics of the Grey Partridge *Perdix perdix* 1793–1993: monitoring, modelling and management. *Ibis* 137: 29–37.

Prater, A.J. (1989) Ringed Plover *Charadrius hiaticula* breeding population of the United Kingdom in 1984. *Bird Study* 36: 154–159.

Prince, P. & Clarke, R. (1993) The hobby's breeding range in Britain. *British Wildlife* 4: 341–346.

Proffitt, F.M., Newton, I., Wilson, J.D. & Siriwardena, G.M. (2004) Bullfinch *Pyrrhula pyrrhula* breeding ecology in lowland farmland and woodland: comparisons across time and habitat. *Ibis* 146 (Suppl. 2): 78–86. doi: [10.1111/j.1474-919X.2004.00363.x](https://doi.org/10.1111/j.1474-919X.2004.00363.x)

Prytherch, R.J. (2013) The breeding biology of the Common Buzzard. *British Birds* 106: 264–279.

Rands, M. (1985) Pesticide use on cereals and the survival of Grey Partridge chicks: a field experiment. *Journal of Applied Ecology* 22: 49–54.

Ratcliffe, D.A. (1993) *The Peregrine Falcon*. Second edition. T. & A.D. Poyser, London.

Rebecca, G.W. (2011) Spatial and habitat-related influences on the breeding performance of Merlins in Britain. *British Birds* 104: 202–216.

Rebecca, G.W. & Bainbridge, I.P. (1998) The breeding status of the Merlin *Falco columbarius* in Britain in 1993–94. *Bird Study* 45: 172–187.

Redpath, S. & Thirgood, S. (1997) *Birds of Prey and Red Grouse* HMSO, London.

Redpath, S. & Thirgood, S. (1999) Numerical and functional responses in generalist predators: Hen Harriers and Peregrines on Scottish grouse moors. *Journal of Animal Ecology* 68: 879–892.

Redpath, S. & Thirgood, S. (2009) Hen harriers and red grouse: moving towards consensus? *Journal of Applied Ecology* 46: 961–963.

Redpath, S., Thirgood, S. & Clarke, R. (2002a) Field vole *Microtus agrestis* abundance and Hen Harrier *Circus cyaneus* diet and breeding in Scotland. *Ibis* 144: E33–E38.

Redpath, S., Arroyo, B., Etheridge, B., Leckie, F., Bouwman, K. & Thirgood, S. (2002b) Temperature and Hen Harrier productivity: from local mechanisms to geographical patterns. *Ecography* 25: 533–540.

Reed, T. (1985) Estimates of British breeding wader populations. *Wader Study Group Bulletin* 45: 11–12.

Reed, T.E., Jenouvrier, S. & Visser, M.E. (2012) Phenological mismatch strongly affects individual fitness but not population demography in a woodland passerine. *Journal of Animal Ecology*. doi: [10.1111/j.1365-2656.2012.02020.x](https://doi.org/10.1111/j.1365-2656.2012.02020.x)

Reed, T.E., Grøtan, V., Jenouvrier, S., Sæther, B.-E., & Visser, M.E. (2013) Population growth in a wild bird is buffered against phenological mismatch. *Science* 340 (6131): 488–491. doi: [10.1126/science.1232870](https://doi.org/10.1126/science.1232870)

Rehlfisch, M.M., Wernham, C.V. & Marchant, J.H. (eds) (1999) *Population, distribution, movements and survival of fish-eating birds in Great Britain*. DETR, London.

Rehlfisch, M.M., Austin, G.E., Holloway, S.J., Allan, J.R. & O'Connell, M. (2002) An approach to the assessment of change in the numbers of Canada Geese *Branta canadensis* and Greylag Geese *Anser anser* in southern Britain. *Bird Study* 49: 50–59.

Rhymer, C.M., Devereux, C.L., Denny, M.J.H. & Whittingham, M.J. (2012) Diet of Starling *Sturnus vulgaris* nestlings on farmland: the importance of Tipulidae larvae. *Bird Study* 59: 426–436.

Richner, H. (1992) The effect of extra food on fitness in breeding Carrion Crows. *Ecology* 73: 330–335.

Riddiford, N. (1983) Recent declines of Grasshopper Warblers *Locustella naevia* at British bird observatories. *Bird Study* 30: 143–148.

Risely, K., Baillie, S.R., Eaton, M.A., Joys, A.C., Musgrove, A.J., Noble, D.G., Renwick, A.R. & Wright, L.J. (2010) *The Breeding Bird Survey 2009*. Research Report 559. BTO, Thetford.

[Full text](#) (PDF, 1.17 MB)

Risely, K., Renwick, A.R., Dadam, D., Eaton, M.A., Johnston, A., Baillie, S.R., Musgrove, A.J. & Noble, D.G. (2011) *The Breeding Bird Survey 2010*. Research Report 597. British Trust for Ornithology, Thetford.

[Full text](#) (PDF, 1.59 MB)

Risely, K., Massimino, D., Johnston, A., Newson, S.E., Eaton, M.A., Musgrove, A.J., Noble, D.G., Procter, D. & Baillie, S.R. (2012) *The Breeding Bird Survey 2011*. Research Report 624. British Trust for Ornithology, Thetford.

[Full text](#) (PDF, 2.85 MB)

Risely, K., Massimino, D., Newson, S.E., Eaton, M.A., Musgrove, A.J., Noble, D.G., Procter, D. & Baillie, S.R. (2013) *The Breeding Bird Survey 2012*. Research Report 645. British Trust for Ornithology, Thetford.

[Full text](#) (PDF, 3.11 MB)

Robertson, D. (2003) Eurasian Reed Warblers in Scotland: a review of probable breeding records. *Scottish Birds* 24: 36–39.

Robertson, H.A. (1990) Breeding of Collared Doves *Streptopelia decaocto* in rural Oxfordshire, England. *Bird Study* 37: 73–83.

Robertson, P.A. (1991) Estimating the nesting success and productivity of British Pheasants *Phasianus colchicus* from nest-record schemes. *Bird Study* 38: 73–79.

Robertson, P.A., Woodburn, M.I.A., Tapper, S.C. & Stoate, C. (1989) *Estimating game densities in Britain from land-use maps*. Institute of Terrestrial Ecology, Grange-over-Sands.

Robinson, R.A., Wilson, J.D. & Crick, H.Q.P. (2001) The importance of arable habitat for farmland birds in grassland landscapes. *Journal of Applied Ecology* 38: 1059–1069.

Robinson, R.A., Siriwardena, G.M. & Crick, H.Q.P. (2002) Status and population trends of the Starling *Sturnus vulgaris* in Great Britain. In *Investigation into the causes of the decline of starlings and house sparrows in Great Britain* (eds Crick, H.Q.P., Robinson, R.A., Appleton, G.F., Clark, N.A. & Rickard, A.D.), pp 11–32. Research Report 290. BTO, Thetford.

Robinson, R.A., Crick, H.Q.P. & Peach, W.J. (2003) Population trends of Swallows *Hirundo rustica* breeding in Britain. *Bird Study* 50: 1–7.

doi: [10.1080/00063650309461283](https://doi.org/10.1080/00063650309461283)

Robinson, R.A., Green, R.E., Baillie, S.R., Peach, W.J. & Thomson, D.L. (2004) Demographic mechanisms of the population decline of the song thrush *Turdus philomelos* in Britain. *Journal of Animal Ecology* 73: 670–682. doi: [10.1111/j.0021-8790.2004.00841.x](https://doi.org/10.1111/j.0021-8790.2004.00841.x)

Robinson, R.A., Siriwardena, G.M. & Crick, H.Q.P. (2005a) Status and population trends of Starling *Sturnus vulgaris* in Great Britain. *Bird Study* 52: 252–260.

Robinson, R.A., Siriwardena, G.M. & Crick, H.Q.P. (2005b) Size and trends of the House Sparrow *Passer domesticus* population in Great Britain. *Ibis* 147: 552–562. doi: [10.1111/j.1474-919x.2005.00427.x](https://doi.org/10.1111/j.1474-919x.2005.00427.x)

Robinson, R.A., Freeman, S.N., Balmer, D.E. & Grantham, M.J. (2007a) Cetti's Warbler *Cettia cetti*: analysis of an expanding population. *Bird Study* 54: 230–235. doi: [10.1080/00063650709461479](https://doi.org/10.1080/00063650709461479)

Robinson, R.A., Baillie, S.R. & Crick, H.Q.P. (2007b) Weather-dependent survival: implications of climate change for passerine population processes. *Ibis* 149: 357–364.

Robinson, R.A., Balmer, D.E. & Marchant, J.H. (2008) Survival rates of hirundines in relation to British and African rainfall. *Ringing & Migration* 24: 1–6.

Robinson, R.A., Kew, J.J. & Kew, A.J. (2010a) Survival of suburban blackbirds *Turdus merula* varies seasonally but not by sex. *Journal of Avian Biology* 41: 83–87. doi: [10.1111/j.1600-048X.2009.04789.x](https://doi.org/10.1111/j.1600-048X.2009.04789.x)

Robinson, R.A., Lawson, B., Toms, M.P., Peck, K.M., Kirkwood, J.K., Chantrey, J., Clatworthy, I.R., Evans, A.D., Hughes, L.A., Hutchinson, O.C., John, S.K., Pennycott, T.W., Perkins, M.W., Rowley, P.S., Simpson, V.R., Tyler, K.M. & Cunningham, A.A. (2010b) Emerging infectious disease leads to rapid population declines of common British birds. *PLoS ONE* 5(8): e12215. [Abstract/Full text](#)

Robinson, R.A., Baillie, S.R. & King, R. (2012) Population processes in European Blackbirds *Turdus merula*: a state-space approach. *Journal of Ornithology* 152: 419–433. [Abstract](#)

Ronka, M.T.H., Saari, C.L.V., Lehtikainen, E.A., Suomela, J. & Hakkila, K. (2005) Environmental changes and population trends of breeding waterfowl in northern Baltic Sea. *Annales Zoologici Fennici* 42: 587–602.

Rossmann, E., Hontsch, K., Blaum, N. & Jeltsch, F. (2007) Reproductive success and nestling diet in the Lesser Spotted Woodpecker (*Picoides minor*): the early bird gets the caterpillar. *Journal of Ornithology* 148: 323–332.

Rowell, H.E. & Spray, C.J. (2004) *The Mute Swan Cygnus olor (Britain and Ireland populations) in Britain and Northern Ireland 1960/61 – 2000/01*. Waterbird Review Series, The Wildfowl & Wetlands Trust/Joint Nature Conservation Committee, Slimbridge.

Rubolini, D., Saino, N. & Møller, A.P. (2010) Migratory behaviour constrains the phenological response of birds to climate change. *Climate Research* 42: 45–55.

- Sage, R.B., Woodburn, M.I.A., Davis, C., Aebischer, N.J. (2002) The effect of an experimental infection of the nematode *Heterakis gallinarum* on hand-reared grey partridges *Perdix perdix*. *Parasitology* 124: 529–535.
- Saino, N., Szép, T., Ambrosini, R., Romano, M., & Møller, A.P. (2004) Ecological conditions during winter affect sexual selection and breeding in a migratory bird. *Proceedings of the Royal Society of London Series B* 271: 681–686. [Full text](#)
- Saino, N., Rubolini, D., Lehikoinen, E., Sokolov, L., Bonisoli-Alquati, A., Ambrosini, R., Boncoraglio, G. & Møller, A. (2009) Climate change effects on migration phenology may mismatch brood parasitic cuckoos and their hosts. *Biology Letters* 5: 539–541.
- Saino, N., Romano, M., Caprioli, M., Ambrosini, R., Rubolini, D., Scandolara, C. & Romano, A. (2012) A ptilochnological study of carry-over effects of conditions during wintering on breeding performance in the barn swallow *Hirundo rustica*. *Journal of Avian Biology* 43: 513–524. doi: [10.1111/j.1600-048X.2012.05622.x](#)
- Sálek, M. & Schröpfer, L. (2008) Population decline of the Little Owl (*Athene noctua* Scop.) in the Czech Republic. *Polish Journal of Ecology* 56: 527–534.
- Sanderson, F.J., Donald, P.F., Pain, D.J., Burfield, I.J. & van Bommel, F.P.J. (2006) Long-term population declines in Afro–Palearctic migrant birds *Biological Conservation* 131: 93–105.
- Sanz, J.J. (2002) Climate change and breeding parameters of Great and Blue Tits throughout the Western Palearctic. *Global Change Biology* 8: 409–422.
- SAS Institute (2009) SAS/STAT user's guide, version 9.2. SAS Inst. Inc., Cary, NC.
- SAS Institute Inc. (2010) *SAS 9.2 Language Reference: Dictionary, Third Edition*. SAS Inst. Inc., Cary, NC.
- Sauter, A., Korner-Nievergelt, F. & Jenni, L. (2010) Evidence of climate change effects on within-winter movements of European Mallards *Anas platyrhynchos*. *Ibis* 152: 600–609.
- Schaub, M., Ullrich, B., Knöttsch, G., Albrecht, P. & Meisser, C. (2006) Local population dynamics and the impact of scale and isolation: a study on different Little Owl populations. *Oikos* 115: 389–400.
- Schaub, M., Jakober, H. & Stauber, W. (2011) Demographic response to environmental variation in breeding, stopover and non-breeding areas in a migratory passerine. *Oecologia* 167: 445–459.
- Selas, V., Steen, R., Kobro, S., Lislevand, T. & Stenberg, I. (2008) Direct and indirect weather impacts on spring populations of Lesser Spotted Woodpecker (*Dendrocopos minor*) in Norway. *Scandinavian Journal of Forest Research* 23: 148–153.
- Sellers, R.M. (2006) Breeding population estimate for Northern Wheatear in Britain. *British Birds* 99: 533–535.
- Sergio, F., Boto, A., Scandolara, C. & Bogliani, G. (2002) Density, nest sites, diet, and productivity of Common Buzzards (*Buteo buteo*) in the Italian pre-Alps. *Journal of Raptor Research* 36: 24–32.
- Sergio, F., Scandolara, C., Marchesi, L., Pedrini, P. & Penteriani, V. (2005) Effect of agro-forestry and landscape changes on Common Buzzards (*Buteo buteo*) in the Alps: implications for conservation. *Animal Conservation* 8: 17–25.
- Setchfield, R.P., Mucklow, C., Davey, A., Bradter, U. & Anderson, G.Q.A. (2012) An agri-environment option boosts productivity of Corn Buntings *Emberiza calandra* in the UK. *Ibis* 154: 235–247. doi: [10.1111/j.1474-919X.2011.01207.x](#)
- Sharpe, F., Clark, J. & Leech, D. (2008) Does variation in demographic parameters account for regional variation in Northern Lapwing *Vanellus vanellus* population declines across Great Britain? *Bird Study* 55: 247–256.
- Shaw, L.M., Chamberlain, D.E. & Evans, M.R. (2008) The house sparrow *Passer domesticus* in urban areas: reviewing a possible link between post-decline distribution and human socioeconomic status. *Journal of Ornithology* 149: 293–299. [Abstract](#) doi: [10.1007/s10336-008-0285-y](#)
- Shaw, L.M., Chamberlain, D., Conway, G. & Toms, M. (2011) *Spatial distribution and habitat preferences of the House Sparrow Passer domesticus in urbanised landscapes*. Research Report 599. BTO, Thetford.
- [Full text](#) (PDF, 628.20 KB)
- Shortall, C.R., Moore, A., Smith, E., Hall, M.J., Woiod, I.P. & Harrington, R. (2009) Long-term changes in the abundance of flying insects *Insect Conservation and Diversity* 2: 251–260. doi: [10.1111/j.1752-4598.2009.00062.x](#)
- Shrubbs, M. (1990) Effects of agricultural change on nesting Lapwings *Vanellus vanellus* in England and Wales. *Bird Study* 37: 115–127.
- Shrubbs, M., Lack, P. & Greenwood, J. (1991) The numbers and distribution of Lapwings *Vanellus vanellus* nesting in England and Wales in 1987. *Bird Study* 38: 20–37.
- Sim, I.M.W., Campbell, L., Pain, D.J. & Wilson, J.D. (2000) Correlates of the population increase of Common Buzzards *Buteo buteo* in the West Midlands between 1983 and 1996. *Bird Study* 47: 154–164.
- Sim, I.M.W., Cross, A.V., Lamacraft, D.L. & Pain, D.J. (2001a) Correlates of Common Buzzard *Buteo buteo* density and breeding success in the West Midlands. *Bird Study* 48: 317–329.
- Sim, I.M.W., Gibbons, D.W., Bainbridge, I.P. & Mattingley, W.A. (2001b) Status of the Hen Harrier *Circus cyaneus* in the UK and the Isle of Man in 1998. *Bird Study* 48: 341–353.
- Sim, I.M.W., Dillon, I.A., Eaton, M.A., Etheridge, B., Lindley, P., Riley, H., Saunders, R., Sharpe, C. & Tickner, M. (2007a) Status of the Hen Harrier *Circus cyaneus* in the UK and Isle of Man in 2004, and a comparison with the 1988/89 and 1998 surveys. *Bird Study* 54: 256–267.

- Sim, I.M.W., Burfield, I.J., Grant, M.C., Pearce-Higgins, J.W. & Brooke, M. de L. (2007b) The role of habitat composition in determining breeding site occupancy in a declining Ring Ouzel *Turdus torquatus* population. *Ibis* 149: 374–385.
- Sim, I., Rollic, C., Arthur, D., Benn, S., Booker, H., Fairbrother, V., Green, M., Hutchinson, K., Ludwig, S., Nicoll, M., Poxton, I., Rebecca, G., Smith, L., Stanbury, A. & Wilson, P. (2010) The decline of the Ring Ouzel in Britain. *British Birds* 103, 229–239.
- Sim, I.M.W., Rebecca, G.W., Ludwig, S.C., Grant, M.C. & Reid, J.M. (2011) Characterizing demographic variation and contributions to population growth rate in a declining population. *Journal of Animal Ecology* 80: 159–170.
- Simms, E. (1989) *The Song Thrush*. Shire Natural History, Aylesbury.
- Siriwardena, G.M. (2004) Possible roles of habitat, competition and avian nest predation in the decline of the Willow Tit *Parus montanus* in Britain. *Bird Study* 51: 193–202. doi: [10.1080/00063650409461354](https://doi.org/10.1080/00063650409461354)
- Siriwardena, G.M. (2006) Avian nest predation, competition and the decline of British Marsh Tits *Parus palustris*. *Ibis* 148: 255–265. doi: [10.1111/j.1474-919X.2006.00525.x](https://doi.org/10.1111/j.1474-919X.2006.00525.x)
- Siriwardena, G.M. & Stevens, D.K. (2004) Effects of habitat on the use of supplementary food by farmland birds in winter *Ibis* 146: 144–154.
- Siriwardena, G.M., Baillie, S.R., Buckland, S.T., Fewster, R.M., Marchant, J.H. & Wilson, J.D. (1998a) Trends in the abundance of farmland birds: a quantitative comparison of smoothed Common Birds Census indices. *Journal of Applied Ecology* 35: 24–43. [Abstract/Full text](#)
- Siriwardena, G.M., Baillie, S.R. & Wilson, J.D. (1998b) Variation in the survival rates of British passerines with respect to their population trends on farmland *Bird Study* 45: 276–292.
- Siriwardena, G.M., Baillie, S.R. & Wilson, J.D. (1999) Temporal variation in the annual survival rates of six granivorous birds with contrasting population trends *Ibis* 141: 621–636.
- Siriwardena, G.M., Baillie, S.R., Crick, H.Q.P., Wilson, J.D. & Gates, S. (2000a) The demography of lowland farmland birds. In *Proceedings of the 1999 BOU Spring Conference: Ecology and Conservation of Lowland Farmland Birds* (eds N.J. Aebischer, A.D. Evans, P.V. Grice & J.A. Vickery), pp 117–133. British Ornithologists' Union, Tring.
- Siriwardena, G.M., Baillie, S.R., Crick, H.Q.P. & Wilson, J.D. (2000b) The importance of variation in the breeding performance of seed-eating birds for their population trends on farmland. *Journal of Applied Ecology* 37: 128–148. [Abstract/Full text](#)
- Siriwardena, G.M., Crick, H.Q.P., Baillie, S.R. & Wilson, J.D. (2000c) Agricultural habitat-type and the breeding performance of granivorous farmland birds in Britain *Bird Study* 47: 66–81.
- Siriwardena, G.M., Freeman, S.N. & Crick, H.Q.P. (2001a) The decline of the Bullfinch *Pyrrhula pyrrhula* in Britain: is the mechanism known? *Acta Ornithologica* 36: 143–152.
- Siriwardena, G., Baillie, S., Crick, H. & Wilson, J. (2001b) Changes in agricultural land-use and breeding performance of some granivorous farmland passerines in Britain. *Agriculture, Ecosystems & Environment* 84: 191–206.
- Siriwardena, G.M., Wilson, J.D., Baillie, S.R. & Crick, H.Q.P. (2001c) Can the historical CBC trend for Skylark be 'recovered' using present-day agricultural habitat preferences and changes in agricultural land use? In *The Ecology and Conservation of Skylarks* (eds Donald, P.F. & Vickery, J.A.), 56–60. RSPB, Sandy.
- Siriwardena, G.M., Robinson, R.A. & Crick, H.Q.P. (2002) Status and population trends of the house sparrow *Passer domesticus* in Great Britain. In *Investigation into the causes of the decline of starlings and house sparrows in Great Britain* (eds Crick, H.Q.P., Robinson, R.A., Appleton, G.F., Clark, N.A. & Rickard, A.D.), pp 33–52. Research Report 290. BTO, Thetford.
- Siriwardena, G.M., Stevens, D.K., Anderson, G.Q.A., Vickery, J.A., Calbrade, N.A. & Dodd, S. (2007) The effect of supplementary winter seed food on breeding populations of farmland birds: evidence from two large-scale experiments. *Journal of Applied Ecology* 44: 920–932.
- Sitters, H.P., Fuller, R.J., Hoblyn, R.A., Wright, M.T., Cowie, N. & Bowden, C.G.R. (1996) The Woodlark *Lullula arborea* in Britain: population trends, distribution and habitat occupancy. *Bird Study* 43: 172–187.
- Smart, J., Taylor, E., Amar, A., Smith, K., Bierman, S., Carpenter, J., Grice, P., Currie, F., Smithers, R., Fuller, R. & Hewson, C. (2007) *Habitat associations of woodland birds: implications for woodland management for declining species*. Research Report 26. RSPB, Sandy. [Full text](#)
- Smart, J., Amar, A., O'Brien, M., Grice, P. & Smith, K. (2008) Changing land management of lowland wet grasslands of the UK: impacts on snipe abundance and habitat quality. *Animal Conservation* 11: 339–351.
- Smart, J., Amar, A., Sim, I.M.W., Etheridge, B., Cameron, D., Christie, G. & Wilson, J.D. (2010) Illegal killing slows population recovery of a re-introduced raptor of high conservation concern – the Red Kite *Milvus milvus*. *Biological Conservation* 143: 1278–1286. doi: [10.1016/j.biocons.2010.03.002](https://doi.org/10.1016/j.biocons.2010.03.002)
- Smart, J., Bolton, M., Hunter, F., Quayle, H., Thomas, G. & Gregory, R.D. (2013) Managing uplands for biodiversity: do agri-environment schemes deliver benefits for breeding lapwing *Vanellus vanellus*? *Journal of Applied Ecology* 50: 794–804. doi: [10.1111/1365-2664.12081](https://doi.org/10.1111/1365-2664.12081)
- Smith, K.W. (1997) Nest site selection of the great spotted woodpecker *Dendrocopos major* in two oak woods in southern England and its implications for woodland management. *Biological Conservation* 80: 283–288.
- Smith, K.W. (2005) Has the reduction in nest-site competition from Starlings *Sturnus vulgaris* been a factor in the recent increase of Great Spotted Woodpecker *Dendrocopos major* numbers in Britain? *Bird Study* 52: 307–313.
- Smith, K.W. (2006) The implications of nest site competition from starlings *Sturnus vulgaris* and the effect of spring temperatures on the timing and breeding performance of great spotted woodpeckers *Dendrocopos major* in southern England. *Annales Zoologici Fennici* 43: 177–185. [Full text](#)

- Smith, K.W. (2007) The utilization of dead wood resources by woodpeckers in Britain. *Ibis* 149: 183–192.
- Smith, K.W. & Charman, E.C. (2012) The ecology and conservation of the Lesser Spotted Woodpecker. *British Birds* 105: 294–307.
- Smith, K.W. & Smith, L. (2013) The effect of supplementary feeding in early spring on the breeding performance of the Great Spotted Woodpecker *Dendrocopos major*. *Bird Study* 60: 169–175.
- Snow, D.W. (1965) The relationship between census results and the breeding population of birds on farmland. *Bird Study* 12: 287–304.
- Soler, M. & Soler, J.J. (1996) Effects of experimental food provisioning on reproduction in the Jackdaw *Corvus monedula*, a semi-colonial species. *Ibis* 138: 377–383.
- Sotherton, N., Robertson, P.A. & Dowell, S. (1993) Manipulating pesticide use to increase the production of wild game birds in Britain. In *Quail III: National Quail Symposium* (eds Church, K.E. & Dailey, T.V.), 92–101. Kansas Department of Wildlife and Parks, Pratt.
- Sotherton, N.W., Aebischer, N.J. & Ewald, J.A. (2014) Research into action: grey partridge conservation as a case study. *Journal of Applied Ecology* 51: 1–5. doi: [10.1111/1365-2664.12162](https://doi.org/10.1111/1365-2664.12162)
- Spray, C.J. (1981) An isolated population of *Cygnus olor* in Scotland. In *Proceedings of 2nd International Swan Symposium, Sapporo, Japan 1980* (eds Matthews, G.V.T. & Smart, M.), 191–203. International Waterfowl Research Bureau, Slimbridge.
- Steen, R., Selas, V. & Stenberg, I. (2006) Impact of weather on annual fluctuations in breeding numbers of Lesser Spotted Woodpecker *Dendrocopos minor* in Norway. *Ardea* 94: 225–231.
- Steven, R., Pickering, C. & Castley, J.G. (2011) A review of the impacts of nature-based recreation on birds. *Journal of Environmental Management* 92: 2287–2294. doi: [10.1016/j.jenvman.2011.05.005](https://doi.org/10.1016/j.jenvman.2011.05.005)
- Stevens, D.K., Anderson, G.Q.A., Grice, P.V. & Norris, K. (2007) Breeding success of Spotted Flycatchers *Muscicapa striata* in southern England – is woodland a good habitat for this species? *Ibis* 149: 214–223.
- Stevens, D.K., Anderson, G.Q.A., Grice, P.V., Norris, K. & Butcher, N. (2008) Predators of Spotted Flycatcher *Muscicapa striata* nests in southern England as determined by digital nest-cameras. *Bird Study* 55: 179–187.
- Stevens, M., Sheehan, D., Wilson, J., Buchanan, G. & Cresswell, W. (2010) Changes in Sahelian bird biodiversity and tree density over a five-year period in northern Nigeria. *Bird Study* 57: 156–174.
- Stoate, C. & Szczur, J. (2001) Could game management have a role in the conservation of farmland passerines? A case study from a Leicestershire farm. *Bird Study* 48: 279–292.
- Stoate, C. & Szczur, J. (2006) Potential influence of habitat and predation on local breeding success and population in Spotted Flycatchers *Muscicapa striata*. *Bird Study* 53: 328–330.
- Stoddart, A. (2013) Redpolls: a review of their taxonomy, identification and British status. *British Birds* 106: 708–736.
- Stone, B.H., Sears, J., Cranswick, P.A., Gregory, R.D., Gibbons, D.W., Rehfish, M.M., Aebischer, N.J. & Reid, J.B. (1997) Population estimates of birds in Britain and in the United Kingdom. *British Birds* 90: 1–22.
- Stroud, D.A., Reed, T.M., Pienkowski, M.W. & Lindsay, R.A. (1987) *Birds, Bogs and Forestry*. Nature Conservancy Council, Peterborough.
- Strubbe, D. & Matthysen, E. (2007) Invasive ring-necked parakeets *Psittacula krameri* in Belgium: habitat selection and impact on native birds. *Ecography* 30: 578–588.
- Strubbe, D. & Matthysen, E. (2009) Experimental evidence for nest-site competition between invasive Ring-necked Parakeets (*Psittacula krameri*) and native Nuthatches (*Sitta europaea*). *Biological Conservation* 142: 1588–1594.
- Strubbe, D., Matthysen, E. & Graham, C.H. (2010) Assessing the potential impact of invasive ring-necked parakeets *Psittacula krameri* on native nuthatches *Sitta europaea* in Belgium. *Journal of Applied Ecology* 47: 549–557. doi: [10.1111/j.1365-2664.2010.01808.x](https://doi.org/10.1111/j.1365-2664.2010.01808.x)
- Summers, R. (1998) The decline in clutch size of Hen Harriers. *BTO News* 218: 23.
- Summers, R.W. (1999) Numerical responses of crossbills *Loxia* spp. to annual fluctuations in cone crops. *Ornis Fennica* 76: 141–144.
- Summers, R.W. & Buckland, S.T. (2010) A first survey of the global population size and distribution of the Scottish Crossbill *Loxia scotica*. *Bird Conservation International*. Available on CJO 11 Jan 2010. doi: [10.1017/S0959270909990323](https://doi.org/10.1017/S0959270909990323)
- Summers-Smith, J.D. (1989) A history of the status of the Tree Sparrow *Passer montanus* in the British Isles. *Bird Study* 36: 23–31.
- Summers-Smith, J.D. (2003) The decline of the House Sparrow: a review. *British Birds* 96: 439–446.
- Summers-Smith, J.D. (2007) Is unleaded petrol a factor in urban House Sparrow decline? *British Birds* 100: 558–559.
- Swann, R.L. & Etheridge, B. (1995) A comparison of breeding success and prey of the Common Buzzard *Buteo buteo* in two areas of northern Scotland. *Bird Study* 42: 37–43.
- Szép, T. (1995) Relationship between west African rainfall and the survival of central European Sand Martins *Riparia riparia*. *Ibis* 137: 162–168.
- Tapper, S. (1992) *Game Heritage: an ecological review from shooting and gamekeeping records*. Game Conservancy, Fordingbridge.

- Tapper, S. (1999) *A question of balance: game animals and their role in the British countryside*. The Game Conservancy Trust, Hampshire, UK.
- Tapper, S. & France, J. (1992) The National Game Bag Census 1991. *Game Conservancy Annual Review* 23: 38–40.
- Tapper, S.C., Potts, G.R. & Brockless, M.H. (1996) The effect of an experimental reduction in predation pressure on the breeding success and population density of grey partridges (*Perdix perdix*). *Journal of Applied Ecology* 33: 965–978.
- Taylor, A.J. & O'Halloran, J. (2002) The decline of the Corn Bunting *Miliaria calandra* in the Republic of Ireland. *Biology and Environment: Proceedings of the Royal Irish Academy* 102: 165–175.
- Taylor, I. & Grant, M. (2004) Long-term trends in the abundance of breeding Lapwing *Vanellus vanellus* in relation to land-use change on upland farmland in southern Scotland. *Bird Study* 51: 133–142.
- Temple, S.A. & Wiens, J.A. (1989) Bird populations and environmental changes: can birds be bio-indicators? *American Birds* 43: 260–270.
- Teunissen, W., Schekkerman, H., Willems, F. & Majoor, F. (2008) Identifying predators of eggs and chicks of Lapwing *Vanellus vanellus* and Black-tailed Godwit *Limosa limosa* in the Netherlands and the importance of predation on wader reproductive output. *Ibis* 150: 74–85.
- Thackeray, S.J., Sparks, T.H., Frederiksen, M., Burthe, S., Bacon, P.J., Bell, J.R., Botham, M.S., Brereton, T.M., Bright, P.W., Carvalho, L., Clutton-Brock, T., Dawson, A., Edwards, M., Elliott, J.M., Harrington, R., Johns, D., Jones, I.D., Jones, J.T., Leech, D.I., Roy, D.B., Scott, W.A., Smith, M., Smithers, R.J., Winfield, I.J. & Wanless, S. (2010) Trophic level asynchrony in rates of phenological change for marine, freshwater and terrestrial environments. *Global Change Biology* 16: 3304–3313. doi: [10.1111/j.1365-2486.2010.02165.x](https://doi.org/10.1111/j.1365-2486.2010.02165.x)
- Tharme, A.P., Green, R.E., Baines, D., Bainbridge, I.P. & O'Brien, M. (2001) The effect of management for Red Grouse shooting on the population density of breeding birds on heather-dominated moorland. *Journal of Applied Ecology* 38: 439–457.
- Thaxter, C.B., Redfern, C.P.F. & Bevan, R.M. (2006) Survival rates of adult Reed Warblers *Acrocephalus scirpaceus* at a northern and southern site in England. *Ringing & Migration* 23: 65–79.
- Thaxter, C.B., Sansom, A., Thewlis, R.M., Calbrade, N.A., Ross-Smith, V.H., Bailey, S., Mellan, H.J. & Austin, G.E. (2010) Wetland Bird Survey Alerts 2006/2007: changes in numbers of wintering waterbirds in the constituent countries of the United Kingdom, Special Protection Areas (SPAs) and Sites of Special Scientific Interest (SSSIs). Research Report 556. BTO, Thetford. [Full text](#)
- Thingstad, P.G., Nyholm, N.E.I. & Fjeldheim, B. (2006) Pied Flycatcher *Ficedula hypoleuca* population dynamics in peripheral habitats in Scandinavia. *Ardea* 94: 211–223.
- Thomas, G.E. (1993) Estimating annual total heron population counts. *Applied Statistics* 42: 473–486.
- Thompson, P.S., Amar, A., Hoccom, D.G., Knott, J. & Wilson, J.D. (2009) Resolving the conflict between driven-grouse shooting and conservation of hen harriers. *Journal of Applied Ecology* 46: 950–954.
- Thomson, D.L., Baillie, S.R. & Peach, W.J. (1997) The demography and age-specific annual survival of British song thrushes *Turdus philomelos* during periods of population stability and decline. *Journal of Animal Ecology* 66: 414–424.
- Thomson, D.L., Green, R.E., Gregory, R.D. & Baillie, S.R. (1998) The widespread declines of songbirds in rural Britain do not correlate with the spread of their avian predators. *Proceedings of the Royal Society of London Series B* 265: 2057–2062. [Full text](#)
- Thorup, K., Sunde, P., Jacobsen, L.B. & Rahbek, C. (2010) Breeding season food limitation drives population decline of the Little Owl *Athene noctua* in Denmark. *Ibis* 152: 803–814.
- Tompkins, D.M., Draycott, R.A.H. & Hudson, P.J. (2000) Field evidence for apparent competition mediated via the shared parasites of two gamebird species *Ecology Letters* 3: 10–14.
- Tompkins, D., Greenman, J., Robertson, P. & Hudson, P. (2000b) The role of shared parasites in the exclusion of wildlife hosts *Heterakis gallinarum* in the Ring-necked Pheasant and the Grey Partridge. *Journal of Animal Ecology* 69: 829–840.
- Toms, M.P., Crick, H.Q.P. & Shawyer, C.R. (2000) *Project Barn Owl Final Report*. BTO Research Report 197/ HOT Research Report 98/1. BTO/Hawk & Owl Trust, Thetford.
- Toms, M.P., Crick, H.Q.P. & Shawyer, C.R. (2001) The status of breeding Barn Owls *Tyto alba* in the United Kingdom 1995–97. *Bird Study* 48: 23–37.
- Tryjanowski, P., Sparks, T.H. & Crick, H.Q.P. (2006) Red-backed Shrike (*Lanius collurio*) nest performance in a declining British population: a comparison with a stable population in Poland. *Ornis Fennica* 83: 181–186.
- Tucker, G.M. & Heath, M.F. (1994) *Birds in Europe: their conservation status*. Conservation Series no. 3. BirdLife International, Cambridge.
- Turner, A. (2009) Climate change: a Swallow's eye view. *British Birds* 102: 3–16.
- Vanhinsbergh, D., Fuller, R. & Noble, D. (2003) *A review of possible causes of recent changes in populations of woodland birds in Britain*. Research Report 245. BTO, Thetford.
- Verhulst, S. (1992) Effects of density, beech crop and winter feeding on survival of juvenile Great Tits: an analysis of Kluyver's removal experiment *Ardea* 80: 285–292.
- Vickery, J.A. (1991) Breeding density of Dippers *Cinclus cinclus*, Grey Wagtails *Motacilla cinerea* and Common Sandpipers *Actitis hypoleucos* in relation to the acidity of streams in south-west Scotland. *Ibis* 133: 178–185.

- Vickery, J.A. (1992) The reproductive success of the Dipper *Cinclus cinclus* in relation to the acidity of streams in south-west Scotland. *Freshwater Biology* 28: 195–205.
- Vickery, J.A., Tallowin, J.T., Feber, R.E., Asteraki, E.A., Atkinson, P.W., Fuller, R.J. & Brown, V.K. (2001) The management of lowland neutral grasslands in Britain: effects of agricultural practices on birds and their food resources. *Journal of Applied Ecology* 38: 647–664.
- Vickery, J.A., Bradbury, R.B., Henderson, I.G., Eaton, M.A. & Grice, P.V. (2004) The role of agri-environment schemes and farm management practices in reversing the decline of farmland birds in England. *Biological Conservation* 119: 19–39.
- Vincent, K.E. (2005) *Investigating the causes of the decline of the urban House Sparrow* *Passer domesticus* population in Britain. Unpublished PhD thesis, De Montfort University. [Full text](#)
- Visser, M.E. & Both, C. (2005) Shifts in phenology due to global climate change: the need for a yardstick. *Proceedings of the Royal Society B* 272: 2561–2569. [Full text](#)
- Visser, M.E. & Holleman, L.J.M. (2001) Warmer springs disrupt the synchrony of oak and winter moth phenology. *Proceedings of the Royal Society of London Series B* 268: 289–294. [Full text](#)
- Visser, M.E., van Noordwijk, A.J., Tinbergen, J.M. & Lessells, C.M. (1998) Warmer springs lead to mistimed reproduction in Great Tits (*Parus major*). *Proceedings of the Royal Society of London Series B* 265: 1867–1870. [Full text](#)
- Visser, M.E., Holleman, L.J.M. & Caro, S.P. (2009) Temperature has a causal effect on avian timing of reproduction. *Proceedings of the Royal Society B Biological Sciences* 276: 2323–2331.
- Vögeli, M., Laiolo, P., Serrano, D. & Tella, J.L. (2011) Predation of experimental nests is linked to local population dynamics in a fragmented bird population. *Biology Letters* 7: 954–957.
- Walsh, P.M., Brindley, E. & Heubeck, M. (1995) *Seabird numbers and breeding success in Britain and Ireland, 1994*. JNCC, Peterborough.
- Wanless, S., Harris, M.P., Redman, P. & Speakman, J.R. (2005) Low energy values of fish as a probable cause of a major seabird breeding failure in the North Sea. *Marine Ecology Progress Series* 294: 1–8.
- Ward, R.M., Cranswick, P.A., Kershaw, M., Austin, G.E., Brown, A.W., Brown, L.M., Coleman, J.T., Chisholm, H.K. & Spray, C.J. (2007) Numbers of Mute Swans *Cygnus olor* in Great Britain: results of the national census in 2002. *Wildfowl* 57: 3–20.
- Watson, A., Perkins, A.J., Maggs, H.E. & Wilson, J.D. (2009) Decline of Corn Buntings *Emberiza calandra* on east Scottish study areas in 1989–2007. *Bird Study* 56: 213–220.
- Watson, M., Aebischer, N.J., Potts, G.R. & Ewald, J.A. (2007) The relative effects of raptor predation and shooting on overwinter mortality of grey partridges in the United Kingdom. *Journal of Applied Ecology* 44: 972–982.
- Wesolowski, T. (1985) The breeding ecology of the Wood Warbler *Phylloscopus sibilatrix* in primeval forest. *Ornis Scandinavica* 16: 49–60.
- Wesolowski, T. & Maziarz, M. (2009) Changes in breeding phenology and performance of Wood Warblers *Phylloscopus sibilatrix* in a primeval forest: a thirty-year perspective. *Acta Ornithologica* 44: 69–80.
- Wesolowski, T. & Stawarczyk, T. (1991) Survival and population dynamics of Nuthatches *Sitta europaea* breeding in natural cavities in a primeval temperate forest. *Ornis Scandinavica* 22: 143–154.
- Wesolowski, T. & Tomialojc, L. (1986) The breeding ecology of woodpeckers in a temperate primeval forest – preliminary data. *Acta Ornithologica* 22: 1–21.
- Whitfield, D., Fielding, A. & Whitehead, S. (2008) Long-term increase in the fecundity of Hen Harriers in Wales is explained by reduced human interference and warmer weather. *Animal Conservation* 11: 144–152.
- Whittingham, M.J., Swetnam, R.D., Wilson, J.D., Chamberlain, D.E. & Freckleton, R.P. (2005) Habitat selection by yellowhammers *Emberiza citrinella* on lowland farmland at two spatial scales: implications for conservation management. *Journal of Applied Ecology* 42: 270–290. [Abstract/Full text](#)
- Wiklander, U., Nilsson, I., Nilsson, S., Olsson, O., Pettersson, B. & Stagen, A. (1992) Occurrence of the Lesser Spotted Woodpecker *Dendrocopos minor* in relation to area of deciduous forest. *Ornis Fennica* 69: 113–118.
- Wiklander, U., Olsson, O. & Nilsson, S. G. (2001) Annual and seasonal reproductive trends in the Lesser Spotted Woodpecker *Dendrocopos minor*. *Ibis* 143: 72–82.
- Wilson, A.M. & Vickery, J.A. (2005) Decline in Yellow Wagtail *Motacilla flava flavissima* breeding on lowland wet grassland in England and Wales between 1982 and 2002. *Bird Study* 52: 88–92.
- Wilson, A.M., Marchant, J.H., Gregory, R.D., Siriwardena, G.M. & Baillie, S.R. (1998) *Enhancements for monitoring of opportunistic bird populations*. Research Report 200. BTO, Thetford.
- [Full text](#) (PDF, 1.66 MB)
- Wilson, A.M., Vickery, J.A. & Browne, S.J. (2001) Numbers and distribution of Northern Lapwings *Vanellus vanellus* breeding in England and Wales in 1998. *Bird Study* 48: 2–17.
- Wilson, A.M., Henderson, A.C.B. & Fuller, R.J. (2002) Status of the Common Nightingale *Luscinia megarhynchos* in England at the end of the 20th century with particular reference to climate change. *Bird Study* 49: 193–204. doi: [10.1080/00063650209461266](https://doi.org/10.1080/00063650209461266)
- Wilson, A.M., Ausden, M. & Milsom, T.P. (2004) Changes in breeding wader populations on lowland wet grasslands in England and Wales: causes and potential solutions. *Ibis* 146: 32–40.

- Wilson, A.M., Vickery, J.A., Brown, A., Langston, R.H.W., Smallshire, D., Wotton, S. & Vanhinsbergh, D. (2005) Changes in the numbers of breeding waders on lowland wet grasslands in England and Wales between 1982 and 2002. *Bird Study* 52: 55–69. doi: [10.1080/00063650509461374](https://doi.org/10.1080/00063650509461374)
- Wilson, J.D. (2001) Weeds as a food resource for farmland birds: what, where and how many should we leave? *BCPC Conference – Weeds 2001*, 1–2: 391–398.
- Wilson, J.D., Taylor, R. & Muirhead, L.B. (1996) Field use by farmland birds in winter: an analysis of field type preferences using resampling methods *Bird Study* 43: 320–332.
- Wilson, J.D., Evans, J., Browne, S.J., & King, J.R. (1997) Territory distribution and breeding success of skylarks *Alauda arvensis* on organic and intensive farmland in Southern England. *Journal of Applied Ecology* 34: 1462–1478. [Abstract](#)
- Wilson, J.D., Morris, A.J., Arroyo, B.E., Clark, S.C. & Bradbury, R.B. (1999) A review of the abundance and diversity of invertebrate and plant foods of granivorous birds in northern Europe in relation to agricultural change. *Agriculture, Ecosystems & Environment* 75: 13–30.
- Wilson, J.D., Boyle, J., Jackson, D.B., Lowe, B. & Wilkinson, N.I. (2007) Effect of cereal harvesting method on a recent population decline of Corn Buntings *Emberiza calandra* on the Western Isles of Scotland. *Bird Study* 54: 362–370.
- Wilson, J.M. & Cresswell, W. (2006) How robust are Palearctic migrants to habitat loss and degradation in the Sahel? *Ibis* 148: 789–800.
- Winstanley, D., Spencer, R. & Williamson, K. (1974) Where have all the Whitethroats gone? *Bird Study* 21: 1–14.
- Wotton, S.R. & Gillings, S. (2000) The status of breeding Woodlarks *Lullula arborea* in Britain in 1997. *Bird Study* 47: 212–224. doi: [10.1080/00063650009461176](https://doi.org/10.1080/00063650009461176)
- Wotton, S., Gibbons, D.W., Dilger, M. & Grice, P.V. (1998) Cetti's Warblers in the United Kingdom and the Channel Islands in 1996 *British Birds* 91: 77–89.
- Wotton, S.R., Langston, R.H.W. & Gregory, R.D. (2002a) The breeding status of the Ring Ouzel *Turdus torquatus* in the UK in 1999. *Bird Study* 49: 26–34.
- Wotton, S.R., Carter, I., Cross, A.V., Etheridge, B., Snell, N., Duffy, K., Thorpe, R. & Gregory, R.D. (2002b) Breeding status of the Red Kite *Milvus milvus* in Britain in 2000. *Bird Study* 49: 278–286.
- Wright, L.J., Hoblyn, R.A., Sutherland, W.J. & Dolman, P.M. (2007) Reproductive success of Woodlarks *Lullula arborea* in traditional and recently colonized habitats. *Bird Study* 54: 315–323. doi: [10.1080/00063650709461491](https://doi.org/10.1080/00063650709461491)
- Wright, L.J., Hoblyn, R.A., Green, R.E., Bowden, C.G.R., Mallord, J.W., Sutherland, W.J. & Dolman, P.M. (2009) Importance of climatic and environmental change in the demography of a multi-brooded passerine, the Woodlark *Lullula arborea*. *Journal of Animal Ecology* 78: 1191–1202. doi: [10.1111/j.1365-2656.2009.01582.x](https://doi.org/10.1111/j.1365-2656.2009.01582.x)
- Wright, P.M. (2005) *Merlins of the south-east Yorkshire Dales*. Tarnmoor, Skipton.
- Wyllie, I. & Newton, I. (1991) Demography of an increasing population of Sparrowhawks. *Journal of Animal Ecology* 60: 749–766.
- Yom-Tov, Y. (1974) The effect of food and predation on breeding density and success, clutch size and laying date of the crow (*Corvus corone* L.). *Journal of Animal Ecology* 43: 479–498.
- Zmihorski, M., Altenburg-Bacia, D., Romanowski, J., Kowalski, M. & Osojca, G. (2006) Long-term decline of the little owl (*Athene noctua* Scop., 1769) in central Poland. *Polish Journal of Ecology* 54: 321–324.

Mute Swan

Cygnus olor

Key facts


Conservation listings:	Europe: no SPEC category (concentrated in Europe, conservation status favourable) (BiE04) UK: green (BoCC3)
Long-term trend:	UK, England: rapid increase
Population size:	6,400 (5,800-7,000) pairs in 2009 (APEP13: 2002 estimate (Ward et al. 2007) updated using BBS trend); 79,000 individuals in winter in 2004-09 (Musgrove et al. 2011)
Migrant status:	Resident
Nesting habitat:	Ground nester
Primary breeding habitat:	Wetland
Secondary breeding habitat:	
Breeding diet:	Vegetation
Winter diet:	Vegetation

Status summary

Mute Swan populations, which had been fairly stable since the 1960s, increased progressively from the mid 1980s to around 2000, when a new plateau was reached. Waterways, likely to be a preferred habitat for breeding swans, show a more moderate rate of increase than CBC/BBS. Winter trends as measured by WeBS have shown a parallel upturn, with little change in Britain after 2000 (Austin et al. 2014). After a spell on the amber list during 2002-09, for reasons unconnected with its UK trend, the species is now green listed once more. There has been widespread moderate increase across Europe since 1980 (PECBMS 2013a).

CBC/BBS UK 1966-2012

Mute Swan







Smoothed population index, relative to an arbitrary 100 in the year given, with 85% confidence limits in green

Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	108	240	47	642		
	25	1986-2011	176	124	66	186		
	10	2001-2011	288	0	-12	10		
	5	2006-2011	320	10	0	19		
CBC/BBS England	44	1967-2011	93	209	45	611		Small CBC sample
	25	1986-2011	151	112	57	197		
	10	2001-2011	244	2	-9	19		
	5	2006-2011	274	5	-5	14		
WBS/WBBS waterways	36	1975-2011	79	88	36	187		
	25	1986-2011	98	65	25	112		
	10	2001-2011	144	1	-19	19		
	5	2006-2011	134	-7	-23	6		
BBS UK	16	1995-2011	246	27	4	57		
	10	2001-2011	287	0	-12	10		
	5	2006-2011	318	10	1	17		
BBS England	16	1995-2011	210	11	-6	36		
	10	2001-2011	244	3	-9	17		
	5	2006-2011	272	5	-6	13		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.






BBS UK graph



BBS England graph

Demographic trends


Fledglings per breeding attempt Mute Swan 1966-2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Mute Swan




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	34	None					
Clutch size	43	1968-2011	32	None					
Brood size	43	1968-2011	67	Curvilinear	4.42 chicks	4.50 chicks	1.8%		
Nest failure rate at egg stage	43	1968-2011	38	None					
Nest failure rate at chick stage	43	1968-2011	42	None					
Laying date	43	1968-2011	17	None			0 days		Small sample


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

Causes of change

The increase in this species has been attributed to the banning of lead weights for fishing and the positive implications of this on survival. Milder winters have also been a factor, increasing overwinter survival and having knock-on effects on breeding success.

Change factor	Primary driver	Secondary driver
Demographic	Increased survival	Increased breeding success
Ecological	Other	Climate change

Further information on causes of change

The main hypothesis relating to the factors causing the increase in this species concerns the use of lead as fishing weights (Rowell & Spray 2004, Ward et al. 2007). In the late 1970s lead poisoning was shown to be the largest single cause of death among Mute Swans in England, accounting for the deaths of 3,000-Å3,500 birds annually (Kirby et al. 1994). There is good evidence showing that lead contamination of Mute Swans in England caused local population declines during the late 1970s and 1980s (Blus 1994, Birkhead & Perrins 1985). The increase in the British Mute Swan population seen between the 1983 and 1990 censuses can thus be explained partly by the ban on the use of lead weights in fishing imposed by the Water Authorities in 1987 (Rowell & Spray 2004). There is no evidence to suggest that lead poisoning was ever a problem in Scotland (e.g. Brown & Brown 1984).

A second, not mutually exclusive, hypothesis is that warmer winter weather has benefited this species. Deaths during the winter due to poor weather are an important cause of mortality in many areas (Spray 1981, Perrins & Sears 1991) and a run of mild winters is likely to have reduced this (Rowell & Spray 2004). Mild winters are not only associated with low mortality but are also followed by high reproductive output (Delany et al. 1992) which has also contributed to the increase in the Mute Swan population. A study examining five years' data on breeding biology found that winter temperature was one of the factors significantly affecting the date of laying, which in turn was related to clutch size, which in itself was the most significant factor determining the number of cygnets fledged (Birkhead et al. 1983), hence demonstrating an effect on breeding performance. Esselink & Beekman (1991) have also shown that mild winters are not only associated with low mortality but are also followed by high reproductive output be enabling adults attain peak body condition. This may have been particularly important in Scotland.

Whilst the recovery of the British Mute Swan population may in large part be attributed to the reduced incidence of lead poisoning, locally other factors may have had an equal or more important contribution to the observed changes (Ward et al. 2007). Recent years have also seen an increase in the availability of suitable breeding habitats, in the form of the large numbers of gravel pits and ponds that have been created. Improvements to the water quality of rivers and canals, as a result of efforts to reduce pollution, may have also helped the species (Coleman et al. 2001, Rowell & Spray 2004). The number and activity of Swan Rescue Centres may also have an effect on the Mute Swan population size (Delany et al. 1992, Perrins & Martin 1999), although there is little documented evidence to support this. Other factors affecting local populations include increased protection of nesting birds; in an English Midlands study area, this was considered a key factor in the reversal of the 1960s and 1970s

decline (Coleman et al. 2001).

In Scotland (and presumably elsewhere), the increased autumn sowing of cereals has improved the winter food supply for swans, enabling a higher proportion of birds to survive the winter (Delany et al. 1992, Ward et al. 2007), although there are no specific analyses to support this.

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts


Conservation listings:	Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe) (BiE04) UK: not listed (re-established population); amber (localised NW Scottish population); amber (in winter, localised and >20% of NW European Flyway population) (BoCC3)
Long-term trend:	UK waterways: rapid increase
Population size:	46,000 pairs in 2004-08 (APEP13)

Status summary

Apart from an indigenous population in northwest Scotland and the Western Isles, and winter visitors mainly from Iceland, the Greylag Goose is a re-established species throughout the UK. Re-established Greylags increased very rapidly, at a rate estimated at 12% per annum in southern Britain between the 1988-91 Atlas period and 1999 (Rehfishch et al. 2002). This equates across Britain to 170%, or 9.4% per annum, in the period to 2000 (Austiret al. 2007). In Scotland, the native population has grown at an annual rate of 11.7% since 1997 and the re-established birds at 9.7% per annum since 1989 (Mitchell et al. 2011). It has become impossible to distinguish native from re-established populations and they are best now treated as a single unit (Mitchell et al. 2012). The WBS sample became large enough for annual monitoring in 1992, since when further steep increase has been recorded along linear waterways with no sign yet of levelling off. Annual breeding-season monitoring in a wider range of habitats through BBS has shown similar strong increases. Winter counts of resident birds have increased rapidly since the late 1960s (Austin et al. 2014).

WBS/WBBS 1992-2012

Greylag Goose






Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
WBS/WBBS waterways	18	1993-2011	40	486	158	1301		
	10	2001-2011	58	78	16	196		
	5	2006-2011	61	35	3	68		

BBS UK Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
	10	2001-2011	231	111	75	145		
	5	2006-2011	282	26	9	51		
BBS England	16	1995-2011	147	248	137	515		
	10	2001-2011	194	111	67	171		
	5	2006-2011	238	50	31	69		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.



Demographic trends

Productivity and survival trends for this species are not currently produced by BTO

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglinton, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts


Conservation listings:	Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe) (BiE04) UK: not listed (introduced)
Long-term trend:	UK waterways: rapid increase
Population size:	62,000 pairs in 2004-08 (APEP13)

Status summary

Canada Geese were first introduced to English parkland around 1665 but have expanded hugely in range and numbers following translocations in the 1950s and 1960s. They increased rapidly, at a rate estimated at 9.3% per annum in Britain between the 1988-91 Atlas period and 2000, with no sign of any slowing in the rate of increase (Austin et al. 2007). Most of this increase, amounting to 166% during that decade alone, has been in areas previously with low goose densities. The WBS sample became large enough for annual monitoring in 1980, since when further, apparently exponential increase has occurred on linear waterways. Annual breeding-season monitoring in a wider range of habitats through BBS has shown similar strong increases in England and in the UK as a whole but with significant reversals over the last five years. Winter monitoring by WeBS shows a strong long-term increase, but with little change since about 2001 (Austin et al. 2014). The economic, social and environmental impacts of rapidly expanding, non-native Canada Goose populations are of growing conservation concern across Europe.

WBS/WBBS 1980-2012

Canada Goose




Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
WBS/WBBS waterways	30	1981-2011	65	262	59	1136		
	25	1986-2011	75	375	121	1002		
	10	2001-2011	120	102	-1	321		

Source BBS UK	5 Period (yrs)	2006-2011 Years 1995-2011	118 Plots 485	54 Change (%)	5 Lower limit	118 Upper limit	Alert	Comment
	10	2001-2011	557	11	-4	31		
	5	2006-2011	623	-21	-34	-1		
BBS England	16	1995-2011	433	42	22	75		
	10	2001-2011	515	3	-10	22		
	5	2006-2011	575	-25	-36	-4		


Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.



WBS/WBBS waterways graph



BBS UK graph



BBS England graph

Demographic trends

Productivity and survival trends for this species are not currently produced by BTO


This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglinton, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts

Table with 2 columns: Fact Category, Fact Detail. Rows include Conservation listings (Europe: no SPEC category, UK: amber), Long-term trend (UK: probable increase), and Population size (15,000 pairs in 2009).

Status summary

Shelducks occurred on relatively few CBC plots, most of which were close to a coast or an estuary, and it is unclear how well the CBC trend represented the UK breeding population. The CBC showed a substantial increase from the mid 1960s until the early 1980s, some decrease during the 1980s, and stability during the 1990s, although the wide confidence intervals provide scope for other interpretations.




Population changes in detail

Table with 9 columns: Source, Period (yrs), Years, Plots (n), Change (%), Lower limit, Upper limit, Alert, Comment. Rows detail population changes for CBC all habitats and BBS UK/England across different time periods.


Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See here for more information.



Smoothed population index, relative to an arbitrary 100 in 2009, with 85% confidence limits in green



Smoothed population index, relative to an arbitrary 100 in 2009, with 85% confidence limits in green



Smoothed population index, relative to an arbitrary 100 in 2009, with 85% confidence limits in green

Demographic trends

Productivity and survival trends for this species are not currently produced by BTO

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts

Conservation listings:	Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe) (BiE04) UK: amber (winter decline) (BoCC3)
Long-term trend:	UK, England: rapid increase
Population size:	61,000-146,000 pairs in 2009 (APEP13: 1988-91 estimate (APEP06) updated using CBC/BBS trend)


Migrant status:	Resident
Nesting habitat:	Ground nester
Primary breeding habitat:	Wetland
Secondary breeding habitat:	
Breeding diet:	Vegetation
Winter diet:	Vegetation

Status summary

The Mallard has increased steadily as a breeding bird in the UK since the 1960s, especially in England. The BBS Austin et al. 2014). The species has recently been moved from the green to the amber list on the strength of this decline in the UK wintering population. There has been widespread moderate increase across Europe since 1980 but the trend since 1990 has been stable (PECBMS 2013a).

CBC/BBS UK 1966–2012



Mallard








Smoothed population index, relative to an arbitrary 100 in the year given, with 85% confidence limits in green

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	567	167	106	249		
	25	1986-2011	914	31	13	53		
	10	2001-2011	1472	-3	-8	2		
	5	2006-2011	1645	0	-5	5		
CBC/BBS England	44	1967-2011	479	203	126	283		
	25	1986-2011	770	40	23	62		
	10	2001-2011	1244	2	-4	7		
	5	2006-2011	1398	-1	-6	3		
WBS/WBBS waterways	36	1975-2011	164	214	137	288		
	25	1986-2011	200	89	50	131		
	10	2001-2011	295	3	-4	10		
	5	2006-2011	274	5	-1	10		
BBS UK	16	1995-2011	1267	20	11	33		
	10	2001-2011	1444	-3	-9	3		
	5	2006-2011	1598	2	-3	8		
BBS England	16	1995-2011	1068	29	20	40		
	10	2001-2011	1223	2	-4	8		
	5	2006-2011	1363	-1	-5	3		
BBS Scotland	16	1995-2011	98	-6	-20	16		
	10	2001-2011	104	-25	-37	-7		
	5	2006-2011	112	12	-8	38		
BBS Wales	16	1995-2011	65	-17	-51	34		
	10	2001-2011	74	19	-16	58		
	5	2006-2011	74	-6	-27	15		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.





Demographic trends

Productivity and survival trends for this species are not currently produced by BTO

Causes of change

There is little good evidence available regarding the drivers of the breeding population increase in this species in the UK.

Change factor	Primary driver	Secondary driver
Demographic	Unknown	
Ecological	Unknown	

Further information on causes of change

There are no demographic trends available for this species and there is very little evidence generally relating to the causes of the population increases in the UK.

Mallards originating from domesticated birds and not resembling wild-type birds in either plumage or behaviour are very abundant but perhaps under-represented in survey data, especially since many individuals appear to be semi-captive. A large part of the increase in breeding numbers may be attributable to such birds, rather than to true-bred stock. It is also likely that increases may be at least partly attributable to ongoing large-scale releases for shooting (Marchant et al. 1990).

Declines in wintering numbers have been linked to a decrease in continental immigration (Mitchell et al. 2002, Sauter et al. 2010). Guillemain et al. (2010) found trends of increasing average body mass of Mallard in France which were large enough to have major fitness consequences with respect to winter survival, suggesting that overwinter survival has not decreased. Overwinter loss was investigated in Mallard at 35 inland waters in the Midlands and southern England (Hill 1984). Duckling mortality was the key factor, explaining 58% of total mortality between years and this was weakly density dependent. Overwinter loss was higher following years when a large number of young were produced and was the main regulatory factor.

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>


Key facts

Conservation listings:	Europe: SPEC category 3 (declining) (BiE04) UK: amber (European decline) (BoCC3)
Long-term trend:	UK waterways: moderate increase
Population size:	16,000-19,000 pairs in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend for England)

Status summary

The colonisation of the UK by Tufted Ducks, which began in 1849, was aided by the spread of the zebra mussel Dreissena polymorpha, a non-native invasive species that had been introduced accidentally to Britain a few decades earlier. The long-term increase shown by WBS/WBBS, and the increase in range in Britain between the three atlas periods (Gibbons et al. 1993, Balmer et al. 2013) indicate that population expansion and in-filling of range are still occurring. BBS data also show significant increase since 1994 in the UK as a whole. The species' winter trend in the UK since the 1960s, which includes many continental visitors, is also shallowly upward, but with little recent change (Austin et al. 2014). In contrast, moderate recent declines elsewhere in northern Europe have resulted in its reclassification as a species of conservation concern (BirdLife International 2004) and have moved the species from the green to the amber list in the UK (Eaton et al. 2009).

WBS/WBBS 1974-2012
Tufted Duck




Smoothed population index, relative to an arbitrary 100 in the year given, with 85% confidence limits in green

Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
WBS/WBBS waterways	36	1975-2011	36	71	-29	275		
	25	1986-2011	43	14	-36	112		
	10	2001-2011	58	-15	-40	19		
	5	2006-2011	53	-15	-32	4		

BBS UK Source	Period (yrs)	1995-2011 Years	P50s (n)	Change (%)	Bower limit	Upper limit	Alert	Comment
	10	2001-2011	163	12	-11	36		
	5	2006-2011	180	8	-10	31		
BBS England	16	1995-2011	131	33	-4	66		
	10	2001-2011	143	17	-7	38		
	5	2006-2011	160	8	-6	22		


Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.



WBS/WBBS waterways graph



BBS UK graph



BBS England graph

Demographic trends

Productivity and survival trends for this species are not currently produced by BTO

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>


Key facts

Conservation listings:	Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe) (BiE04) UK: green (BoCC3)
Long-term trend:	UK waterways: moderate increase
Population size:	3,500 (3,100-3,800) pairs in 2009 (APEP13: 1987 estimate (Gregory et al. 1997) updated using WBS/WBBS trend)

Status summary

Goosanders were first discovered to have colonised the UK in Perthshire in 1871, and spread from Scotland into northern England in the 1940s (Holloway 1996). Between the first two breeding atlases, the species expanded its range in northern England, and colonised Wales and southwest England. WBS samples became large enough for annual monitoring in 1980, and showed sustained population increase, although this may now have levelled off. The BTO's two national surveys of sawbills demonstrated an average increase in population size of 3% per annum between 1987 and 1997 (Rehfisch et al. 1999). There has been considerable further range expansion since 1990 (Balmer et al. 2013). Reasons for the colonisation of the UK, and the subsequent range expansion and population increase, are unknown. The species' winter trend in Britain, comprising British breeders and continental visitors, rose steeply from the late 1960s to the mid 1990s, but has since fallen back to 1980s levels (Austin et al. 2014).

WBS/WBBS 1980–2012
Goosander




Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
WBS/WBBS waterways	30	1981-2011	42	81	12	189		
	25	1986-2011	47	36	-6	107		
	10	2001-2011	71	4	-14	25		

Source	5 Period (yrs)	2006-2011 Years	66 Plots (n)	6 Change (%)	-10 Lower limit	28 Upper limit	Alert	Comment
--------	----------------------	--------------------	--------------------	--------------------	-----------------------	----------------------	-------	---------

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.



Demographic trends

Productivity and survival trends for this species are not currently produced by BTO

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts


Conservation listings:	Europe: SPEC category 2 (declining) (BiE04) UK: not listed (introduced)
Long-term trend:	UK, England: possible shallow decline
Population size:	82,000 territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)

Status summary

Since Red-legged Partridge is a non-native species released in the UK for the purpose of being shot by hunters, its possible population decrease over the recent 25-year period raises no conservation concern. Moreover, BBS data indicate that significant increase has occurred in the UK and England since 1994. PACEC 2006). The effects on native fauna of releases of such vast scale of this species and Watson et al. 2007). Numbers have shown widespread moderate decline across Europe since 1990, but no longer-term trend is available (PECBMS 2013a).

CBC/BBS UK 1966–2012

Red-legged Partridge







Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	232	-19	-47	19		
	25	1986-2011	383	-18	-36	1		
	10	2001-2011	644	13	5	21		
	5	2006-2011	727	-4	-9	1		
CBC/BBS England	44	1967-2011	226	-23	-54	19		

Source	25 Period (yrs)	1986-2011 Years 2001-2011	373 Plots 621	-21 Change (%)	-39 Lower Limit	2 Upper Limit	Alert	Comment
	5	2006-2011	698	-4	-9	0		
BBS UK	16	1995-2011	545	24	14	37		
	10	2001-2011	634	13	5	21		
	5	2006-2011	711	-4	-10	1		
BBS England	16	1995-2011	530	20	9	31		
	10	2001-2011	613	12	5	20		
	5	2006-2011	685	-4	-10	1		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.



Demographic trends

Productivity and survival trends for this species are not currently produced by BTO

Key facts


Conservation listings:	Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe) (BiE04) UK: amber (25-50% population decline) (BoCC3) UK Biodiversity Action Plan: priority species
Long-term trend:	UK: decline
Population size:	230,000 pairs in 2009 (APEP13: 1988-91 Atlas estimate updated using National Gamebag Census to 1995 and then by BBS trend)

Status summary

The distinctive dark-winged race scotica is endemic to Britain and Ireland and has the vast bulk of its population within the UK, thus conferring global significance to the UK trend. It is economically very important to some rural communities as a game bird and has benefited from intensive management of many moorlands that was designed specifically to increase the numbers of grouse available to be shot. BBS shows fluctuations but no overall trend since 1994. Hudson 1992, Newton 2004), which prompted the move of the species from the green to the amber list in 2002. Longer-term trends in Red Grouse abundance are overlain by cycles, with periods that vary regionally, linked to the dynamics of infection by a nematode parasite Trichostrongylus tenuis (Dobson & Hudson 1992, Gibbonset al. 1993) and to interrelated variations in the aggressiveness of males in autumn (Martinez-Padilla et al. 2014). Montane Fennoscandian populations also declined during 2002-12 (Lehikoinenet al. 2014).




Raptor predation is believed not to affect breeding populations significantly, although it can reduce numbers in the post-breeding period (Redpath & Thirgood 1997). Thompson et al. 2009). Finding a solution to the harrier-grouse conflict would bring considerable benefits to the management of the UK's heather moorlands and have broad implications for the conservation of predators (Redpath & Thirgood 2009).

Laying dates in the Scottish Highlands advanced by about ten days between 1992 and 2011, and were inversely correlated with pre-laying temperatures, but no overall effect of climate change on chick survival could be identified (Fletcher et al. 2013).



Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	16	1995-2011	135	6	-10	22		
	10	2001-2011	151	-8	-22	7		
	5	2006-2011	184	22	6	39		
BBS England	16	1995-2011	77	-4	-23	20		
	10	2001-2011	99	-9	-25	7		
	5	2006-2011	130	19	3	33		
BBS Scotland	16	1995-2011	52	-3	-22	20		
	10	2001-2011	46	-18	-35	-1		
	5	2006-2011	46	15	-3	35		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.



Demographic trends

Productivity and survival trends for this species are not currently produced by BTO


Key facts

Conservation listings:	Europe: SPEC category 3 (vulnerable) (BiE04) UK: red (>50% population decline) (BoCC3) UK Biodiversity Action Plan: click here , priority species
Long-term trend:	UK, England: rapid decline
Population size:	43,000 territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)
Migrant status:	Resident
Nesting habitat:	Ground nester
Primary breeding habitat:	Farmland
Secondary breeding habitat:	
Breeding diet:	Animal
Winter diet:	Vegetation

Status summary

This native gamebird has declined enormously and, despite years of research and the application of a government biodiversity action plan, the continuing decline shown by CBC/BBS suggests that all efforts to boost the population in the wider countryside have so far been unsuccessful. Grey Partridge is one of the most strongly decreasing bird species in Europe, with rapid declines evident in all regions (Kuijper et al. 2009, PECBMS 2009, 2013a). Numbers can be increased within shooting estates where nesting habitat can be provided and pesticide use restricted, but at the expense of corvids, mustelids and foxes (Sotherton et al. 2014).

CBC/BBS UK 1966–2012
Partridge







Smoothed population index, relative to an arbitrary 100 in the year given, with 85% confidence limits in green

Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	132	-90	-93	-86	>50	
	25	1986-2011	176	-76	-82	-71	>50	
	10	2001-2011	225	-31	-42	-23	>25	
	5	2006-2011	237	-24	-33	-14		
CBC/BBS England	44	1967-2011	118	-90	-93	-86	>50	
	25	1986-2011	158	-75	-81	-68	>50	
	10	2001-2011	204	-24	-34	-12		
	5	2006-2011	215	-22	-31	-12		
BBS UK	16	1995-2011	225	-53	-60	-45	>50	
	10	2001-2011	222	-32	-42	-23	>25	
	5	2006-2011	231	-24	-33	-14		
BBS England	16	1995-2011	201	-49	-55	-41	>25	
	10	2001-2011	201	-24	-34	-14		
	5	2006-2011	209	-22	-32	-12		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.





Demographic trends

Productivity and survival trends for this species are not currently produced by BTO

Causes of change

The ultimate factor behind the decline is the deterioration of the bird's agricultural habitat. There is convincing evidence showing that a steep drop in chick survival rate as a result of decreasing chick food availability due to agricultural intensification is the primary driver of population declines. A reduction of hen survival rate during incubation, lower nest success and reduction of winter survival, related to increased predation rates, have all been reported as also playing secondary roles.

Change factor	Primary driver	Secondary driver
Demographic	Decreased breeding success	Reduced adult survival
Ecological	Agricultural intensification	Increased predation

Further information on causes of change

The ultimate factor behind the decline of this species is the deterioration of the bird's agricultural habitat (Aebischer & Ewald 2004). A detailed field and modelling study in the 1980s provides excellent evidence relating to the ecology and population dynamics of the Grey Partridge in a large (62 sq km) study area in Sussex (Potts 1980, Potts 2012). Potts (1980, 2012) identified a reduction in chick survival during the first six weeks after hatching due to a herbicide-induced fall in cereal invertebrate abundance as the primary reason for the decline. More recently, the intensive use of broad-spectrum insecticides on cereals in the summer has been associated with a further reduction in average chick survival rate (Aebischer & Potts 1998). A field study involving an experimental set-up using sprayed and non-sprayed fields confirmed that invertebrate food supplies were important as it was shown that use of pesticides reduced food available to chicks, resulting in lower chick survival and thus depleting numbers of birds being recruited into the population (Rands 1985). Further support for this comes from Sotherton et al. (1993), who also both found that chick survival rate was lower in sprayed than in unsprayed areas.

Potts also identified two other causes for the decline: the disappearance of nesting cover as field boundaries were removed to improve farming efficiency and lower brood production resulting from increased predation. There is evidence from various sources indicating that a reduction of hen survival rate during incubation, lower nest success and a reduction of winter survival, related to increased predation rates, have been influential in the continued population decrease from the 1970s (Potts & Aebischer 1995, Tapper et al. 1996, Bro et al. 2000, De Leo et al. 2004, Panek 2005).

Aebischer & Ewald (2010) offer convincing evidence that, since 2002, local Grey Partridge recoveries have been made possible by sympathetic management of rotational set-aside to provide cover for chicks. In an area of nearly 1,000 ha in Hertfordshire, set-aside was used for habitat creation and Grey Partridge breeding density increased sixfold. However, the disappearance of rotational set-aside in 2007, which halved the amount of brood-rearing habitat, with concurrent poor weather, reversed the increase and effectively removed this potential mechanism for national population recovery.

Overshooting due to the failure of hunters to separate Grey Partridges from [Red-legs](#) can have local population effects, but this is not likely to be a national problem (Aebischer & Ewald 2004). Aebischer & Ewald (2010) showed that on Partridge Count Scheme (PCS) sites, the annual change in spring density in recent years was not related to either shooting pressure or intensity of Red-legged Partridge releasing and suggest that provision of brood-rearing habitats and game cover increased with the latter, which probably counteracted the shooting losses of Grey Partridges on Red-legged Partridge shoots.

In some areas, parasite-mediated apparent competition with the *Tompkinset* al. 2000a, b). However, the evidence for this is conflicting, as Sage et al. (2002) found no deleterious fitness effects of the parasite and Browne et al. (2006) found that poor wild brood survival was indicative of low habitat and food quality rather than of a high rate of parasite infection.


Key facts

Conservation listings:	Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe) (BiE04) UK: not listed (introduced)
Long-term trend:	England: moderate increase
Population size:	2.3 million females in 2009 (APEP13: 1988-91 Atlas estimate (Robertson et al. 1989) updated using CBC/BBS trend for England); at least 35 million captive-reared birds released each autumn (PACEC 2006)
Migrant status:	Resident
Nesting habitat:	Ground nester
Primary breeding habitat:	Farmland
Secondary breeding habitat:	Woodland
Breeding diet:	Vegetation
Winter diet:	Vegetation

Status summary

Pheasants have increased steeply in abundance since the 1960s. The BBS records increase in England and Wales, but little change in Scotland since 1994. During 1968-88, a period when the total biomass of birds in Britain fell by an estimated 10%, CBC data indicate that Pheasant biomass rose by about 2,500 tonnes - more than ten times more than any other species (Dolton & Brooke 1999). The increase has been fuelled by a concurrent steep rise in the numbers of Pheasants released onto shooting estates ([game-bag data](#)). BBS has recorded stability or slight decrease in England and Wales since 2006.

CBC/BBS England 1966-2012
Pheasant



Smoothed population index, relative to an arbitrary 100 in the year given, with 85% confidence limits in green


Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	44	1967-2011	655	90	53	154		
	25	1986-2011	1074	51	33	72		
	10	2001-2011	1768	14	10	18		
	5	2006-2011	2022	-3	-6	0		
BBS UK	16	1995-2011	1767	32	25	40		
	10	2001-2011	2047	17	13	21		
	5	2006-2011	2308	-1	-4	2		
BBS England	16	1995-2011	1493	33	26	40		
	10	2001-2011	1728	15	10	19		
	5	2006-2011	1954	-2	-6	0		
BBS Scotland	16	1995-2011	131	19	0	41		
	10	2001-2011	147	23	10	39		
	5	2006-2011	168	9	-1	19		
BBS Wales	16	1995-2011	91	35	6	76		
	10	2001-2011	108	32	10	55		
	5	2006-2011	115	-9	-17	1		
BBS N.Ireland	16	1995-2011	41	144	41	231		
	10	2001-2011	52	43	17	75		
	5	2006-2011	59	10	-8	31		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




CBC/BBS England 1966–2012
Pheasant




CBC/BBS England graph

BBS UK 1994–2012
Pheasant




BBS UK graph


BBS England 1994–2012
Pheasant




BBS England graph



BBS Scotland graph



BBS Wales graph



BBS N.Ireland graph

Demographic trends

Productivity and survival trends for this species are not currently produced by BTO

Causes of change

The population size of this species is principally determined by releases of reared birds for shooting, which have increased sixfold since 1960. Little is known about the impacts of changes in demographic parameters among wild-breeding birds.

Change factor	Primary driver	Secondary driver
Demographic	Unknown	
Ecological	Other	

Further information on causes of change

It must be noted that numbers of this introduced gamebird are determined principally by releases of reared birds for shooting (Marchant et al. 1990). Such releases have increased approximately sixfold since 1960 ([game-bag data](#)) and were recently running at around 35 million birds annually (PACEC 2006). Robertson (1991) studied records of Pheasant nests from the Nest Record Scheme and found that productivity is probably too low to sustain a population. There is little else known about changes in demographic parameters of Pheasants in the UK.

High Pheasant densities potentially have negative effects, which have not been adequately studied, on native UK birds: these include their effect on the structure of the field layer in woodland, the spread of disease and parasites and competition for food (Fuller et al. 2005). Infection with caecal nematodes from farm-reared Pheasants may be contributing to the decline of Tompkins et al. 2000b), although Sage et al. (2002) found that this had no population impact.

Red-throated Diver

Gavia stellata

Key facts

Conservation listings:	Europe: SPEC category 3 (depleted) (BiE04) UK: amber (European status) (BoCC3); an RBBP species
Long-term trend:	UK: increase
Population size:	1,300 (1,000-1,600) pairs in 2006 (APEP13: Dillon et al. 2009)

Status summary

Population trends are not monitored by the BTO, but JNCC's Mavor et al. 2008). Complete surveys of Shetland indicated a decrease of 36% there between 1983 and 1994, however (Gibbons et al. 1997). The estimated breeding population in 2006 had increased significantly by 34% since the first national survey in 1994, with stability in Shetland and Orkney but increase across the Hebrides and Scottish mainland (Dillon et al. 2009). Since the 1980s, there may have been some tendency for more pairs to hatch a second chick, although two-chick broods are only occasional in Orkney and changes in the distribution of nests recorded might have influenced the results. In 2011, however, there were fewer two-chick broods in Shetland than in any year since at least 1979 (Holling et al. 2013). Nest losses at the egg stage have possibly increased (although samples are small).


Population changes in detail

Annual breeding population changes for this species are not currently monitored by BTO


Demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Clutch size	31	1980-2011	18	None					Small sample
Brood size	31	1980-2011	28	Linear increase	1.25 chicks	1.54 chicks	23.3%		Small sample
Nest failure rate at egg stage	31	1980-2011	10	Linear increase	0.67% nests/day	2.33% nests/day	247.8%		Small sample
Nest failure rate at chick stage	31	1980-2011	16	None					Small sample


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts

Conservation listings:	Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe) (BiE04) UK: green (species level); amber (race <i>carbo</i> , >20% of European breeders; race <i>sinensis</i> , localised breeding) (BoCC3)
Long-term trend:	UK: increase
Population size:	9,000 pairs in 1998-2002 (APEP13: Mitchell et al. 2004)


Status summary

The Cormorant was almost exclusively a coastal breeder in the UK until 1981, but has since established colonies in many inland areas of eastern and central England (Rehfishch et al. 1999; Newson et al. 2006). Breeding had been recorded at 89 inland sites by 2012, and the inland population had risen to about 2,130 pairs by 2005 and 2,362 pairs in 2012 (Newson et al. 2007, 2013). Inland breeding in England is thought to have been sparked by birds of the continental *racesinensis* from the Netherlands and Denmark, although many nominate *carbo* from coastal colonies in Wales and England have contributed to its development.

Breeding numbers and productivity at sample colonies have been monitored annually since 1986 by JNCC's JNCC 2012). Trends during 1986-2005 show decreases in Scotland and in northeast and southwest England, but no trend in Wales, and steep increases inland in England and in regions bordering the northern part of the Irish Sea (Mavor et al. 2008). Reasons for recent decline probably include increased mortality from licensed and unlicensed shooting. BBS counts are very largely of immature or other non-breeding birds inland and away from breeding sites and, until we have better information on the proportions of breeding and non-breeding birds recorded on BBS, the generally upward long-term trend probably adds little information about breeding numbers. The winter trend in Britain, comprising British and Irish breeders and continental visitors, has shown strong increase since the late 1980s but is now stable or in shallow decline, with an increase in shooting under licence since 2004 possibly taking an effect on population growth (Chamberlain et al. 2013, Austin et al. 2014). Although the species is now green listed, both races that occur in the UK qualify for amber listing, for reasons unconnected with the UK trend.



BBS UK 1994–2012

Cormorant



Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	16	1995-2011	237	21	-8	57		Non-breeders included
	10	2001-2011	288	-9	-23	10		Non-breeders included
	5	2006-2011	318	-4	-18	12		Non-breeders included
BBS England	16	1995-2011	199	16	-5	48		Non-breeders included
	10	2001-2011	243	2	-14	18		Non-breeders included
	5	2006-2011	272	-4	-17	12		Non-breeders included

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.



Demographic trends

Productivity and survival trends for this species are not currently produced by BTO

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Grey Heron


Ardea cinerea

Key facts

Conservation listings:	Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe) (BiE04) UK: green (BoCC3)
Long-term trend:	UK, Wales, Scotland: shallow increase England: moderate increase
Population size:	13,000 pairs in 2007-11 (APEP13); 11,322 (10,792-12,006) apparently occupied nests in 2012 (Heronries Census)

Status summary

The BTO Heronries Census, which has monitored Grey Herons since 1928, shows the species to have been more abundant in the early 2000s than at any time in the last 80 years. The effects of harsh winters, which induce severe mortality in this species (Besbeas et al. 2002), are clearly visible in the long-term trend. The general increase that underlies these fluctuations may stem from reduced persecution, improvements in water quality, the provision of new habitat as new lakes and gravel pits mature, and increased feeding opportunities at freshwater fisheries (Gibbons et al. 1993, Marchant et al. 2004). A downturn evident since 2001 is, as yet, unexplained, though recent cold winter weather appears to have accelerated the decline. High rates of nest failure at the chick stage were noted in the late 1960s, but not subsequently. Clutch and brood sizes have fallen in the long term. In the latest special survey of UK heronries, carried out in 2003 to mark the 75th anniversary of the Heronries Census, a record total of more than 10,441 Grey Heron nests were counted, around 75% of the estimated total population. Numbers have shown widespread moderate increase across Europe since 1980 (PECBMS 2013a).






Population changes in detail






Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Heronries UK	82	1929-2011	334	35	4	61		
	25	1986-2011	535	7	0	15		
	10	2001-2011	619	-16	-21	-12		
	5	2006-2011	635	-14	-17	-10		
Heronries England and Wales	82	1929-2011	276	39	7	62		
	25	1986-2011	437	6	-1	13		
	10	2001-2011	497	-17	-21	-13		
	5	2006-2011	508	-15	-17	-11		
Heronries England	82	1929-2011	233	38	7	62		
	25	1986-2011	366	8	1	13		
	10	2001-2011	426	-16	-20	-12		
	5	2006-2011	437	-14	-17	-11		
Heronries Scotland	76	1935-2011	47	11				
	25	1986-2011	75	31				
	10	2001-2011	91	1				

Source	Period (yrs)	2006-2011 Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Heronries Wales	76	1935-2011	43	-17				
	25	1986-2011	69	-8				
	10	2001-2011	70	-19				
	5	2006-2011	71	-20				
BBS UK	16	1995-2011	657	-8	-17	2		Non-breeders included
	10	2001-2011	755	-24	-30	-18		Non-breeders included
	5	2006-2011	811	-20	-26	-14		Non-breeders included
BBS England	16	1995-2011	542	-15	-23	-4		Non-breeders included
	10	2001-2011	630	-20	-27	-14		Non-breeders included
	5	2006-2011	688	-22	-26	-17		Non-breeders included
BBS Scotland	16	1995-2011	49	5	-19	38		Non-breeders included
	10	2001-2011	52	-31	-47	-11	>25	Non-breeders included
	5	2006-2011	55	-22	-41	-2		Non-breeders included
BBS Wales	16	1995-2011	43	-20	-36	9		Non-breeders included
	10	2001-2011	47	-35	-45	-21	>25	Non-breeders included
	5	2006-2011	43	-27	-40	-10	>25	Non-breeders included

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.



Grey Heron Scotland graph




Demographic trends


Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	17	None					
Fledglings per breeding attempt	43	1968-2011	17	None					

Clutch size	43	1968-2011	15	Linear decline	4.05 eggs	3.44 eggs	-15.1%		Small sample
Variable	Period	Years	Mean annual	Trend	Modelled in	Modelled in	Change	Alert	Comment
Brood size	43	1968-2011	99	Linear decline	2.84 chicks	2.43 chicks	-14.7%		
Nest failure rate at egg stage	43	1968-2011	17	Curvilinear	0.00% nests/day	0.02% nests/day			Small sample
Nest failure rate at chick stage	43	1968-2011	40	None					


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

Key facts


Conservation listings:	Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe) (BiE04) UK: amber (25-50% population decline) (BoCC3)
Long-term trend:	UK: uncertain
Population size:	3,900-7,800 pairs in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)

Status summary

The moderate decline shown by the WBS/WBBS may reveal problems among birds on linear waterways during the early 1980s and since the late 1990s, but stability in the long term BBS trend suggests that wider populations (including birds on small still waters) have been more healthy. Because of the shortage of data, and the conflict between WBS and BBS assessments, the rapid decline indicated by WBS in the 1980s did not initially trigger a conservation listing. The species was moved from the green to the amber list in 2009, however, on the strength of its UK decline. BBS data are now showing significant decline in the recent five-year period. In an analysis of nest record cards, Moss & Moss (1993) found that nests on ponds and lakes were significantly more successful than those on rivers and streams and that nests on rivers, subject to fluctuating water levels, experienced significantly higher failure rates through flooding than those on canals, where water levels are artificially maintained. Winter numbers, as monitored by WeBS, showed sustained shallow increase until 2008, followed by a minor decline (Austin et al. 2014).

WBS/WBBS 1974-2012

Little Grebe




Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
WBS/WBBS waterways	36	1975-2011	21	-49	-73	10		
	25	1986-2011	21	-23	-53	43		
	10	2001-2011	23	-6	-35	55		

Source	Period (yrs)	Years	Plots	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	5	2006-2011	22	-16	-36	18		
		1995-2011	70					
	10	2001-2011	81	-12	-31	8		
	5	2006-2011	87	-17	-31	-1		
BBS England	16	1995-2011	56	-4	-32	41		
	10	2001-2011	64	-16	-36	3		
	5	2006-2011	67	-15	-33	5		


Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.



WBS/WBBS waterways graph



BBS UK graph



BBS England graph

Demographic trends

Productivity and survival trends for this species are not currently produced by BTO

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Great Crested Grebe


Podiceps cristatus

Key facts

Conservation listings:	Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe) (BiE04) UK: green (BoCC3)
Long-term trend:	UK: probable increase
Population size:	5,300 pairs in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)

Status summary

This species was believed to be on the verge of extinction in Britain around 1860, when only 32-72 pairs were known in England (Holloway 1996). A subsequent increase followed reductions in persecution, aided by statutory protection, and the creation of extensive new habitat in the form of gravel pits (Gibbons et al. 1993). Increase was tracked by special surveys to around 7,000 adult birds in Britain by 1975 (Hughes et al. 1979). The BBS provides the first annual, national monitoring of this species and indicates shallow increase since 1994. Winter numbers, monitored by WeBS, have shown a long-term shallow increase but may now be in shallow decline (Austin et al. 2014).





Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	16	1995-2011	70	11	-26	40		
	10	2001-2011	77	32	1	60		
	5	2006-2011	83	-11	-36	16		
BBS England	16	1995-2011	64	-8	-23	12		

Source	10 Period (yrs)	2001-2011 Years 2006-2011	71 Plots (%)	14 Change (%)	-5 Lower limit	32 Upper limit	Alert	Comment
--------	-----------------------	---------------------------------	--------------------	---------------------	----------------------	----------------------	-------	---------

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.



Demographic trends

Productivity and survival trends for this species are not currently produced by BTO

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts


Conservation listings:	Global: Near Threatened (BiE04) UK: amber (European decline) (BoCC3); an RBBP species
Long-term trend:	UK, England: rapid increase
Population size:	1,600 pairs in 2006-10 (APEP13: RBBP data); probably over 2,200 pairs in 2010 (Holling & RBBP 2012)

Status summary

Red Kite was historically widespread across Britain but, following widespread persecution, fewer than ten breeding pairs remained by the 1930s and 1940s, concentrated into a small area of mid Wales. Through careful husbandry organised by a 'Kite Committee' of local conservationists and landowners, including RSPB bounties paid to farmers for successful nests during 1922-90, the Welsh population rose to 100 pairs by 1993. Most birds were descended from a single female that had continued to breed successfully during the population bottleneck (Carter 2001). As a step towards restoring the original breeding range, birds were introduced in 1989 into the Chilterns (Oxfordshire and Buckinghamshire) and into the Black Isle in Easter Ross (Evans & Pienkowski 1991). Successful breeding populations quickly established in both areas. Further releases were begun in Northamptonshire in 1995, central Scotland in 1996, Yorkshire in 1999, Dumfries & Galloway in 2001, northeast England in 2004 and County Down in 2008. Each of these centres has given rise to a productive breeding group, in some cases benefiting from large-scale provision of food or the development of a well-established communal roost. Introduced birds and their offspring wander widely across Britain and Ireland but, as yet, pairs have been slow to set up breeding sites distant from the release areas (Balmer et al. 2013). BBS sightings have shown an exponential rise since 1994. Illegal killing is continuing and in northern Scotland the use of poisoned baits deliberately to kill raptors has severely limited the growth of the Red Kite population (Smart et al. 2010).

BBS UK 1994–2012

Red Kite





Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	16	1995-2011	87	676	369	1265		

Source	10 Period (yrs)	2001-2011 Years 2006-2011	130 Plots (174)	356 Change (193)	242 Lower limit	518 Upper limit	Alert	Comment
BBS England	16	1995-2011	62	11260	4751	10585		
	10	2001-2011	96	1099	811	1570		
	5	2006-2011	135	153	125	197		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.



Demographic trends

Productivity and survival trends for this species are not currently produced by BTO

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Hen Harrier

Circus cyaneus

Key facts

Conservation listings:	Europe: SPEC category 3 (vulnerable) (BiE04) UK: red (historical decline) (BoCC3); an RBBP species
Long-term trend:	UK: probable increase
Population size:	662 (576-770) pairs in UK and Isle of Man in 2010 (Hayhow et al. 2013)

Migrant status:	Resident
Nesting habitat:	Ground nester
Primary breeding habitat:	Moorland
Secondary breeding habitat:	
Breeding diet:	Animal
Winter diet:	Animal

Status summary

This species was red listed because of substantial declines over the last two centuries. The UK population was unchanged between surveys in 1988-89 and 1998, with declines in Orkney and England but increases in Northern Ireland and the Isle of Man (Sim et al. 2001). A 41% increase was recorded in the UK and Isle of Man during 1998-2004, possibly due to increased use of non-moorland habitats, but with decreases in the Southern Uplands, east Highlands and England, all being areas with many managed grouse moors (Sim et al. 2007a). The latest survey, in 2010, revealed a decline of around 18% since the 2004 survey: a notable decrease in Scotland might stem from habitat change and illegal persecution, while illegal persecution continues to limit harriers in England to very low levels of population (Hayhow et al. 2013).


Population changes in detail

Annual breeding population changes for this species are not currently monitored by BTO


Demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Clutch size	43	1968-2011	12	None					Small sample
Brood size	43	1968-2011	19	None					Small sample
Nest failure rate at egg stage	43	1968-2011	10	Curvilinear	0.02% nests/day	0.05% nests/day	150.0%		Small sample
Nest failure rate at chick stage	43	1968-2011	13	None					Small sample


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

Causes of change

Based on multiple field studies providing good evidence, the main driver of declines in Hen Harrier populations appears to have been illegal persecution, causing a reduction in nesting success, annual productivity and survival of breeding females.

Change factor	Primary driver	Secondary driver
Demographic	Decreased breeding success	Decreased survival
Ecological	Other	

Further information on causes of change

Demographic data presented here show that clutch size decreased by 12% between 1968 and 2008 (although further investigation has shown that this trend is due to the increased proportions in recent years of records from Orkney, where clutch sizes tend to be smaller than on the mainland: Summers 1998, Crick 1998). The daily failure rate of nests at the egg stage have increased by 500%. However, this is based on extremely small sample sizes so these figures have to be treated with caution.

There is good evidence showing that, although the Hen Harrier has been protected under UK law since 1961, many are still unlawfully killed or disturbed in efforts to protect the economic viability of driven shooting of Redpath & Thirgood 1997, Thompson et al. 2009). A study combining Atlas data and a two-year field study provided good evidence that nesting success, annual productivity and survival of female Hen Harriers was lower on grouse moors than on other moorland or in young conifer forests, due to destruction by humans (Bibby & Etheridge 1993, Etheridge et al. 1997). Fielding et al. (2011) conclude that illegal killing is the biggest single factor affecting the species and that it is having a dramatic impact on the population in core areas of its range in northern England and Scotland. Keepering that remains within the law, however, can benefit harrier populations by increasing their prey and reducing their nest predators, especially crows and foxes (Baines & Richardson 2013).

Recovery of the Welsh harrier population, in contrast to those elsewhere in the UK, has been attributed to an increase in the breeding productivity, apparently due to a combination of cessation of human interference in recent years and warmer temperatures, leading to increased productivity (Whitfield et al. 2008). Whitfield et al. (2008) also provide strong field-based evidence from the Welsh harrier population that human interference has been the primary driver of population change, through its impact on breeding productivity (specifically, an increased proportion of breeding females laying eggs, combined with a general increase in the average number of young fledged).

In areas where illegal persecution is minimal, food availability restricts numbers. Good-quality recent studies found that rough grass, a preferred habitat for field voles, is a critical foraging habitat for Orkney Hen Harriers (Amar & Redpath 2005, Amar et al. 2008a) and that habitat characteristics around harrier nest-sites (at a 1-km radius) can have a strong influence on breeding performance (Amar et al. 2002).

A field experiment showed that food shortage just before the laying period resulted in low levels of polygyny and reduced nesting success among secondary females, resulting in reduced productivity (Amar & Redpath 2002, Amar et al. 2005). The area of rough grassland has decreased during the same period as sheep numbers have increased and this is thought have reduced food supplies (Amar et al. 2003, 2005, Amar & Redpath 2005), but there was no detectable effect of rough grass area on fledging success or fledged brood size (Amar et al. 2008). Further, these studies provide no evidence that the effects on breeding success have an impact on abundance. However, Redpath et al. (2002a) present good evidence from a different field study in Scotland which also shows that food availability, notably numbers of field voles, can influence population change in Hen Harriers, where there is no persecution. Harrier densities were highest in areas and years where their small prey animals were most abundant. Clutch size was positively correlated with the number of field voles, although fledging success was not significantly correlated with the relative abundance of small prey (Redpath & Thirgood 1999, Redpath et al. 2002a). Madders (2000) also highlighted the importance of foraging habitat in Scotland, finding that the extent of young first-rotation forestry, the preferred foraging habitat in this area, is currently in decline and states that this has contributed to many of the reported changes in local Hen Harrier populations (although no specific research into demographic parameters were presented).

There is some evidence that climate also affects demography, although this is secondary to drivers outlined above and there is no evidence for effects on abundance. In Scotland, chick mortality increased in cold temperatures and annual values of harrier fledged brood size were positively related to summer temperature (Redpath et al. 2002b) and warmer temperatures led to increased productivity (in the absence of persecution) in Wales (Whitfield et al. 2008).

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Sparrowhawk

Accipiter nisus

Key facts


Conservation listings:	Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe) (BiE04) UK: green (BoCC3)
Long-term trend:	England: rapid increase
Population size:	35,000 pairs in 2009 (APEP13: 1988-91 Atlas estimate (Newton 1986) updated using CBC/BBS trend for England)
Migrant status:	Resident
Nesting habitat:	Above-ground nester
Primary breeding habitat:	Woodland
Secondary breeding habitat:	
Breeding diet:	Animal
Winter diet:	Animal

Status summary

Between the 1970s and the mid 1990s, the CBC charted a steep increase in this species. Many former haunts especially in the Midlands and east of England were reoccupied between the first two atlas periods (Gibbons et al. 1993). The population has stabilised since the mid 1990s, though population fluctuations are now evident. Nest productivity has risen, especially during the period of strong population increase. There has been little long-term change across Europe since 1980 (PECBMS 2013a).


CBC/BBS England 1974–2012

Sparrowhawk




Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	36	1975-2011	162	153	49	312		
	25	1986-2011	219	35	3	65		
	10	2001-2011	325	-7	-17	1		
	5	2006-2011	355	-7	-16	0		
BBS UK	16	1995-2011	349	0	-10	12		
	10	2001-2011	392	-8	-16	2		
	5	2006-2011	429	-5	-13	4		
BBS England	16	1995-2011	286	-4	-15	6		
	10	2001-2011	320	-8	-17	1		
	5	2006-2011	346	-7	-14	0		


Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.



CBC/BBS England graph




BBS UK graph



BBS England graph

Demographic trends


Fledglings per breeding attempt Sparrowhawk 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012


Sparrowhawk




More on demographic trends


Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	32	Curvilinear	2.64 fledglings	3.15 fledglings	19.2%		
Clutch size	43	1968-2011	34	None					
Brood size	43	1968-2011	69	Curvilinear	3.16 chicks	3.39 chicks	7.4%		
Nest failure rate at egg stage	43	1968-2011	32	Linear decline	0.47% nests/day	0.06% nests/day	-87.2%		
Nest failure rate at chick stage	43	1968-2011	45	None					
Laying date	43	1968-2011	13	None			0 days		Small sample

For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).






Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

Causes of change

There is good evidence that improved breeding success due to a decline in organochlorine pesticide use is the most likely cause of the increase in this species, but that reduced survival, especially of young birds, may be driving the decline in Scottish populations.

Change factor	Primary driver	Secondary driver
Demographic	Increased breeding success	Increased survival
Ecological	Other	

Further information on causes of change

Sparrowhawks suffered a severe population crash caused by organochlorine pesticides in the 1950s and 1960s, when the species was extinguished from large areas of lowland Britain (Newton 1986, 2013). Studies of this species in eastern England confirmed this, and the recovery of the Sparrowhawk in this area was primarily dependent on declining organochlorine contamination which resulted in an improvement of breeding success mainly due to an increase in hatching success, itself associated with improved eggshell thickness and reduced egg breakage (Newton & Wyllie 1992). The figures above support this, showing improving numbers of fledglings per breeding attempt, a fall in failure rates at the egg stage and increases in brood size.

Comparison of an increasing population in east-central England with stable and decreasing populations in southern Scotland showed that differences in population trend were associated mainly with differences in the recruitment of new breeders (greatest in the increasing and lowest in the decreasing population) and in age of first breeding (earliest in the increasing and latest in the decreasing population). There were also differences in the annual survival of breeders (greater in the increasing population) while differences in breeding success between areas were slight and non-significant (Wyllie & Newton 1991). A comprehensive long-running study of Sparrowhawks in Scotland during 1972-86 provides further detailed evidence. Overwinter loss operating in the period between the fledging of young and subsequent recruitment to the breeding population was identified as the key factor, explaining 77% of the variance in total annual loss, and largely accounting for the pattern of change in breeding numbers (Newton 1988). Work by Newton & Marquiss (1986) found that annual survival of established breeders and breeding performance was the same in both a declining and increasing population, but that recruitment of incoming breeders was lower in the declining population and state that this was the main proximate cause of decline.

The population has stabilised since the mid 1990s and, possibly through the effects of intraspecific competition, average brood size has begun to fall again (see above).

Key facts


Conservation listings:	Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe) (BiE04) UK: green (BoCC3)
Long-term trend:	England: rapid increase
Population size:	57,000-79,000 pairs in 2009 (APEP13: 2001 estimate (Clements 2002) updated using BBS trend)
Migrant status:	Resident
Nesting habitat:	Above-ground nester
Primary breeding habitat:	Woodland
Secondary breeding habitat:	Farmland
Breeding diet:	Animal
Winter diet:	Animal

Status summary

The Buzzard has shown a substantial eastward range expansion since the 1988-91 Atlas and is now an almost ubiquitous breeding bird in the UK (Balmer et al. 2013). For more than a decade it has been the most abundant UK raptor (Clements 2002). The increasing trend identified by the CBC relates especially to the spread of range into central and eastern Britain, where CBC was strongly represented. If anything, however, the upsurge has been amplified with the addition of the more widely representative BBS data since 1994. The BBS PECBMS 2013a). Though breeding success is still rising overall, a decrease in productivity has been documented in Avon, per pair but not per unit area, as the population has risen (Prytherch 2013) .

CBC/BBS England 1966–2012

Buzzard




Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	44	1967-2011	236	712	433	1829		Small CBC sample
	25	1986-2011	408	683	436	1317		
	10	2001-2011	811	83	70	99		
	5	2006-2011	1015	35	28	40		
BBS UK	16	1995-2011	921	80	62	96		
	10	2001-2011	1181	30	22	40		
	5	2006-2011	1419	17	12	22		
BBS England	16	1995-2011	607	167	136	210		
	10	2001-2011	811	84	73	98		
	5	2006-2011	1015	35	29	40		
BBS Scotland	16	1995-2011	144	31	8	60		
	10	2001-2011	170	-14	-25	1		
	5	2006-2011	197	-1	-15	12		
BBS Wales	16	1995-2011	140	4	-16	22		
	10	2001-2011	159	-1	-13	11		
	5	2006-2011	161	-1	-11	10		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




CBC/BBS England 1966-2012
Buzzard




CBC/BBS England graph

BBS UK 1994-2012
Buzzard




BBS UK graph


BBS England 1994-2012
Buzzard



BBS England graph




BBS Scotland graph



BBS Wales graph

Demographic trends


Fledglings per breeding attempt Buzzard 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Buzzard




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	28	Linear increase	1.51 fledglings	2.07 fledglings	36.9%		
Clutch size	43	1968-2011	35	Curvilinear	2.20 eggs	2.16 eggs	-2.0%		
Brood size	43	1968-2011	109	Curvilinear	1.89 chicks	1.97 chicks	4.4%		
Nest failure rate at egg stage	43	1968-2011	29	Linear decline	0.85% nests/day	0.05% nests/day	-94.1%		Small sample
Nest failure rate at chick stage	43	1968-2011	54	None					


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

Causes of change

There is good evidence that the increase in population numbers is associated with rapidly improving nesting success, which has been linked to reduced persecution (and therefore improved survival) and increased food supplies due to the recovery of rabbit populations from the effects of myxomatosis. It is not possible to say which is the more important driver.

Change factor	Primary driver	Secondary driver
Demographic	Improved breeding success	Increased survival
Ecological	Other	

Further information on causes of change

As the figures above show, there has been an increase in the number of fledglings per breeding attempt and a decrease in daily failure rates at the egg stage. As such, the increase in population numbers has been associated with rapidly improving nesting success, through reduced persecution, the recovery of rabbit populations from the effects of myxomatosis and release from the deleterious effects of organochlorine pesticides (Elliott & Avery 1991, Sim et al. 2000, 2001a, Clements 2002). Numbers of Buzzard were relatively stable until the late 1980s when the population size began increasing steeply. Elliott & Avery (1991) analysed data collected by the RSPB to provide good evidence that, during 1975-89, persecution was a factor in restricting the Buzzard's range. Halley (1993) found that levels of persecution in Scotland had fallen and postulated that this was a factor in the increase in Buzzard population size. In a study of two local populations in Scotland, Swann & Etheridge (1995) provided some evidence to show that persecution was a factor in restricting population density at the site that benefited from higher productivity, although they did not specifically analyse the effects of persecution. Sim et al. (2000) provide good evidence from Buzzard populations in the West Midlands that persecution levels, especially poisonings, were lower in the 1990s when the population started increasing and state that higher survival rate due to reduced persecution was likely to be one of the main factors responsible for the rapid increase in the Buzzard population in this area. Gibbons et al. (1995) found that Buzzards were less common in the uplands where grouse moors were most frequent, stating that this was due to either persecution, unsuitable habitat management or lack of food, although did not specify which was the most important driver.

There is also good evidence to support the role of changing food availability in population increases. Graham et al. (1995) showed that Buzzard breeding density was positively related to lagomorph abundance and Swann & Etheridge (1995) found that Buzzards laid larger clutches, produced bigger broods and had significantly higher productivity where rabbits were more common. Sim et al. (2000, 2001a) also provided good evidence that increased productivity coincided with an increase in rabbit abundance. Other studies have also found that breeding success is related to food availability (Kostrzewa & Kostrzewa 1991, Austin & Houston 1997, Goszczynski 1997, 2001). It is, therefore, plausible that Buzzard distribution is influenced by rabbit abundance, which has increased since rabbits have overcome the effects of myxomatosis.

Habitat change may have played some role in the increases. High Buzzard breeding densities were associated with high proportions of unimproved pasture and mature

woodland within estimated territories (Sim et al. 2000) and Sergio et al. (2002, 2005) found that Buzzard productivity benefited from the conversion of coppice woodland to mature forest in Italy. There is also some evidence that breeding success is related to climate, although there is little evidence for this from the UK. In Germany, Kostrzewa & Kostrzewa (1990) provide evidence to show that the number of young fledged was negatively correlated with rainfall in April and May. Although there is no evidence to support this, it is worth noting that these possible habitat/climate effects and food effects are not mutually exclusive.

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts


Conservation listings:	Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe) (BiE04) UK: green (BoCC3)
Long-term trend:	UK: probable shallow decline
Population size:	270,000 territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)

Status summary

Trends for this species show wide fluctuations that are related to its high potential for reproduction and to its susceptibility to cold winter weather. The BBS Leech & Barimore 2008); there has been shallow decline in the number of fledglings per breeding attempt. There has been little long-term change across Europe since 1980 (PECBMS 2013a).

CBC/BBS UK 1966–2012

Moorhen







Smoothed population index, relative to an arbitrary 100 in the year given, with 85% confidence limits in green


Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	321	-2	-25	26		
	25	1986-2011	485	-13	-24	2		
	10	2001-2011	727	-22	-26	-16		
	5	2006-2011	798	-25	-28	-20		
CBC/BBS England	44	1967-2011	295	6	-19	30		
	25	1986-2011	449	-12	-24	2		

Source	Period (yrs)	2001-2011 Years	Birds (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
	5	2006-2011	749	-23	-26	-18		
WBS/WBBS waterways	36	1975-2011	123	-30	-45	-4	>25	
	25	1986-2011	145	-6	-26	23		
	10	2001-2011	198	-20	-30	-11		
	5	2006-2011	182	-22	-29	-15		
BBS UK	16	1995-2011	650	-9	-15	0		
	10	2001-2011	723	-22	-27	-16		
	5	2006-2011	792	-24	-27	-19		
BBS England	16	1995-2011	602	-10	-16	-3		
	10	2001-2011	673	-21	-25	-16		
	5	2006-2011	743	-22	-25	-17		


Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.





Demographic trends


Fledglings per breeding attempt Moorhen 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012


Moorhen




More on demographic trends


Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	52	Linear decline	2.51 fledglings	2.13 fledglings	-15.1%		
Clutch size	43	1968-2011	114	None					
Brood size	43	1968-2011	108	None					
Nest failure rate at egg stage	43	1968-2011	140	Linear increase	1.10% nests/day	2.17% nests/day	97.3%		
Nest failure rate at chick stage	43	1968-2011	52	Linear increase	0.04% nests/day	0.18% nests/day	350.0%		
Laying date	43	1968-2011	83	Linear decline	May 10	May 4	-6 days		

For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).






Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

Key facts


Conservation listings:	Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe) (BiE04) UK: green (BoCC3)
Long-term trend:	UK, England: rapid increase
Population size:	31,000 pairs in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)
Migrant status:	Resident
Nesting habitat:	Ground nester
Primary breeding habitat:	Wetland
Secondary breeding habitat:	
Breeding diet:	Vegetation
Winter diet:	Vegetation

Status summary

WBS/WBBS and CBC/BBS trends for Coot indicate a long-term increase, although the magnitude of the change is not clear. Small CBC samples, mainly of birds on small water-bodies, suggested a rapid rise in the late 1960s. WBS/WBBS and BBS include more birds on larger waters, and so may be more representative of Coot populations, but WBS/WBBS has not recorded the strong increase found by BBS observers since 1994. The five-year trend is downward in all indices. The combination of CBC and BBS data suggests that the long-term increase in the UK and England has been rapid. There has been widespread moderate increase across Europe since 1980, though with little change since 1990 (PECBMS 2013a). Winter abundance on large still waters, as monitored by WeBS, showed shallow increase from the mid 1980s to around 2000/01 but has since declined, especially in Northern Ireland (Austin et al. 2014).

CBC/BBS UK 1966–2012




Coot




Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	123	182	75	550		Small CBC sample
	25	1986-2011	195	34	5	77		Small CBC sample
	10	2001-2011	308	-7	-17	5		
	5	2006-2011	341	-8	-18	4		
CBC/BBS England	44	1967-2011	112	170	72	490		Small CBC sample
	25	1986-2011	176	29	3	86		Small CBC sample
	10	2001-2011	279	-10	-21	0		
	5	2006-2011	309	-11	-21	-1		
WBS/WBBS waterways	36	1975-2011	61	83	13	228		
	25	1986-2011	74	33	-19	101		
	10	2001-2011	98	1	-27	28		
	5	2006-2011	88	-9	-24	7		
BBS UK	16	1995-2011	267	29	11	51		
	10	2001-2011	306	-6	-17	7		
	5	2006-2011	338	-6	-17	5		
BBS England	16	1995-2011	242	25	7	46		
	10	2001-2011	277	-10	-22	2		
	5	2006-2011	307	-12	-22	-1		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.





BBS UK graph



BBS England graph

Demographic trends

Productivity and survival trends for this species are not currently produced by BTO

Causes of change

There are no demographic trends available for this species and very little evidence regarding the ecological drivers of change.

Change factor	Primary driver	Secondary driver
Demographic	Unknown	
Ecological	Unknown	

Further information on causes of change

There is very little information available regarding the demographic or ecological drivers of population change in Coot.

Brinkhof & Cave (1997) conducted a supplementary feeding experiment and found that seasonal variation in offspring production was in essence the result of seasonal variation in food availability. Thus, increases in food supply may have improved breeding success, but there is no evidence to support this.

Work from Finland (Ronka et al. 2005) has suggested that Coot are sensitive to overwinter weather: thus it is possible that this species may have benefited from milder winters.


This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts

Conservation listings:	Europe: no SPEC category (concentrated in Europe, conservation status favourable) (BiE04) UK: amber (>20% of European breeding population, >20% of East Atlantic Flyway population in winter, localised wintering population) (BoCC3)
Long-term trend:	UK waterways: moderate increase
Population size:	110,000 pairs in 2009 (APEP13: 1985-99 estimate (O'Brien 2004) updated using CBC/BBS trend)

Status summary

Oystercatchers increased along linear waterways between 1974 and about 1986, as the species colonised inland sites across England and Wales (Gibbons et al. 1993). Thereafter, the WBS/WBBS index stabilised and now appears to be in decline, so showing a pattern similar to that in winter abundance revealed by WeBS (Austin et al. 2014). Surveys in England and Wales revealed an increase of 47% in breeding birds in wet meadows between 1982 and 2002 (Wilson et al. 2005). BBS data since 1994, which include birds in a broader range of locations and habitats, show strong increase in England but a significant decline in Scotland. The increase in nest failure rates during the 27-day egg stage (25 days for incubation and 2 days for laying) probably results from the spread of the species into less favourable habitats, where nest losses through predation or trampling may be more likely. The trend towards earlier laying may be linked to recent climate change (Crick & Sparks 1999). There have been widespread moderate declines across Europe since 1980 (PECBMS 2013a).




Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
WBS/WBBS waterways	36	1975-2011	47	62	26	158		
	25	1986-2011	59	-19	-39	21		
	10	2001-2011	96	-22	-32	-10		

Source	Period (yrs)	Years	Plots	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	5	2006-2011	92	-17	-27	-8		
	10	2001-2011	375	-9	-17	1		
	5	2006-2011	428	-4	-12	7		
BBS England	16	1995-2011	176	47	20	73		
	10	2001-2011	215	23	6	41		
	5	2006-2011	253	5	-5	17		
BBS Scotland	16	1995-2011	127	-30	-39	-20	>25	
	10	2001-2011	130	-19	-29	-8		
	5	2006-2011	137	-7	-19	6		


Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




WBS/WBBS waterways graph



BBS UK graph




BBS England graph




BBS Scotland graph

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Clutch size	43	1968-2011	146	None					
Nest failure rate at egg stage	43	1968-2011	160	Curvilinear	1.46% nests/day	4.41% nests/day	202.1%		
Laying date	43	1968-2011	65	Curvilinear	May 18	May 13	-5 days		

For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).



Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>


Key facts

Conservation listings:	Europe: no SPEC category (concentrated in Europe, conservation status favourable) (BiE04) UK: amber (BoCC3)
Long-term trend:	UK: probable decline
Population size:	38,400-59,400 pairs in 1980-2000 (APEP13: BiE04)

Status summary

There was no annual monitoring of the breeding population before the inception of BBS. Since 1994, BBS has shown stability or minor decrease in the UK and Scotland, but this is believed to follow an earlier decline (Gibbons et al. 1993). A detailed survey has confirmed that a sharp decline has occurred in Wales since the 1980s, with just 36 pairs located in 2007 (Johnstone et al. 2008). Nest survival on grass moors, unlike that on heather moors, may have declined over time (Crick 1992), perhaps linked to increased stocking densities of sheep (Fuller 1996). Clutch size has decreased slightly, though a large number of late-season nest records, which provide higher proportions of two- and three-egg clutches, were submitted from an intensive study during 1996-98 (J.W. Pearce-Higgins, pers. comm.). Warmer springs are reported to advance the breeding phenology of Golden Plovers and of their tipulid prey (Pearce-Higgins et al. 2005). There has been no long-term change across Europe since 1981, but moderate increase since 1990 (PECBMS 2013a). Winter numbers counted by WeBS, although mainly at coastal sites and omitting some big concentrations inland, increased strongly in Britain between the mid 1980s and 2006, since when there has been a sharp fall (Austin et al. 2014); these birds are mainly of Fennoscandian or Russian origin. The species has recently been restored to the amber list because of the international importance of the UK's wintering population.

BBS UK 1994-2012
Golden Plover





Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	16	1995-2011	62	-9	-31	11		
	10	2001-2011	64	4	-16	25		

Source	Period (yrs)	2006-2011 Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS Scotland	16	1995-2011	38	-18	-40	8		
	10	2001-2011	32	-2	-28	18		
	5	2006-2011	32	-14	-36	3		


Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.



Demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Clutch size	43	1968-2011	12	Linear decline	3.98 eggs	3.84 eggs	-3.6%		Small sample

For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).



Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts


Conservation listings:	Europe: SPEC category 2 (vulnerable) (BiE04) UK: red (BoCC3) UK Biodiversity Action Plan: priority species
Long-term trend:	UK: rapid decline England: moderate decline
Population size:	140,000 pairs in 2009 (APEP13: 1985-99 estimate (O'Brien 2004) updated using CBC/BBS trend)

Migrant status:	Short-distance migrant
Nesting habitat:	Ground nester
Primary breeding habitat:	Wetland
Secondary breeding habitat:	Farmland
Breeding diet:	Animal
Winter diet:	Animal

Status summary

Although CBC recorded some increase in its early years, Lapwings have declined continuously on lowland farmland since the mid 1980s. National surveys in England and Wales showed a 49% population decline between 1987 and 1998 (Wilson et al. 2001). Population declines of more than 50% over 15 years in Northern Ireland (Henderson et al. 2002) mirror similar declines throughout grassland areas of Wales and southeast England (Wilson et al. 2001, 2005). The BBS Austin et al. 2014); these birds are mainly of continental origin. Lapwing is one of the most strongly declining bird species in Europe, having decreased in all regions since 1980, although with differing regional timing (PECBMS 2009, 2013a). The 2009 review moved this species from amber to the UK red list, for which it qualifies on the strength of its UK decline.


CBC/BBS UK 1966–2012 Lapwing




Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	301	-54	-71	-29	>50	
	25	1986-2011	481	-60	-69	-47	>50	
	10	2001-2011	766	-34	-39	-26	>25	
	5	2006-2011	839	-35	-38	-28	>25	
CBC/BBS England	44	1967-2011	252	-36	-68	-7	>25	
	25	1986-2011	402	-50	-62	-37	>50	
	10	2001-2011	651	-26	-32	-19	>25	
	5	2006-2011	720	-31	-34	-25	>25	
WBS/WBBS waterways	31	1980-2011	69	-47	-72	-7	>25	
	25	1986-2011	76	-67	-79	-50	>50	
	10	2001-2011	111	-45	-52	-34	>25	
	5	2006-2011	102	-32	-42	-23	>25	
BBS UK	16	1995-2011	666	-41	-47	-33	>25	
	10	2001-2011	733	-33	-37	-25	>25	
	5	2006-2011	784	-32	-36	-26	>25	
BBS England	16	1995-2011	555	-24	-32	-15		
	10	2001-2011	622	-26	-30	-19	>25	
	5	2006-2011	671	-30	-34	-24	>25	
BBS Scotland	16	1995-2011	87	-56	-64	-44	>50	
	10	2001-2011	87	-40	-51	-25	>25	
	5	2006-2011	91	-35	-41	-23	>25	


Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




CBC/BBS UK graph




CBC/BBS England graph






BBS UK graph



BBS England graph




BBS Scotland graph


Demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Clutch size	43	1968-2011	195	Linear increase	3.71 eggs	3.80 eggs	2.6%		
Nest failure rate at egg stage	43	1968-2011	215	Curvilinear	1.58% nests/day	2.38% nests/day	50.6%		
Laying date	43	1968-2011	48	None			0 days		

For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).



Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

Causes of change

There is good evidence that declines have resulted from habitat loss and degradation due to changes in agricultural practice, in particular change from spring to autumn sowing, drainage of grasslands and loss of mixed farmland, which have led to breeding productivity dropping below a sustainable level. Chick mortality is thought to be the main determinant of poor Lapwing productivity, and therefore of population decline.

Change factor	Primary driver	Secondary driver
Demographic	Decreased breeding success	
Ecological	Agricultural intensification	

Further information on causes of change

Although there has been a slight increase in clutch size for this species, nest failures at the egg stage have increased dramatically (see above). Adult and first-year survival rates show no trend through time (Peach et al. 1994, Catchpole et al. 1999) and are unlikely to be the main driver of population declines (Sharpe et al. 2008). Chick mortality is thought to be the main determinant of poor Lapwing productivity and therefore of population decline (Sharpe et al. 2008), although evidence to support this is largely circumstantial and further empirical research is needed.

There is a good deal of research supporting the hypothesis that habitat loss and degradation due to the intensification of farming are likely to have been the main driver in the decline of this species, by reducing breeding productivity (e.g. Galbraith 1988, Shrubb 1990, Hotker 1991, Hudson et al. 1994, Siriwardena et al. 2000a, Taylor & Grant 2004, Wilson et al. 2005, Milsom 2005, Fuller & Ausden 2008). These changes include extensive drainage, increased use of pesticides and fertilisers, re-seeding, earlier and more frequent mowing, increased grazing pressure and loss of spring cereals. Increases in intensity of grazing have reduced the habitat quality for Lapwing (Shrubb 1990, Fuller & Ausden 2008), whilst fertilisation has led to earlier spring grass growth, earlier cutting dates and higher stocking levels, which have increased egg and chick mortality and reduced relaying opportunities (Durant et al. 2008). Drainage and loss of wet features on grassland have also had a negative impact, reducing food supplies (Taylor & Grant 2004, Eglington et al. 2010).

Loss of mixed farming systems and extensive grazing have reduced the availability of high-quality foraging habitat close to nesting habitat, i.e. unimproved pasture and meadows, to birds breeding in arable areas, resulting in reduced breeding success (Galbraith 1988, Hudson et al. 1994, Henderson et al. 2004).

In the uplands, afforestation has also resulted in habitat loss (Fuller & Ausden 2008). On arable land, spring-sown cereals were once favoured nesting crops but these have been widely replaced by autumn-sown cereals, which are less suitable breeding habitats (Shrubb 1990, Shrubb et al. 1991, Mason & Macdonald 1999, Fuller & Ausden 2008).

Lapwing population declines may be explained partly by increased nest predation rates resulting from habitat changes due to agricultural intensification (Baines 1990, Liker & Szekely 1997, Jackson & Green 2000, Chamberlain & Crick 2003, Evans 2004, Jackson et al. 2004, Milsom 2005, Bolton et al. 2007, Teunissen et al. 2008, MacDonald & Bolton 2008b, Bellebaum & Bock 2009). Long-term nest record card analysis has shown that the proportion of nests lost to predators was substantially higher in the 1990s than in previous decades (Sharpe et al. 2008). Recent empirical evidence suggests that levels of predation on wader nests are unsustainably high in many cases, even in some situations where breeding habitat is otherwise favourable (MacDonald & Bolton 2008a).

Declines among Lapwings are unlikely to be ameliorated by either habitat improvement or predator control in isolation (Bodey et al. 2011, Smart et al. 2013).

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Ringed Plover

Charadrius hiaticula

Key facts

Conservation listings:	Europe: no SPEC category (concentrated in Europe, conservation status favourable) (BiE04) UK: amber (species level and race <i>hiaticula</i> , 25-50% decline, >20% East Atlantic Flyway population in winter) (BoCC3)
Long-term trend:	UK: decline
Population size:	5,400 (5,300-5,600) pairs in 2007 (APEP13: Conway et al. 2008)

Status summary

The breeding population is monitored at intervals by special surveys. A BTO survey in 1984 showed increases throughout the UK since the previous survey in 1973-74 (Prater 1989). The spread of the breeding distribution inland between the first two atlas periods, especially in England, was probably associated with the increase in number of gravel pits and reservoirs (Gibbons et al. 1993). The 1984 survey revealed that over 25% of the UK population nested on the Western Isles, especially on the machair, but breeding waders there have subsequently suffered greatly from predation by introduced hedgehogs (Jackson et al. 2004) - a problem that appears increasingly severe (Jackson 2007). Surveys in England and Wales revealed an increase of 12% in breeding birds in wet meadows between 1982 and 2002 (Wilson et al. 2005). The BTO's repeat national survey in 2007 found an overall decrease in UK population of around 37% since 1984, with the greatest decreases in inland areas (Burton & Conway 2008, Conway et al. 2008, Conway & Burton 2009; [click here](#)). Ringed Plovers that choose beaches for nesting are especially vulnerable to disturbance, however, and already in 1984 were largely confined in some regions to wardened reserves (Prater 1989). Human usage of beach areas severely restricts the availability of this habitat to nesting plovers (Liley & Sutherland 2007). The marked increase in nest failures at the egg stage has earned Ringed Plover a place on the NRS concern list (Leech & Barimore 2008). Wintering numbers have been in decline since the late 1980s (Austin et al. 2014).


Population changes in detail

Annual breeding population changes for this species are not currently monitored by BTO


Demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Clutch size	43	1968-2011	92	None					
Nest failure rate at egg stage	43	1968-2011	126	Linear increase	2.22% nests/day	3.26% nests/day	46.8%		
Laying date	43	1968-2011	40	None			0 days		

For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).



Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>


Key facts

Conservation listings:	Global: Near Threatened (BiE04) UK: amber (>20% of European breeding and winter populations) (BoCC3) UK Biodiversity Action Plan: priority species
Long-term trend:	England: moderate decline
Population size:	68,000 pairs in 2009 (APEP13: 1985-99 estimate (O'Brien 2004) updated using CBC/BBS trend)

Status summary

Curlews monitored by CBC were mostly in lowland habitats and may have been affected primarily by drainage of farmland (Gibbons et al. 1993). Surveys of breeding birds in wet meadows in England and Wales revealed a decrease of 39% between 1982 and 2002 (Wilson et al. 2005). A 2006 survey in Wales highlighted the rapid decline of the species across all habitats, with low breeding success as a plausible mechanism (Johnstone et al. 2007). In Northern Ireland, a breeding decline of around 60% occurred between the mid 1980s and 1999 (Henderson et al. 2002). The BBS Austin et al. 2014). There has been widespread moderate decline across Europe since 1980 (PECBMS 2013a).

CBC/BBS England 1966-2012
Curlew



Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	44	1967-2011	136	-39	-71	24		Small CBC sample
	25	1986-2011	227	-28	-57	4		Small CBC sample
	10	2001-2011	368	-23	-29	-17		
	5	2006-2011	432	-13	-18	-7		

WBS/WBBS waterways Source	Period (yrs)	1980-2011 Years	# of Plots (n)	% Change	Lower limit	Upper limit	Alert	Comment
	25	1986-2011	51	-37	-55	-5	>25	
	10	2001-2011	77	-44	-52	-33	>25	
	5	2006-2011	74	-24	-33	-15		
BBS UK	16	1995-2011	510	-45	-51	-39	>25	
	10	2001-2011	545	-33	-41	-26	>25	
	5	2006-2011	612	-13	-20	-6		
BBS England	16	1995-2011	326	-32	-38	-23	>25	
	10	2001-2011	368	-23	-30	-16		
	5	2006-2011	432	-13	-18	-8		
BBS Scotland	16	1995-2011	120	-56	-63	-47	>50	
	10	2001-2011	115	-40	-50	-30	>25	
	5	2006-2011	118	-10	-22	7		
BBS Wales	16	1995-2011	35	-58	-70	-43	>50	
	10	2001-2011	33	-43	-57	-25	>25	
	5	2006-2011	30	-24	-42	-3		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




CBC/BBS England 1966–2012
Curlew




CBC/BBS England graph

WBS/WBBS 1979–2012
Curlew






WBS/WBBS waterways graph

BBS UK 1994–2012
Curlew




BBS UK graph




Demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Clutch size	43	1968-2011	20	None					Small sample
Nest failure rate at egg stage	43	1968-2011	22	Curvilinear	2.80% nests/day	2.91% nests/day	3.9%		Small sample

For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).



Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglinton, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Common Sandpiper

Actitis hypoleucos

Key facts


Conservation listings:	Europe: SPEC category 3 (declining) (BiE04) UK: amber (25-50% population decline) (BoCC3)
Long-term trend:	UK waterways: moderate decline
Population size:	15,000 pairs in 2009 (APEP13: previous estimate (Dougall et al. 2004) updated using WBS/WBBS trend)

Status summary

WBS/WBBS results for this species show a decline from 1985 onwards (after a more gradual increase) that has yet to be explained, though decreases are less evident in BBS squares. Poorer breeding success and reduced survival of first-year birds over the winter in West Africa were both suggested as possible reasons for the failure of the Peak District population to recover after a hard-weather event in 1989 (Holland & Yalden 2002). The reasons for poor recruitment to the breeding population are hard to assess in the absence of firm information on where British birds spend the winter (Dougall et al. 2010). UK clutch sizes appear to have shown a slight decline since the 1960s. Following declines during the 1990s in the large Swedish and Finnish populations, the European status of this species is no longer considered 'secure' (BirdLife International 2004). Widespread moderate decline has occurred across Europe since 1980 (PECBMS 2013a). The species was moved from the green to the amber list in 2009 on the strength of its declines in UK and across Europe.

WBS/WBBS 1974–2012

Common Sandpiper




Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
WBS/WBBS waterways	36	1975-2011	47	-44	-58	-32	>25	
	25	1986-2011	57	-49	-60	-40	>25	
	10	2001-2011	84	-22	-32	-9		

Source BBS UK	5 Period (yrs)	2006-2011 Years 1995-2011	77 Plots (#)	-15 Change (%)	-25 Lower limit	-6 Upper limit	Alert	Comment
	10	2001-2011	66	3	-17	26		
	5	2006-2011	74	4	-16	26		
BBS Scotland	16	1995-2011	33	-8	-29	16		
	10	2001-2011	30	3	-18	25		
	5	2006-2011	33	2	-22	25		


Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.



WBS/WBBS waterways graph



BBS UK graph




BBS Scotland graph


Demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Clutch size	43	1968-2011	12	Curvilinear	3.98 eggs	3.85 eggs	-3.3%		Small sample
Nest failure rate at egg stage	43	1968-2011	13	None					Small sample

For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).



Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts


Conservation listings:	Europe: SPEC category 2 (declining) (BiE04) UK: amber (>50% population decline but data possibly unrepresentative, >20% of East Atlantic Flyway population in winter) (BoCC3)
Long-term trend:	UK: decline UK waterways: rapid decline
Population size:	25,000 pairs in 2009 (APEP13: 1985-99 estimate (O'Brien 2004) updated using CBC/BBS trend)

Status summary

In 2009, UK population decline was added to the criteria by which Redshank qualifies for amber listing; the scale of decline reported here already meets the red-list criterion, however. Range contraction had occurred from considerable areas of the UK by 1988-91, probably as a result of the drainage of farmland (Gibbons et al. 1993). WBS/WBBS results show a decline along waterways that apparently accelerated during the 1990s. BBS shows continuing overall decrease. Surveys in England and Wales revealed a decrease of 29% in breeding birds in wet meadows between 1982 and 2002 (Wilson et al. 2005). The substantial section of the British population that nests on saltmarshes decreased by 23% between 1985 and 1996, apparently as a result of increased grazing pressure (Brindley et al. 1998, Norris et al. 1998). By 2011, fewer than 12,000 breeding pairs remained on saltmarshes, a decrease of 53% from the 1985 survey: a better understanding of saltmarsh grazing practices and longer-term management of this habitat is urgently needed (Malpas et al. 2013). Wintering populations (augmented by many Icelandic and some other northern European breeders) have shown some increase since the 1970s but have been in decline since about 2001 (Austin et al. 2014). The success of nests at the egg stage has risen steeply since the 1960s. Numbers have shown widespread moderate decrease across Europe since 1980 (PECBMS 2013a). In Scotland at least, agri-environment schemes can benefit this species (O'Brien & Wilson 2011).

WBS/WBBS 1974-2012

Redshank




Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
WBS/WBBS waterways	36	1975-2011	24	-63	-89	-36	>50	

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
	10	2001-2011	31	-39	-56	-20	>25	
	5	2006-2011	27	-24	-44	-3		
BBS UK	16	1995-2011	84	-42	-56	-25	>25	
	10	2001-2011	93	-30	-44	-13	>25	
	5	2006-2011	101	-27	-41	-13	>25	
BBS England	16	1995-2011	60	-32	-49	-8	>25	
	10	2001-2011	68	-40	-53	-22	>25	
	5	2006-2011	75	-23	-38	-3		


Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.



WBS/WBBS waterways graph



BBS UK graph




BBS England graph


Demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Clutch size	43	1968-2011	29	None					Small sample
Nest failure rate at egg stage	43	1968-2011	30	Curvilinear	4.46% nests/day	2.06% nests/day	-53.8%		

For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).



Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Woodcock


Scolopax rusticola

Key facts

Conservation listings:	Europe: SPEC category 3 (declining) (BiE04) UK: amber (European status) (BoCC3)
Long-term trend:	UK: probable rapid decline
Population size:	81,000 (64,000-100,000) males in 2003 (APEP13: Hoodless et al. 2009)

Status summary

The Woodcock declined rapidly and significantly on CBC plots for the three decades up to 2000. Because CBC did not include many coniferous forests and its plots were concentrated in lowland Britain, however, it is not certain how clearly this trend represents the whole UK population. Range contractions, that might have had the same cause as the decline in abundance, were recorded concurrently with part of the CBC decline (Gibbons et al. 1993). Recreational disturbance, the drying out of natural woodlands, overgrazing by deer, declining woodland management, and the maturation of new plantations are possible causes of the Woodcock's decline, but there is no strong hypothesis as yet (Fuller et al. 2005). BBS is inefficient at recording this scarce, mainly crepuscular species, and cannot continue the index series. The first special survey aimed at monitoring the UK's breeding Woodcock took place in 2003 and has provided a new, much higher baseline population estimate for future monitoring (Hoodless et al. 2009; also, BoCC3), which now rests on the breeding declines recorded across Europe, especially European Russia (BiE04). Annual [humbers shot](#) in the UK, which include winter visitors from declining populations in Europe, have increased around threefold since 1945 and are currently running at a historically high level.



Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC all habitats	31	1968-1999	20	-74	-88	-49	>50	Small CBC sample
	25	1974-1999	20	-76	-88	-51	>50	Small CBC sample
	10	1989-1999	13	-40	-62	-11	>25	Small CBC sample
	5	1994-1999	13	-24	-44	-3		Small sample

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.



CBC all habitats graph

Demographic trends

Productivity and survival trends for this species are not currently produced by BTO


This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglinton, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts

Conservation listings:	Europe: SPEC category 3 (declining) (BiE04) UK: amber (European status) (BoCC3)
Long-term trend:	UK waterways: rapid decline
Population size:	80,000 pairs in 2009 (APEP13: 1985-99 estimate (O'Brien 2004) updated using CBC/BBS trend)

Status summary

Snipe were monitored by the CBC mainly in lowland England, where numbers have fallen rapidly since the 1970s as farmland has been drained (Gibbons et al. 1993, Siriwardena et al. 2000a). The CBC index fell from the early 1970s until 1984, when the number of occupied plots became too small for further monitoring (Marchant et al. 1990), and the graph is not included here. In Northern Ireland, a breeding decline of around 30% occurred between the mid 1980s and 1999 (Henderson et al. 2002). Surveys in England and Wales revealed a decrease of 62% in breeding birds in wet meadows between 1982 and 2002, with the remaining birds becoming highly aggregated into a tiny number of suitable sites (Wilson et al. 2005). Birds were more likely to persist where soils remained soft and wet; the fact that Snipe have continued to decline, despite soil conditions being improved for them at many lowland wetland reserves, suggests that other key aspects of habitat quality, such as prey abundance, are more likely to be driving the decline (Smart et al. 2008). The trend in the upland and moorland strongholds of the species is not fully known, but the 1988-91 atlas documented range loss widely in Wales, Northern Ireland and Scotland, as well as lowland England, and atlas work during 2008-11 confirmed that range loss or population decrease has been evident almost everywhere (Balmer et al. 2013). The BBS showed initial increases from 1994, especially in Scotland, but a sharp downturn over the recent decade. Daily nest failure rates at the egg stage appear to have halved. Following moderate decline across Europe since 1980 (PECBMS 2013a), this previously 'secure' species is now provisionally evaluated as 'declining' (BirdLife International 2004). In Scotland at least, agri-environment schemes can benefit this species (O'Brien & Wilson 2011).



Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
--------	--------------	-------	-----------	------------	-------------	-------------	-------	---------

WBS/WBBS waterways Source	Period (yrs)	1975-2011 Years	Plots (n)	Change (%)	Lower limit	Upper limit	>50 Alert	Small sample Comment
	25	1986-2011	15	-91	-98	-80	>50	Small sample
	10	2001-2011	23	-50	-71	-20	>25	
	5	2006-2011	21	-29	-58	-3	>25	
BBS UK	16	1995-2011	160	8	-13	38		
	10	2001-2011	184	-22	-32	-8		
	5	2006-2011	211	-25	-31	-13		
BBS England	16	1995-2011	86	-20	-34	0		
	10	2001-2011	106	-32	-43	-16	>25	
	5	2006-2011	130	-26	-34	-13	>25	
BBS Scotland	16	1995-2011	55	15	-8	51		
	10	2001-2011	56	-19	-33	-1		
	5	2006-2011	59	-23	-30	-11		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




WBS/WBBS 1974-2012
Snipe




WBS/WBBS waterways graph

BBS UK 1994-2012
Snipe




BBS UK graph

BBS England 1994-2012
Snipe



BBS England graph

BBS Scotland 1994-2012
Snipe




BBS Scotland graph


Demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Clutch size	43	1968-2011	12	None					Small sample
Nest failure rate at egg stage	43	1968-2011	14	Linear decline	3.18% nests/day	1.29% nests/day	-59.4%		Small sample

For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).



Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts


Conservation listings:	Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe) (BiE04) UK: green (Rock Dove C. l. livia) (BoCC3)
Long-term trend:	UK: possible increase
Population size:	550,000 (450,000-650,000) pairs in 2009 (APEP13: distance sampling estimate for 2006 (Newson et al. 2008) updated using BBS trend)

Status summary

CBC samples for Feral Pigeon were consistently too small for annual monitoring, and there was no trend information before BBS began in 1994. Breeding atlas data show a 39% increase in occupied 10-km squares between 1968-72 and 1988-91 (Gibbons et al. 1993) and a further 5% or so by 2008-11 (Balmer et al. 2013), suggesting that Feral Pigeons may be on an upward trajectory, like the other Columba species in the UK. At the time of the first atlas, however, Feral Pigeons were more commonly overlooked during bird surveys, and some of the reported subsequent range increase may have been due to greater observer awareness. It is now clear that Feral Pigeons are almost ubiquitous in the UK, nesting in rural as well as urban habitats, and avoiding only the highest ground. No distinction can realistically be drawn between feral birds of domestic origin and true wild-type Rock Doves, although birds of wild-type plumage still predominate on some more-remote Scottish islands. In field conditions, it is often not possible to distinguish between pure native Rock Doves, wild-nesting Feral Pigeons, semicaptive dove-cote breeders, and passing racing pigeons, nor between adults and young of the year, and BBS counts are likely to include birds from all of these groups. BBS indices suggest that a minor decrease has occurred in recent years.

BBS UK 1994–2012

Rock Dove/Feral pigeon




Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	16	1995-2011	679	-15	-24	-1		
	10	2001-2011	746	-19	-26	-10		

Source	Period (yrs)	2006-2011 Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS England	16	1995-2011	565	-23	-31	-13		
	10	2001-2011	615	-26	-33	-18	>25	
	5	2006-2011	669	-20	-27	-13		
BBS Scotland	16	1995-2011	62	7	-31	66		
	10	2001-2011	68	0	-22	32		
	5	2006-2011	75	5	-13	23		
BBS Wales	16	1995-2011	33	56	-6	106		
	10	2001-2011	38	5	-23	34		
	5	2006-2011	38	14	-3	39		


Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




BBS UK graph



BBS England graph



BBS Scotland graph



BBS Wales graph

Demographic trends

Productivity and survival trends for this species are not currently produced by BTO

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglinton, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>


Key facts

Conservation listings:	Europe: no SPEC category (concentrated in Europe, conservation status favourable) (BiE04) UK: amber (>20% of European breeding population) (BoCC3)
Long-term trend:	England: rapid increase
Population size:	260,000 territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend for England)

Status summary

Following release from the lethal and sublethal effects of the organochlorine seed-dressings used in the 1950s and early 1960s, Stock Dove populations have increased very substantially (O'Connor & Mead 1984). Numbers appeared to level off in the early 1980s, and entered a further increasing phase in the early 1990s which may now have ceased. The BBS Siriwardena et al. 2000b). Overall, nest failure rates have fallen substantially since the 1980s and there has been a major increase in the number of fledglings raised per breeding attempt. There has been moderate increase across Europe since 1980 but no detectable trend since 1990 (PECBMS 2013a).

CBC/BBS England 1966–2012 Stock Dove






Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	44	1967-2011	322	187	102	307		
	25	1986-2011	516	16	1	42		
	10	2001-2011	802	-3	-8	5		
	5	2006-2011	889	10	4	18		
BBS UK	16	1995-2011	766	11	2	23		


Source	10 Period (yrs)	2001-2011 Years	857 Plots	2 Change (%)	5 Lower limit	11 Upper limit	Alert	Comment
BBS England	16	1995-2011	705	9	-4	19		
	10	2001-2011	789	-2	-9	6		
	5	2006-2011	868	10	3	18		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.



Fledglings per breeding attempt


Stock Dove 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Stock Dove




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	72	Linear increase	0.98 fledglings	1.40 fledglings	42.4%		
Clutch size	43	1968-2011	116	Curvilinear	2.07 eggs	2.10 eggs	1.4%		
Brood size	43	1968-2011	176	Curvilinear	1.82 chicks	1.83 chicks	0.5%		
Nest failure rate at egg stage	43	1968-2011	109	Linear decline	1.57% nests/day	0.53% nests/day	-66.2%		
Nest failure rate at chick stage	43	1968-2011	72	Linear decline	1.14% nests/day	0.69% nests/day	-39.5%		
Laying date	43	1968-2011	24	None			0 days		Small sample


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

Key facts


Conservation listings:	Europe: no SPEC category (concentrated in Europe, conservation status favourable) (BiE04) UK: green (BoCC3)
Long-term trend:	UK, England: rapid increase
Population size:	5.4 (5.1-5.7) million pairs in 2009 (APEP13: distance sampling estimate for 2006 (Newson et al. 2008) updated using BBS trend)
Migrant status:	Resident
Nesting habitat:	Above-ground nester
Primary breeding habitat:	Farmland
Secondary breeding habitat:	
Breeding diet:	Vegetation
Winter diet:	Vegetation

Status summary

The CBC/BBS trend for this species is of a steady, steep increase since at least the mid 1970s. Since 1994, BBS has recorded significantly upward trends in the UK, and in England, Wales and Northern Ireland separately, but stability in Scotland. The BBS PECBMS 2013a).



CBC/BBS UK 1966–2012






Woodpigeon




Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	1004	173	55	490		
	25	1986-2011	1722	75	55	91		
	10	2001-2011	2801	22	18	26		
	5	2006-2011	3149	9	6	11		
CBC/BBS England	44	1967-2011	807	192	57	533		
	25	1986-2011	1384	84	63	110		
	10	2001-2011	2242	24	19	29		
	5	2006-2011	2542	8	5	11		
BBS UK	16	1995-2011	2438	40	33	47		
	10	2001-2011	2745	21	18	25		
	5	2006-2011	3057	9	6	11		
BBS England	16	1995-2011	1958	46	39	53		
	10	2001-2011	2206	24	20	28		
	5	2006-2011	2483	8	5	11		
BBS Scotland	16	1995-2011	196	6	-12	24		
	10	2001-2011	215	7	-6	18		
	5	2006-2011	241	13	3	22		
BBS Wales	16	1995-2011	186	41	17	65		
	10	2001-2011	210	28	10	52		
	5	2006-2011	212	9	-7	32		
BBS N.Ireland	16	1995-2011	83	93	50	145		
	10	2001-2011	99	21	9	34		
	5	2006-2011	104	15	2	24		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.






Fledglings per breeding attempt Woodpigeon 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Woodpigeon




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	77	Curvilinear	0.55 fledglings	0.74 fledglings	35.2%		
Clutch size	43	1968-2011	74	None					
Brood size	43	1968-2011	118	None					
Nest failure rate at egg stage	43	1968-2011	97	Curvilinear	4.48% nests/day	2.90% nests/day	-35.3%		
Nest failure rate at chick stage	43	1968-2011	78	None					
Laying date	43	1968-2011	87	Linear increase	Jun 1	Jun 23	22 days		


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

Causes of change

There is some evidence that the increase in this species has been due to the spread of intensive winter cereal and rape cultivation, probably by increasing food availability over winter, reflecting the species' ability to subsist on green vegetation, unlike other granivores.

Change factor	Primary driver	Secondary driver
Demographic	Increased overwinter survival	
Ecological	Agricultural intensification	

Further information on causes of change

There are few studies specifically examining demographic and ecological drivers of the long-term increase in this species but the spread of intensive arable cultivation, especially of oilseed rape and winter-sown cereal, which has been shown to reduce overwinter mortality, may explain the rise in numbers (Gibbons et al. 1993, Inglis et al. 1997). Inglis et al. (1997) conducted fieldwork to provide good evidence that, in their study area in Cambridgeshire, the overwintering population size was determined by the area of oilseed rape. Inglis et al. state that, since the introduction of oilseed rape, the number of fledged young produced has a more important effect upon the Woodpigeon population size than does overwinter mortality from starvation, i.e. winter food availability no longer limits the population.

The number of Woodpigeons feeding in gardens has also increased (Glue 1993, 1995, 1997), suggesting that this species may benefit from the trend of increasing urban feeding sites, although there is no direct evidence to support this.

The species is adaptable and O'Connor & Shrubbs (1986) found that the breeding season had advanced in response to the switch to autumn sowing, and thus earlier ripening, of cereals, with more pairs nesting in May and June and relatively fewer during July-September. Climate change may have also permitted earlier nesting. A trend toward earlier nesting could have led CBC, with its fieldwork finishing in early July, to overestimate the rate of increase (Marchant et al. 1990). Newly available data indicate, however, that the species is now nesting almost three weeks later, on average, than it did in the 1960s.

Collared Dove

Streptopelia decaocto

Key facts


Conservation listings:	Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe) (BiE04) UK: green (BoCC3)
Long-term trend:	UK: rapid increase
Population size:	990,000 (900,000-1,090,000) pairs in 2009 (APEP13: distance sampling estimate for 2006 (Newson et al. 2008) updated using BBS trend)
Migrant status:	Resident
Nesting habitat:	Above-ground nester
Primary breeding habitat:	Human habitats
Secondary breeding habitat:	
Breeding diet:	Vegetation
Winter diet:	Vegetation

Status summary

Collared Dove abundance has increased rapidly since the species first colonised Britain in 1955. From just four birds known to be present in that year, the population was put conservatively at 15,000-25,000 pairs by 1970 (Hudson 1972). The CBC index showed an almost exponential rise as colonisation continued during the early 1970s, but the CBC index had levelled off by about 1980. BBS shows continuing increases, at least in England and Wales, with a recent downturn nationally to pre-2000 population levels. The BBS PECBMS 2013a).




CBC/BBS UK 1971-2012


Collared Dove




Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1972-2011	630	372	201	552		
	25	1986-2011	939	63	33	93		
	10	2001-2011	1539	-2	-6	2		
	5	2006-2011	1713	-9	-12	-5		
BBS UK	16	1995-2011	1345	19	12	26		
	10	2001-2011	1529	-1	-5	3		
	5	2006-2011	1696	-8	-11	-4		
BBS England	16	1995-2011	1181	18	11	26		
	10	2001-2011	1332	-4	-7	1		
	5	2006-2011	1479	-9	-12	-6		
BBS Scotland	16	1995-2011	50	-9	-33	31		
	10	2001-2011	59	17	-9	53		
	5	2006-2011	67	-6	-21	14		
BBS Wales	16	1995-2011	73	37	-1	84		
	10	2001-2011	85	6	-5	23		
	5	2006-2011	91	-13	-20	-2		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.






BBS Scotland graph



BBS Wales graph

Demographic trends


Fledglings per breeding attempt Collared Dove 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Collared Dove




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	54	Curvilinear	0.81 fledglings	0.91 fledglings	11.2%		
Clutch size	43	1968-2011	43	Linear decline	1.95 eggs	1.88 eggs	-3.5%		
Brood size	43	1968-2011	72	None					
Nest failure rate at egg stage	43	1968-2011	61	Curvilinear	3.14% nests/day	2.79% nests/day	-11.1%		
Nest failure rate at chick stage	43	1968-2011	54	None					
Laying date	43	1968-2011	43	None			0 days		


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

Causes of change

There is little evidence available relating to the drivers of the increase in this species but it appears to have been able to fill an empty niche and exploit the intermittent seed resources available in gardens and may also benefit from milder winters. Given the long-term rise, there is no baseline of 'stability' against which to compare demographic rates that might be causing a change but there have been increases in nesting productivity.

Change factor	Primary driver	Secondary driver
Demographic	Increased breeding success	
Ecological	Other	Climate change

Further information on causes of change

There are very few studies from the UK looking at the causes of population change in Collared Dove. There has been possible increase in brood size and a decline in daily failure rate at the egg stage (see graphs above). The species appears to have filled a previously empty niche, perhaps because it is able to adapt to new environments, and it has shown a rapid [increase in gardens](#), exploiting the intermittent seed resources available. It may also benefit from milder winters, which the species can exploit with its long breeding seasons. However, evidence for this is anecdotal.

Robertson (1990) measured high productivity and a long breeding season in rural Collared Doves in Oxfordshire and suggested that these were made possible by feeding on superabundant, predictable and persistent supplies of commercial crop seed in and around farmyards. However, there is little evidence based on specific analyses to support this.

There is evidence that the recent slowing of population increase may be due to increasing numbers of grey squirrels, as Newson et al. (2010b) provided good evidence from nest record data which showed a positive relationship between nest failure at the egg stage and squirrel abundance. They may also have been approaching the saturation of their niche. The outbreak of trichomonosis first noted in 2006 is thought to have affected this species quite severely and may be the primary cause of the current downturn. Population trends have been different in Scotland but the reasons for this are unclear.

Key facts


Conservation listings:	Europe: SPEC category 3 (declining) (BiE04) UK: red (>50% population decline) (BoCC3) UK Biodiversity Action Plan: click here , priority species
Long-term trend:	UK, England: rapid decline
Population size:	14,000 territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)
Migrant status:	Long-distance migrant
Nesting habitat:	Above-ground nester
Primary breeding habitat:	Farmland
Secondary breeding habitat:	
Breeding diet:	Vegetation
Winter diet:	Vegetation

Status summary

The CBC/BBS trend shows severe declines in Turtle Dove abundance, beginning in the late 1970s and continuing steeply to the present. Atlas data show that more than half the 10-km squares occupied in 1968-72 had been lost by 2008-11, with the population withdrawing towards East Anglia and Kent (Balmer et al. 2013). These trends, if not reversed, would bring the species close to extinction in the UK within the next two decades. Turtle Dove has been one of the most strongly declining bird species across Europe since 1980 (PECBMS 2013a).

CBC/BBS UK 1966–2012

Turtle Dove







Smoothed population index, relative to an arbitrary 100 in the year given, with 85% confidence limits in green

Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	106	-94	-97	-91	>50	
	25	1986-2011	132	-91	-94	-87	>50	
	10	2001-2011	133	-77	-81	-72	>50	
	5	2006-2011	113	-63	-69	-57	>50	
CBC/BBS England	44	1967-2011	105	-94	-96	-91	>50	
	25	1986-2011	130	-91	-93	-88	>50	
	10	2001-2011	131	-78	-81	-73	>50	
	5	2006-2011	111	-63	-68	-58	>50	
BBS UK	16	1995-2011	158	-85	-88	-82	>50	
	10	2001-2011	132	-77	-81	-72	>50	
	5	2006-2011	112	-63	-68	-56	>50	
BBS England	16	1995-2011	156	-85	-88	-81	>50	
	10	2001-2011	130	-78	-81	-72	>50	
	5	2006-2011	110	-63	-68	-57	>50	


Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.





Demographic trends


Fledglings per breeding attempt Turtle Dove 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Turtle Dove




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	11	None					
Fledglings per breeding attempt	43	1968-2011	11	None					
Clutch size	43	1968-2011	11	None					Small sample
Brood size	43	1968-2011	15	Curvilinear	1.83 chicks	1.73 chicks	-5.4%		Small sample
Nest failure rate at egg stage	43	1968-2011	15	None					Small sample
Nest failure rate at chick stage	43	1968-2011	11	None					Small sample
Laying date	43	1968-2011	12	None			0 days		Small sample


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

Causes of change

There is good evidence to support the hypothesis that the primary demographic driver of Turtle Dove declines is a shortened breeding period, which has reduced the number of nesting attempts. This is thought to be driven by reduced food availability due to increased herbicide use, although analyses that test this directly are lacking. Note, however, that data do not permit analyses of variation in annual survival rates, but mortality both on the wintering grounds (due to habitat deterioration) and on migration (particularly through hunting) could be important.

Change factor	Primary driver	Secondary driver
Demographic	Reduced breeding success	
Ecological	Agricultural intensification	

Further information on causes of change

A four-year intensive field study in East Anglia provided good evidence that the role of breeding productivity in the decline of Turtle Doves is likely to be through a reduction in the average number of nesting attempts per pair (Browne & Aebischer 2005). Browne & Aebischer (2003, 2004, 2005) concluded that Turtle Doves today have a substantially earlier close to the breeding season and consequently produce fewer clutches and young per pair than they did in the 1960s. Reduced food availability due to increased herbicide use and efficacy may make birds more likely to cease breeding earlier than during the 1960s and reduce their number of nesting attempts (Browne & Aebischer 2001, 2002), although this was not specifically tested. Browne & Aebischer (2003) state that it may be a change in phenology of Turtle Doves and their food species which has resulted in reduced availability of food supplies, although they do not support this with any specific analyses of these two factors. Loss of quality and quantity of breeding habitat are also thought to contribute to declines. Browne et al. (2004) used long-term CBC data to provide good evidence that breeding density fell in proportion to loss of nesting, rather than feeding, habitat and that changes in Turtle Dove density were positively related to changes in the amount of hedgerow and woodland edge. Dunn & Morris (2012) suggest however that, although established scrub and large hedgerows were important in retaining Turtle Dove territories, it may be foraging habitat that is limiting their distribution.

There is good evidence to suggest that the population decline experienced by Turtle Doves breeding in Britain is not due to lower success of individual nesting attempts. Analysis of nest record cards and ringing data for farmland Turtle Doves shows a non-significant increase in productivity per nesting attempt while annual survival has fallen (Siriwardena et al. 2000a, 2000b, Browne et al. 2005) so this may have also contributed to the decline. The demographic trends shown here support the view that nesting success per attempt is not the main driver of population change, with only a slight decrease in brood size being reported (see above).

Turtle Dove is a quarry species in many European countries and hunting during migration has been cited as another possible cause of the UK decline, although there is little evidence to support this (Browne & Aebischer 2004). Ring-recovery sample sizes are small and there is only weak evidence suggesting a decrease in annual survival (Siriwardena et al. 2000b). Nevertheless, survival could also have been negatively affected by a reduction in the quality of wintering habitat: this is thought to have

contributed to the decline (Marchant et al. 1990) and one recent study has demonstrated a positive correlation between survival rate among breeding adults in France and food supply in West Africa, as measured by cereal production (Eraud et al. 2009). Further work on the ecology of Turtle Doves on their wintering grounds is needed to investigate the relevance of this result for UK birds.

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>


Key facts

Conservation listings:	Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe) (BiE04) UK: red (BoCC3) UK Biodiversity Action Plan: priority species
Long-term trend:	England: rapid decline
Population size:	16,000 (9,000-24,000) pairs in 2009 (APEP13: distance sampling estimate for 2006 (Newson et al. 2008) updated using BBS trend)
Migrant status:	Long-distance migrant
Nesting habitat:	Host-specific
Primary breeding habitat:	Woodland
Secondary breeding habitat:	
Breeding diet:	Animal
Winter diet:	Animal

Status summary

The CBC/BBS trend shows Cuckoo abundance to have been in decline since the early 1980s. The species was moved in 2002 from the green to the amber list, and in the 2009 review met red-list criteria. The sensitivity of CBC to change in this species may have been relatively low, mainly because Cuckoo territories were typically larger than census plots (Marchant et al. 1990). BBS shows a continuing strong decline in England and Wales, but not in Scotland, where numbers appeared until very recently to be stable. The BBS Newson et al. 2009). There has been widespread moderate decrease across Europe since 1980, but with little change since 1990 (PECBMS 2013a).

CBC/BBS England 1966–2012 Cuckoo



Smoothed population index, relative to an arbitrary 100 in the year given, with 85% confidence limits in green


Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	44	1967-2011	292	-73	-81	-63	>50	
	25	1986-2011	437	-72	-76	-67	>50	
	10	2001-2011	532	-44	-49	-40	>25	
	5	2006-2011	553	-27	-32	-23	>25	
BBS UK	16	1995-2011	721	-50	-55	-46	>50	
	10	2001-2011	686	-32	-38	-27	>25	
	5	2006-2011	712	-21	-28	-15		
BBS England	16	1995-2011	562	-65	-67	-62	>50	
	10	2001-2011	519	-43	-47	-38	>25	
	5	2006-2011	530	-26	-31	-22	>25	
BBS Scotland	16	1995-2011	73	-5	-25	20		
	10	2001-2011	75	-13	-31	6		
	5	2006-2011	86	-17	-32	-2		
BBS Wales	16	1995-2011	56	-34	-48	-21	>25	
	10	2001-2011	57	-19	-31	-7		
	5	2006-2011	56	-7	-19	10		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




CBC/BBS England 1966-2012
Cuckoo




CBC/BBS England graph

BBS UK 1994-2012
Cuckoo




BBS UK graph


BBS England 1994-2012
Cuckoo



BBS England graph



BBS Scotland graph



BBS Wales graph

Demographic trends

Productivity and survival trends for this species are not currently produced by BTO

Causes of change

It is unclear what is the main driver of population decline in Cuckoos. Given the lack of demographic trends for this species it is not possible to identify a specific mechanism behind the declines.

Change factor	Primary driver	Secondary driver
Demographic	Unknown	
Ecological	Unknown	

Further information on causes of change

Cuckoo abundance may be related to their breeding success, which might in turn be determined by the abundance of breeding success of host species. Evidence from BBS data show strong variation in Cuckoo population trends between habitats, which may reflect regional differences in the main hosts and differing trends in Cuckoo breeding success among those host species (Newson et al. 2009). Douglas et al. (2010b) found a strong positive correlation between change in Cuckoo numbers and numbers of Brooke & Davies 1987) but the authors also thought that this was unlikely to be the main cause of population decline. There has perhaps been a disproportionate emphasis on the role of brood parasitism aspects in Cuckoo decline.

Another hypothesis for the decline of Cuckoos relates to phenological mismatch in the timing of host and Cuckoo breeding. There is evidence relating to climate-induced changes in phenology, although the extent to which this may be driving population declines is unclear. Douglas et al. (2010b) used BBS data and found that in recent decades, earlier breeding Douglas et al. 2010b). In Europe, other recent studies have suggested that climate change might disrupt the association between the life cycles of the Cuckoo and its short-distance migrant hosts and they state that this mismatch may contribute to the decline in Cuckoo (Saino et al. 2009, Moller et al. 2011). Thus, evidence at European scale at least is equivocal.

Given that the Cuckoo is a migrant, and the fact that many long-distance migrants have been found to be declining (Sanderson et al. 2006, Hewson & Noble 2009), factors operating on wintering grounds have been suggested as a possible primary driver of Cuckoo declines (Glue 2006, Payevsky 2006, Newson et al. 2009). However, little work has focused on this area to date. Decreased food supplies on the breeding grounds has also been suggested as a possible cause (Glue 2006), following the rapid declines of many British moth species (Conrad et al. 2006), important prey items in Cuckoo diet, but detailed research on this is lacking.

Key facts


Conservation listings:	Europe: SPEC category 3 (declining) (BiE04) UK: amber (25-50% distribution decline) (BoCC3)
Long-term trend:	UK: probable decline
Population size:	4,000 (3,000-5,000) pairs in 1995-97 (APEP13: Toms et al. 2001)

Status summary

Distributional data provide good evidence for a decline in this species that lasted throughout the 20th century, although annual monitoring started only very recently. Productivity has tended to improve since the 1950s and 1960s, when Barn Owls appear to have been affected by organochlorine pesticides (Percival 1990). A national census during 1995-97, organised jointly by Hawk & Owl Trust and BTO, provided a replicable baseline population estimate (Toms et al. 2000, 2001; for more information, clickDadam et al. 2011). In earlier decades, the plight of such a charismatic and popular bird led to extensive releasing of captive-bred birds in unguided attempts at restocking: by 1992, when licensing became a requirement for such schemes, it was estimated that between 2,000 and 3,000 birds were being released annually by about 600 operators, although many birds died quickly and never joined the nesting population (Balmer et al. 2000). There is some evidence that releases may have aided population recovery (Meek et al. 2003). Alongside the restocking, the erection of Barn Owl nest boxes, already numbering c. 25,000 by the mid 1990s, has enabled the species to occupy areas (notably the Fens) that were previously devoid of nesting sites, and may have been a factor in the improving nesting success. Numbers of Barn Owls recorded via BBS have increased strongly since 1995 and reached a peak around 2009. As BBS is a diurnal survey, the detectability of primarily nocturnal species is low and could be influenced quite markedly by changes in behaviour: thus the trends should be interpreted with extra care. The number of nest records for Barn Owl has also increased rapidly over the same period, strengthening the evidence that a national population increase has indeed occurred since Project Barn Owl in 1995-97. There is likely to be some regional variation in population trends, however. RBBP provide a county breakdown of 2005 nesting totals (Holling & RBBP 2008).

BBS UK 1994–2012

Barn Owl





Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
--------	--------------	-------	-----------	------------	-------------	-------------	-------	---------

BBS UK Source	Period (yrs)	1995-2011 Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
	10	2001-2011	64	105	48	178		
	5	2006-2011	79	-9	-20	7		
BBS England	16	1995-2011	43	285	181	469		
	10	2001-2011	61	104	52	162		
	5	2006-2011	77	-10	-23	5		


Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.



Demographic trends

Fledglings per breeding attempt


Barn Owl 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Barn Owl




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	30	Linear increase	2.33 fledglings	3.25 fledglings	39.7%		
Clutch size	43	1968-2011	44	Curvilinear	4.29 eggs	4.85 eggs	13.1%		
Brood size	43	1968-2011	360	Curvilinear	3.05 chicks	2.95 chicks	-3.3%		
Nest failure rate at egg stage	43	1968-2011	30	Linear decline	0.81% nests/day	0.05% nests/day	-93.8%		Small sample
Nest failure rate at chick stage	43	1968-2011	136	Linear decline	0.21% nests/day	0.02% nests/day	-90.5%		
Laying date	43	1968-2011	18	None			0 days		Small sample


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

Key facts


Conservation listings:	Europe: SPEC category 3 (declining) (BiE04) UK: not listed (introduced)
Long-term trend:	UK, England: rapid decline
Population size:	5,700 (3,700-7,700) pairs in 2009 (distance sampling estimate for 2006 (Newson et al. 2008) updated using BBS trend)
Migrant status:	Resident
Nesting habitat:	Cavity nester
Primary breeding habitat:	Farmland
Secondary breeding habitat:	
Breeding diet:	Animal
Winter diet:	Animal

Status summary

The CBC/BBS trend for Little Owl in the UK shows very wide variation, but a downturn in recent decades suggests that a rapid decline now lies behind the observed fluctuations. Trends are unusually uncertain, however, because the species has large breeding territories and, being largely inactive during the day, is difficult to detect except by dedicated surveys. A figure of c. 7,000 pairs from the BTO/Hawk & Owl Trust's Toms et al. 2000) was the first replicable population estimate for Little Owls in the UK. An independent BBS estimate is for c5,700 pairs in 2009, since when further decrease is apparent.




CBC/BBS UK 1966–2012


Little Owl



Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	60	-64	-80	-40	>50	
	25	1986-2011	83	-65	-75	-53	>50	
	10	2001-2011	103	-50	-59	-38	>50	
	5	2006-2011	108	-34	-42	-21	>25	
CBC/BBS England	44	1967-2011	57	-58	-75	-34	>50	
	25	1986-2011	81	-59	-70	-46	>50	
	10	2001-2011	101	-48	-59	-35	>25	
	5	2006-2011	106	-34	-42	-23	>25	
BBS UK	16	1995-2011	99	-44	-54	-32	>25	
	10	2001-2011	102	-49	-59	-38	>25	
	5	2006-2011	106	-32	-41	-21	>25	
BBS England	16	1995-2011	96	-42	-52	-29	>25	
	10	2001-2011	99	-47	-58	-35	>25	
	5	2006-2011	103	-32	-42	-22	>25	


Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.





Demographic trends


Fledglings per breeding attempt Little Owl 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012


Little Owl




More on demographic trends


Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	18	None					
Clutch size	43	1968-2011	22	Linear increase	3.38 eggs	3.65 eggs	8.1%		Small sample
Brood size	43	1968-2011	47	Linear increase	2.52 chicks	2.84 chicks	12.8%		
Nest failure rate at egg stage	43	1968-2011	18	None					Small sample
Nest failure rate at chick stage	43	1968-2011	22	None					Small sample

For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).







Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of adult birds surviving to following year - error bars represent 95% confidence limits

Causes of change

There is little evidence available from the UK but studies from Europe suggest that the main demographic driver of declines in Little Owl is falling rates of juvenile survival. Circumstantial evidence suggests that this may be occurring due to loss of habitat and changes in farming practices.

Change factor	Primary driver	Secondary driver
Demographic	Decreased juvenile survival	
Ecological	Agricultural intensification	

Further information on causes of change

No trends are evident in the number of fledglings per breeding attempt, but this is based on a very small sample as few nest records are available. Clutch and brood sizes have shown linear increase, but no trends were apparent in nest failure rates (see above). There is very little evidence available from the UK regarding causes of the population decline. However, evidence from mainland Europe suggests that population changes are driven mainly by changes in survival. Le Gouar et al. (2011) analysed 35 years of ringing data from the Netherlands and found that juvenile survival rates decreased with time and that years when the population declined were associated with low juvenile survival. More than 60% of the variation in juvenile survival was explained by the increase in road traffic intensity or in average spring temperature. However, they state that these correlations reflect a gradual decrease in juvenile survival coinciding with long-term global change, rather than direct causal effects. The regular occurrence of years with poor adult survival (dry, cold years) was also important. In north-eastern France, Letty et al. (2001) also found that population dynamics were highly sensitive to adult and first-year survival and, in Switzerland and Southern Germany, Schaub et al. (2006) reported that variation of adult survival contributed most to

variation of population growth rate while variation in fecundity contributed least. Thus, evidence from Europe at least suggests that changes in populations of Little Owl are largely due to changes outside of the breeding season (although note that survival can also be affected by breeding-season conditions).

However, in Denmark, Thorup et al. (2010) found, in a declining population, that first-year annual survival rates were much lower than values previously reported, but also that the mean number of fledglings per pair had declined. Measures of reproductive success were higher closer to important foraging habitats and were positively correlated with the amount of seasonally changing land cover (mostly farmland) around nests, as well as temperatures before and during the breeding season. Experimental food supplementation to breeding pairs increased the proportion of eggs that produced fledged chicks, suggesting that the main reason for the ongoing population decline is reduced productivity induced by energetic constraints after egg-laying.

In terms of ecological drivers, in Poland, there is anecdotal evidence that changes in the agricultural landscape associated with disappearance of traditional farming and management of grassland habitats were the main factors in the long-term population decline (Salek & Schropfer 2008). Evidence from Spain has also suggested that habitat loss has played a role in population declines, due to increasing urbanisation (Martinez & Zuberogitia 2004) and in Denmark the extent of contraction of Little Owl distribution varied across the country and local disappearance was associated with reduced areas of agricultural land (Thorup et al. 2010). In Poland, Zmihorski et al. (2006) concluded that the reduction in nesting sites and decreased food availability were the potential factors of the Little Owl decline, although this evidence was circumstantial. There is little evidence relating to the UK population and the drivers in Europe may not necessarily be the same here.

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>


Key facts

Conservation listings:	Europe: no SPEC category (concentrated in Europe, conservation status favourable) (BiE04) UK: green (BoCC3)
Long-term trend:	UK: shallow decline England: fluctuating, with no long-term trend
Population size:	50,000 pairs in 2005 (APEP13: Freeman et al. 2007a)

Status summary

As a nocturnal species, Tawny Owl is covered relatively poorly by the BTO's monitoring schemes. The pattern shown by CBC/BBS is a relatively stable one, however, in keeping with the longevity, sedentary behaviour, and slow breeding rate of this species. There has been a shallow downward trend in the index since the early 1970s. It may be relevant to this possible long-term decline that Gibbons et al. (1993) found evidence for a contraction of the species' UK range between the first two atlas periods, though these losses are now largely reversed (Balmer et al. 2013a). The substantial improvements in nest success during the c.29-day egg stage could be linked to the declining impact of organochlorine pesticides, which were banned in the early 1960s. The numbers of fledglings per breeding attempt have increased steeply. Special post-breeding surveys of this species were conducted in autumn 2005 (Freeman et al. 2007a), following methodology established by an earlier survey in 1989 (Percival 1990).

CBC/BBS UK 1966–2012
Tawny Owl







Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	79	-12	-48	26		
	25	1986-2011	95	-27	-45	-9	>25	
	10	2001-2011	97	-14	-29	2		

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	44	1967-2011	67	-5	-45	41		
	25	1986-2011	81	-17	-35	8		
	10	2001-2011	85	-12	-27	6		
	5	2006-2011	92	3	-14	20		
BBS UK	16	1995-2011	90	-18	-35	1		Nocturnal species
	10	2001-2011	95	-15	-30	5		Nocturnal species
	5	2006-2011	102	-5	-23	10		Nocturnal species
BBS England	16	1995-2011	77	-10	-25	16		Nocturnal species
	10	2001-2011	83	-12	-28	4		Nocturnal species
	5	2006-2011	89	2	-18	19		Nocturnal species

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




Fledglings per breeding attempt Tawny Owl 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Tawny Owl




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	68	Linear increase	1.39 fledglings	1.97 fledglings	41.4%		
Clutch size	43	1968-2011	101	None					
Brood size	43	1968-2011	203	None					
Nest failure rate at egg stage	43	1968-2011	68	Curvilinear	1.09% nests/day	0.23% nests/day	-78.9%		
Nest failure rate at chick stage	43	1968-2011	103	Curvilinear	0.33% nests/day	0.09% nests/day	-72.7%		
Laying date	43	1968-2011	20	None			0 days		Small sample


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts

Conservation listings:	Europe: SPEC category 2 (declining) (BiE04) UK: red (>50% distribution decline) (BoCC3) UK Biodiversity Action Plan: click here , priority species
Long-term trend:	UK: uncertain
Population size:	4,600 (3,700-5,500) males in 2004 (APEP13: Conway et al. 2007)


Status summary

Following a catastrophic decline in range of more than 50% of 10-km squares between the 1968-72 and 1988-91 breeding atlases, the 1992 national survey revealed a welcome increase of 50% in population size since an earlier survey in 1981, probably due to the increased availability of young forest habitat as plantations were felled and replanted (Morris et al. 1994). A [National Nightjar Survey](#) in 2004 revealed that a further 36% increase had taken place in the UK population in 12 years, with a 2.6% increase in the number of 10-km squares occupied (Conway et al. 2007). There was evidence of population declines and range contractions since 1992, however, in North Wales, northwest England, and Scotland. Atlas data from 2008-11 show an 18% range increase in Britain since 1988-91 but some parts of the 1968-72 range remain unoccupied (Balmer et al. 2013). Although annual nest record sample are very small, the increases in nest failure rates and decreases in clutch size have resulted in the inclusion of Nightjar on the NRS concern list (Leech & Barimore 2008). A steep linear decrease is evident in the number of fledglings per breeding attempt. Nest failure is most likely in areas heavily frequented by walkers and dogs (Langston et al. 2007).

Population changes in detail


Annual breeding population changes for this species are not currently monitored by BTO

Demographic trends



Laying date 1966–2012


Nightjar




More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	21	Linear decline	1.42 fledglings	0.71 fledglings	-50.0%		
Clutch size	43	1968-2011	19	Linear decline	1.97 eggs	1.88 eggs	-4.6%		Small sample
Brood size	43	1968-2011	27	None					Small sample
Nest failure rate at egg stage	43	1968-2011	25	Linear increase	1.28% nests/day	3.79% nests/day	196.1%		Small sample
Nest failure rate at chick stage	43	1968-2011	22	Curvilinear	0.15% nests/day	0.86% nests/day	473.3%		Small sample
Laying date	43	1968-2011	21	None			0 days		Small sample


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

Key facts


Conservation listings:	Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe) (BiE04) UK: amber (25-50% decline) (BoCC3)
Long-term trend:	UK: decline
Population size:	87,000 (64,000-111,000) pairs in 2009 (APEP13: distance sampling estimate for 2006 (Newson et al. 2008) updated using BBS trend)

Status summary

Swifts were not monitored before the inception of the BBS. Their monitoring is complicated by the difficulty of finding occupied nests, by the weather-dependent and sometimes extraordinary distances from the nest at which breeding adults may forage, and by the often substantial midsummer influx of non-breeding individuals to the vicinity of breeding colonies. Since Swifts do not normally begin breeding until they are four years old, non-breeding numbers can be large. BBS results indicate that steep declines have occurred in England, Scotland and Wales since 1994. Many Swifts seen on BBS visits will not be nesting nearby, however, and the relationship between BBS transect counts and nesting numbers has not yet been investigated. The BBS Eaton et al. 2009). Modern building design and refurbishment of old buildings can unnecessarily deprive Swifts of nest sites and may be contributing to population decline: the provision of nest boxes and integration of potential nest sites into new buildings and renovations are strongly supported by Crowe 2012). There has been little long-term change across Europe since 1980 (PECBMS 2013a).

BBS UK 1994–2012

Swift







Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	16	1995-2011	1026	-39	-46	-32	>25	
	10	2001-2011	1111	-28	-35	-19	>25	
	5	2006-2011	1186	-21	-26	-16		

BBS England Source	16 Period (yrs)	1995-2011 Years	887 Plots	-39 Change (%)	-45 Lower limit	-31 Upper limit	>25 Alert >25	Comment
	5	2006-2011	1031	-19	-24	-14		
BBS Scotland	16	1995-2011	50	-57	-68	-37	>50	
	10	2001-2011	56	-26	-47	12		
	5	2006-2011	59	-38	-51	-11	>25	
BBS Wales	16	1995-2011	66	-44	-61	-11	>25	
	10	2001-2011	70	-41	-55	-13	>25	
	5	2006-2011	67	-18	-34	17		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.



Demographic trends

Productivity and survival trends for this species are not currently produced by BTO

Key facts


Conservation listings:	Europe: SPEC category 3 (depleted) (BiE04) UK: amber (European status) (BoCC3)
Long-term trend:	UK waterways: fluctuating, with no long-term trend
Population size:	3,800-6,400 pairs in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)

Status summary

The Kingfisher declined along linear waterways (its principal habitat) until the mid 1980s, since when it seems to have made a complete recovery. The decline was associated with a contraction of range in England (Gibbons et al. 1993). Kingfishers suffer severe mortality during harsh winters but, with up to three broods in a season, and up to six chicks in a brood, their potential for rapid population growth is unusually high. It is likely, therefore, that winter weather is the main driver of population change. Amber listing of this species in the UK results from its 'depleted' status in Europe as a whole, following declines between 1970 and 1990 (BirdLife International 2004).

WBS/WBBS 1974-2012

Kingfisher




Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
WBS/WBBS waterways	36	1975-2011	54	-13	-45	27		
	25	1986-2011	66	55	9	137		
	10	2001-2011	96	-12	-32	14		
	5	2006-2011	86	-14	-31	6		

BBS UK Source	16 Period (yrs)	1995-2011 Years	54 Plots (n)	-39 Change (%)	-54 Lower limit	-15 Upper limit	>25 Alert >25	Comment
	5	2006-2011	65	-43	-52	-33	>25	
BBS England	16	1995-2011	48	-33	-49	-11	>25	
	10	2001-2011	55	-33	-46	-16	>25	
	5	2006-2011	59	-42	-50	-30	>25	


Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.



WBS/WBBS waterways graph



BBS UK graph



BBS England graph

Demographic trends

Productivity and survival trends for this species are not currently produced by BTO

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts


Conservation listings:	Europe: SPEC category 2 (depleted) (BiE04) UK: amber (European status) (BoCC3)
Long-term trend:	England: rapid increase
Population size:	52,000 (47,000-58,000) pairs in 2009 (APEP13: distance sampling estimate for 2006 (Newson et al. 2008) updated using BBS trend)
Migrant status:	Resident
Nesting habitat:	Cavity nester
Primary breeding habitat:	Woodland
Secondary breeding habitat:	
Breeding diet:	Animal
Winter diet:	Animal

Status summary

Green Woodpecker populations have risen steadily in Britain since 1966, except for a period of stability or shallow decline centred around 1980. There was considerable range expansion in central and eastern Scotland between the 1968-72 and 1988-91 atlas periods. Recent atlas results indicate that expansion is continuing across England and Scotland, but not in Wales, where major retraction from some western regions was detected in 2008-11 (Balmer et al. 2013). Similarly, the BBS PECBMS 2013a).

CBC/BBS England 1966–2012

Green Woodpecker




Smoothed population index, relative to an arbitrary 100 in the year given, with 85% confidence limits in green

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	44	1967-2011	339	199	119	335		
	25	1986-2011	548	115	87	164		
	10	2001-2011	889	3	-2	8		
	5	2006-2011	1012	-4	-8	0		
BBS UK	16	1995-2011	805	38	28	51		
	10	2001-2011	941	2	-2	8		
	5	2006-2011	1059	-5	-8	-2		
BBS England	16	1995-2011	751	49	39	62		
	10	2001-2011	882	3	-2	8		
	5	2006-2011	1001	-4	-7	0		
BBS Wales	16	1995-2011	48	-19	-40	4		
	10	2001-2011	52	-11	-29	8		
	5	2006-2011	53	-17	-33	-1		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




CBC/BBS England 1966-2012
Green Woodpecker




CBC/BBS England graph

BBS UK 1994-2012
Green Woodpecker




BBS UK graph

BBS England 1994-2012
Green Woodpecker



BBS England graph

BBS Wales 1994-2012
Green Woodpecker



Demographic trends

Productivity and survival trends for this species are not currently produced by BTO

Causes of change

There is little evidence available regarding the demographic or ecological causes of population increase in this species.

Change factor	Primary driver	Secondary driver
Demographic	Unknown	
Ecological	Unknown	

Further information on causes of change

No information on demographic trends for this species is available. The ecological factors underlying the increase in population size are not yet known but, given the species' susceptibility to cold weather, it may be related to climate change. Smith (2007) found that Green Woodpeckers were not limited by nest-sites in his study woods in southern England and linked the upward trend in numbers to the availability of food outside the woods and higher survival due to a series of mild winters.

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Great Spotted Woodpecker

Dendrocopos major

Key facts

Conservation listings:	Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe) (BiE04) UK: green (species level); amber (race <i>anglicus</i> , >20% of European breeders) (BoCC3)
Long-term trend:	UK, England: rapid increase
Population size:	140,000 (130,000-150,000) pairs in 2009 (APEP13: distance sampling estimate for 2006 (Newson et al. 2008) updated using BBS trend)


Migrant status:	Resident
Nesting habitat:	Cavity nester
Primary breeding habitat:	Woodland
Secondary breeding habitat:	
Breeding diet:	Animal
Winter diet:	Animal

Status summary

This species increased rapidly in the 1970s and began a further increase in the mid 1990s. The BBS Balmer et al. 2013). Numbers have shown widespread moderate increase across Europe since 1980 (PECBMS 2013a).




CBC/BBS UK 1966–2012




Great Spotted Woodpecker



Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	463	405	265	630		
	25	1986-2011	751	149	121	185		
	10	2001-2011	1301	58	51	66		
	5	2006-2011	1535	12	8	16		
CBC/BBS England	44	1967-2011	411	360	230	600		
	25	1986-2011	663	129	102	165		
	10	2001-2011	1138	45	39	51		
	5	2006-2011	1340	6	3	10		
BBS UK	16	1995-2011	1047	139	125	155		
	10	2001-2011	1301	58	50	65		
	5	2006-2011	1535	12	8	15		
BBS England	16	1995-2011	912	117	104	130		
	10	2001-2011	1124	44	38	52		
	5	2006-2011	1317	6	3	9		
BBS Scotland	16	1995-2011	48	329	223	465		
	10	2001-2011	66	124	83	181		
	5	2006-2011	88	28	6	59		
BBS Wales	16	1995-2011	76	198	136	266		
	10	2001-2011	94	116	83	157		
	5	2006-2011	104	43	24	53		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.






Demographic trends

Fledglings per breeding attempt


Great Spotted Woodpecker 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Great Spotted Woodpecker




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Brood size	43	1968-2011	31	Curvilinear	3.70 chicks	3.82 chicks	3.2%		
Nest failure rate at chick stage	43	1968-2011	36	None					


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

Causes of change

There is good evidence that nest survival has increased, most likely due to decreased competition with Starlings. This is based on one local study but supported by more extensive analysis of nest record cards. Use of garden feeders may be another of many factors contributing to their population increase.

Change factor	Primary driver	Secondary driver
Demographic	Increased breeding success	
Ecological	Decreased competition	

Further information on causes of change

The initial increase in Great Spotted Woodpeckers during the 1970s has been attributed to Dutch elm disease, which greatly increased the amount of standing dead timber, thereby increasing associated insects and so improving food supplies and providing nest sites (Marchant et al. 1990). However, studies giving demographic evidence supporting the effects of this are sparse. There has been speculation that the storms of 1987 and 1990 also benefited Great Spotted Woodpeckers by increasing the availability of dead wood, although a detailed study by Smith (1997), in two study woodlands, reported no specific link between woodpecker increase and the storms, despite the increase in dead wood.

A long-term study of the breeding success of an increasing population of Great Spotted Woodpeckers in southern England provides good evidence that nest survival has increased dramatically over the last 20 years (Smith 2005, 2006). Nest-site interference by 2005 analysed national nest record cards and found similar trends in nest survival, supporting the hypothesis that reduced competition with Starlings has led to the increase in woodpecker population. The decline in Starling numbers in recent decades may also have allowed Great Spotted Woodpeckers to expand their breeding distribution into less-wooded habitats (Smith 2005). Great Spotted Woodpeckers appear limited in their ability to advance their breeding period to maintain synchrony with their natural prey and thus their ready use of garden feeders has the potential to increase breeding success (Smith & Smith 2013).

It is possible that recent increases of Great Spotted Woodpeckers, are also, at least in part, driven by changing climate (Fuller et al. 2005). In Scandinavia (Nilsson et al. 1992) and Białowieża Forest, Poland (Wesołowski & Tomiałojć 1986), breeding numbers were found to be related to the severity of the preceding winter and the availability of conifer seeds on which the birds then feed. No similar relationship has been found in Britain (Marchant et al. 1990), which is probably not surprising given our relatively mild winters (Smith 1997). Smith (2006) found no evidence that increasing spring temperatures impacted on clutch size, nesting success or number of young fledged.

Lesser Spotted Woodpecker


Dryobates minor

Key facts

Conservation listings:	Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe) (BiE04) UK: red (>50% population decline) (BoCC3); an RBBP species UK Biodiversity Action Plan: priority species
Long-term trend:	UK: rapid decline
Population size:	1,000-2,000 pairs in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)
Migrant status:	Resident
Nesting habitat:	Cavity nester
Primary breeding habitat:	Woodland
Secondary breeding habitat:	
Breeding diet:	Animal
Winter diet:	Animal

Status summary

The Lesser Spotted Woodpecker has declined significantly and very rapidly since around 1980, following a shallower increase; it had already contracted in range between the first two atlas periods (Gibbons et al. 1993), and has subsequently disappeared from many more of its former localities (Balmer et al. 2013). It has become so rare that BBS observers have been unable to continue the annual monitoring that was possible until 2000 through CBC. The species qualifies easily for red listing. All UK breeding records since 2010 should be forwarded to the Rare Breeding Birds Panel, who have established PECBMS 2007, 2013a).



Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC all habitats	31	1968-1999	17	-60	-81	40		Small CBC sample
	25	1974-1999	18	-73	-86	-31	>50	Small CBC sample
	10	1989-1999	11	-51	-75	-22	>50	Small CBC sample
	5	1994-1999	9	-33	-56	0		Small sample

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See here for more information.



CBC all habitats graph

Demographic trends

Productivity and survival trends for this species are not currently produced by BTO

Causes of change

The demographic causes of decline are not yet known and, although there is low breeding success in some populations, the reasons for the decline are unclear.

Change factor	Primary driver	Secondary driver
Demographic	Unknown	
Ecological	Unknown	

Further information on causes of change

The demographic causes of decline are not yet known, and although there is low breeding success in some populations the reasons for the decline in the UK and elsewhere in Europe are unclear (Charman et al. 2009). A detailed field study in Sweden provided good evidence that neither clutch size, brood size in successful nests, fledging success in successful nests nor mean nestling weight differed significantly between years, despite a threefold difference in population variation (Wiktander et al. 2001).

Loss of open woodland is one factor that has been suggested to have contributed to declines in this species. Lesser Spotted Woodpecker is a species that requires mature, open woodland and large areas of woodland at a landscape scale (Wiktander et al. 2001, Charman et al. 2010). Wiktander et al. postulate that the decrease in the area of deciduous forest in Sweden is probably one cause of this species' decline, although they present no specific evidence to support this (Wiktander et al. 1992). Loss of dead wood within woodlands has been proposed as another factor; however, given that dead wood has increased in Britain (Amar et al. 2010) this seems an unlikely cause here. A field study in Poland provided evidence that Lesser Spotted Woodpecker presence is closely correlated with the amount of dead wood and large deciduous trees (Angelstam et al. 2002). In their review of the causes of declines of woodland birds Fuller et al. (2005) state that reductions in small-diameter dead wood suitable for foraging may be a factor in the decline, although recent surveys provided evidence that there was no difference in dead-wood abundance between occupied and unoccupied woods (Charman et al. 2010). However, dead snags have a high turnover and were found to be suitable for nesting sites by woodpeckers for only a few years after death and, furthermore, dead-wood conditions may now be more favourable for Smith 2007).

A third hypothesis relates to competition and predation. A field study in Sweden found that Great Spotted Woodpeckers compete with Lesser Spotted for insect food in dead wood when spruce seed crops are low (Nilsson et al. 1992), but evidence for this in Britain is limited (Charman et al. 2010). The two species may compete for nest sites, since they overlap considerably in their use of nesting substrates (Glue & Boswell 1994). Amar et al. (2006) found that Lesser Spotted Woodpecker decreased more heavily in woods with relatively high numbers of grey squirrel dreys but there was no other evidence that squirrel density was a significant factor in declines.

Changing climate has been found to have an impact on survival and reproduction in some populations. In Norway, a positive relationship between spring numbers of Lesser Spotted Woodpecker and previous June temperatures has been interpreted as an effect of temperatures on woodpecker survival and reproduction during the breeding season (Steen et al. 2006, Selas et al. 2008). Steen et al. (2006) also found that winter temperatures exhibit a direct positive effect on winter survival. However, given that there has been a general trend for increasing temperatures in the UK (see [here](#)), it seems unlikely that changes in climate have been responsible for Lesser Spotted Woodpecker declines. Work in Sweden and Germany suggests that changes in phenology could play a role in breeding success, finding that declines in food availability during the breeding season are likely to be related to seasonal declines in reproductive performance as woodpeckers adjust their timing of breeding to coincide with the seasonal food peak (Wiktander et al. 2001, Rossmann et al. 2007). However, there is little further evidence for this. In Britain, breeding success has fallen and is lower than in recent studies in Germany and Sweden; chick mortality is especially high, most probably related to food shortages in the breeding period (Charman et al. 2012, Smith & Charman 2012).

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglinton, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Riebel, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts


Conservation listings:	Europe: SPEC category 3 (declining) (BiE04) UK: amber (25-50% population decline) (BoCC3)
Long-term trend:	England: fluctuating, with no long-term trend
Population size:	46,000 pairs in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend for England)

Status summary

Kestrels had recovered from the lethal and sublethal effects of organochlorine pesticides by the mid 1970s, the recovery probably driven by improving nesting success, but subsequently entered a decline which has been linked to the effects of agricultural intensification on farmland habitats and their populations of small mammals (Gibbons et al. 1993). Since the mid 1980s, the English population has fluctuated without a long-term trend being apparent but there are significant declines over the BBS period in England and especially in Scotland. The BBS Leech & Barimore 2008). Despite its decline since the mid 1970s, the Kestrel breeds at high density in mixed farmland across much of England, suggesting that the British population might have exceeded 50,000 pairs (Clements 2008). A moderate decrease has been recorded in the Republic of Ireland since 1998 (Crowe 2012). There has been widespread moderate decrease across Europe since 1980 (PECBMS 2013a).

CBC/BBS England 1966-2012

Kestrel



Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	44	1967-2011	276	-8	-37	28		
	25	1986-2011	426	-4	-19	10		
	10	2001-2011	657	-10	-16	-4		
	5	2006-2011	739	-15	-20	-11		

BBS UK Source	Period (yrs)	1995-2011 Years	BBS (n)	Change (%)	Lower limit	Upper limit	>25 Alert	Comment
	10	2001-2011	720	-20	-27	-12		
	5	2006-2011	786	-23	-29	-16		
BBS England	16	1995-2011	568	-15	-21	-8		
	10	2001-2011	636	-10	-15	-4		
	5	2006-2011	704	-16	-20	-11		
BBS Scotland	16	1995-2011	43	-57	-72	-35	>50	
	10	2001-2011	44	-40	-62	-12	>25	
	5	2006-2011	44	-40	-60	-17	>25	

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




CBC/BBS England 1966–2012
Kestrel




CBC/BBS England graph

BBS UK 1994–2012
Kestrel




BBS UK graph

BBS England 1994–2012
Kestrel



BBS England graph


BBS Scotland 1994–2012
Kestrel



BBS Scotland graph

Fledglings per breeding attempt


Kestrel 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Kestrel




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	41	Curvilinear	2.92 fledglings	3.60 fledglings	23.2%		
Clutch size	43	1968-2011	59	None					
Brood size	43	1968-2011	148	None					
Nest failure rate at egg stage	43	1968-2011	41	Curvilinear	0.74% nests/day	0.12% nests/day	-83.8%		
Nest failure rate at chick stage	43	1968-2011	70	Linear decline	0.21% nests/day	0.08% nests/day	-61.9%		
Laying date	43	1968-2011	23	Linear decline	May 5	Apr 26	-9 days		Small sample


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

Key facts

Conservation listings:	Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe) (BiE04) UK: amber (historical decline) (BoCC3); an RBBP species
Long-term trend:	UK: probable increase
Population size:	1,200 (900-1,500) pairs in 2008 (APEP13: Ewing et al. 2011)


Status summary

Having declined substantially over the past two centuries, Merlin shows indications of a recent doubling of UK population (Rebecca & Bainbridge 1998). This increase may be associated with an increased use of forest edge as a nesting habitat (Parr 1994, Little et al. 1995, Rebecca 2011). Because of its recent population upturn, the species was moved from the red to the amber list in 2002. It remains much too scarce, however, for annual population monitoring via BBS: dedicated observers and specialised field methods are required, as described by Hardey et al. (2009). Submissions to the Rare Breeding Birds Panel fall well short of the estimated UK total population but show an average of 1.86 young fledged per occupied territory during 1996-2004 (Holling & RBBP 2007a). Breeding performance has tended to improve since the 1960s, probably linked to the declining influence of organochlorine pesticides (Crick 1993, Newton 2013). Hatching rates in the southeast Yorkshire Dales were consistently higher than had been recorded in earlier studies in Northumberland (Wright 2005). A repeat survey of Merlin's British breeding status undertaken in 2008 found a non-significant decline of around 13% since the previous survey in 1993-94, with decline most noticeable in northern England (Ewing et al. 2011).

Population changes in detail

Annual breeding population changes for this species are not currently monitored by BTO


Demographic trends



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Merlin




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	21	Linear increase	2.44 fledglings	3.38 fledglings	38.4%		
Clutch size	43	1968-2011	36	None					
Brood size	43	1968-2011	55	Linear increase	3.51 chicks	3.83 chicks	9.1%		
Nest failure rate at egg stage	43	1968-2011	23	Linear decline	0.72% nests/day	0.17% nests/day	-76.4%		Small sample
Nest failure rate at chick stage	43	1968-2011	28	Linear decline	0.90% nests/day	0.22% nests/day	-75.6%		Small sample


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts


Conservation listings:	Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe) (BiE04) UK: green (BoCC3); an RBBP species
Long-term trend:	UK, England: increase
Population size:	2,800 pairs in 2009 (APEP13: 2000 estimate (Clements 2001) updated using BBS trend)

Status summary

This species used to be too rare and unobtrusive for wide-scale monitoring but, following population increase, BBS is now able to produce a trend. Many BBS sightings must, however, refer to migrants, first-summer non-breeders, or to breeding birds from distant nests. To establish whether nesting occurs in a locality, dedicated observers and specialised field methods are required, as described by Hardey et al. (2009). The Rare Breeding Birds Panel collects annual data on nesting pairs, which under-represent the true population to unknown degrees, but adequately establish the long-term upward trend (eg Holling & RBBP 2012). Numbers in parts of southeast England could be considerably higher than previously recognised (Clements & Everett 2011). The Hobby's distribution has spread markedly northwards in England since the 1970s (Gibbons et al. 1993), perhaps linked to increases in its dragonfly prey supplies (Prince & Clarke 1993) and to a decreasing dependency on its traditional heathland habitat, but the reasons underlying the increase are still only speculative (Clements 2001). The species is now widespread north to Lancashire and Co Durham (Balmer et al. 2013). A success rate of more than 90% was recorded for nests in Derbyshire during 1992-2001, with successful nests fledging a mean of 2.44 young (Messenger & Roome 2007). The small annual samples of nest record cards indicate no long-term change in either brood size or nest success.

BBS UK 1994–2012

Hobby





Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	16	1995-2011	42	12	-20	51		
	10	2001-2011	49	18	-9	51		

Source	Period (yrs)	2006-2011 Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS England	16	1995-2011	40	19	-8	62		
	10	2001-2011	48	29	2	72		
	5	2006-2011	55	24	2	54		


Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




Demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Brood size	43	1968-2011	24	None					Small sample
Nest failure rate at chick stage	43	1968-2011	14	None					Small sample


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).



Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>


Key facts

Conservation listings:	Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe) (BiE04) UK: green (species level); amber (race <i>peregrinus</i> , >20% of European breeders, European status) (BoCC3); an RBBP species
Long-term trend:	UK, England: increase Northern Ireland, northwest Scotland, North Wales: decline since 1991
Population size:	1,500 pairs in 2002 (APEP13: Banks et al. 2010)

Status summary

The UK population size, distribution and breeding performance have all largely recovered from the detrimental effects of organochlorine pesticides in the 1950s and 1960s (Newton 2013). Populations and breeding performance have declined recently, however, in northwest Scotland and the Northern Isles (Crick & Ratcliffe 1995). Nest record information for the UK as a whole shows a significant rise in the number of fledglings per breeding attempt. In northern England, breeding productivity on grouse moors has been 50% lower than at nests in other habitats, indicating that illegal persecution on land managed for Amar et al. 2012). The number of UK breeding pairs has been censused every decade since 1961 by BTO/JNCC/RSPB/Raptor Study Groups, and has been estimated as follows: 1961 - 385 pairs; 1971 - 489 pairs; 1981 - 728 pairs; 1991 - 1,283 pairs (Ratcliffe 1993). The Banks et al. 2003, 2010); around 50 pairs were missed in Wales, however (Dixon et al. 2008). Similar increases across Europe have resulted in a downgrading of conservation listing from 'SPEC 3 (rare)' to 'secure' (BirdLife International 2004), and consequently the species has recently been moved from the amber to the green list in the UK. BTO is organising a new national [Peregrine Survey in spring 2014](#).

BBS UK 1994–2012 Peregrine




Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	16	1995-2011	45	-28	-47	3		
	10	2001-2011	55	-7	-28	23		

Source	Period (yrs)	2006-2011 Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
--------	--------------	-----------------	-----------	------------	-------------	-------------	-------	---------


Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.



BBS UK graph

Demographic trends


Fledglings per breeding attempt Peregrine 1966-2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Peregrine




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	22	Linear increase	1.77 fledglings	2.27 fledglings	28.4%		
Clutch size	43	1968-2011	18	Curvilinear	3.89 eggs	3.27 eggs	-15.9%		Small sample
Brood size	43	1968-2011	46	Linear increase	2.35 chicks	2.60 chicks	10.6%		
Nest failure rate at egg stage	43	1968-2011	23	Curvilinear	0.15% nests/day	0.13% nests/day	-13.3%		Small sample
Nest failure rate at chick stage	43	1968-2011	26	None					Small sample
Laying date	43	1968-2011	10	Linear decline	Apr 14	Apr 3	-11 days		Small sample


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

Key facts


Conservation listings:	Europe: not evaluated (introduced) (BIE04) UK: not listed (introduced)
Long-term trend:	England: rapid increase
Population size:	8,600 pairs in 2012 (APEP13: H. Peck pers. comm., Project Parakeet)

Status summary

Following escapes and releases over many decades, this African and Asian parrot began breeding annually in the UK in 1969. Substantial but highly localised self-sustaining populations have since built up, with the two largest being in Greater London and in the Isle of Thanet, east Kent. Population modelling has revealed that populations in Greater London have increased by approximately 30% per year, and those in Thanet by 15% per year, but that the range has expanded by only 0.4 km per year in the Greater London area and hardly at all in Thanet (Butler 2003). National BBS data indicate a tenfold increase since 1995. There have been recent post-breeding estimates of more than 30,000 birds at large in the UK (Holling & RBBP 2011a). From 108 nests located during 2001-03, the mean first-egg date was 26 March, median clutch size was 4, and overall nest success 72%, making productivity sufficient to account for the observed population rise, assuming mortality rates remained low (Butler et al. 2013). The species has already been reported causing economic damage to crops, as has occurred elsewhere in its native and introduced range (Butler 2003). A recent study in Belgium has identified negative effects on breeding Strubbe & Matthysen 2007, 2009, Strubbe et al. 2010). No such effects have yet been detectable in Britain, however (Newson et al. 2011).

BBS UK 1994–2012

Ring-necked Parakeet





Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	16	1995-2011	59	1057	458	3284		
	10	2001-2011	88	252	108	494		

Source	Period (yrs)	2006-2011 Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS England	16	1995-2011	59	1058	451	3122		
	10	2001-2011	88	252	107	516		
	5	2006-2011	110	62	41	84		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.



Demographic trends

Productivity and survival trends for this species are not currently produced by BTO


This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts

Conservation listings:	Europe: no SPEC category (not concentrated in Europe, conservation status favourable) (BiE04) UK: green (BoCC3)
Long-term trend:	UK, England: rapid increase
Population size:	600,000 territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)
Migrant status:	Resident
Nesting habitat:	Above-ground nester
Primary breeding habitat:	Woodland
Secondary breeding habitat:	Human habitats
Breeding diet:	Animal
Winter diet:	Vegetation

Status summary

Magpies increased steadily until the late 1980s, after which abundance stabilised (Gregory & Marchant 1996). The BBS PECBMS 2013a).







Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
--------	--------------	-------	-----------	------------	-------------	-------------	-------	---------


CBC/BBS UK Source	Period (yrs)	1967-2011 Years	825 Plots (n)	+03 Change (%)	68 Lower limit	148 Upper limit	Alert	Comment
	25	1986-2011	1337	0	-9	8		
	10	2001-2011	2118	-5	-8	-3		
	5	2006-2011	2356	1	-1	4		
CBC/BBS England	44	1967-2011	699	107	67	173		
	25	1986-2011	1129	1	-8	9		
	10	2001-2011	1773	-4	-7	-1		
	5	2006-2011	1997	1	-2	3		
BBS UK	16	1995-2011	1868	-1	-5	3		
	10	2001-2011	2102	-6	-9	-3		
	5	2006-2011	2330	1	-1	3		
BBS England	16	1995-2011	1566	-1	-6	3		
	10	2001-2011	1760	-4	-7	-1		
	5	2006-2011	1975	1	-2	3		
BBS Scotland	16	1995-2011	46	24	-3	72		
	10	2001-2011	54	4	-17	34		
	5	2006-2011	62	9	-9	31		
BBS Wales	16	1995-2011	160	-10	-20	1		
	10	2001-2011	176	-7	-16	2		
	5	2006-2011	177	5	-2	13		
BBS N.Ireland	16	1995-2011	83	19	-8	43		
	10	2001-2011	97	-15	-24	-6		
	5	2006-2011	101	-4	-13	8		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.







BBS England graph



BBS Scotland graph




BBS Wales graph



BBS N. Ireland graph


Fledglings per breeding attempt Magpie 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Magpie




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	43	Curvilinear	1.14 fledglings	2.36 fledglings	107.1%		
Clutch size	43	1968-2011	43	Linear decline	5.77 eggs	4.83 eggs	-16.3%		
Brood size	43	1968-2011	83	Curvilinear	3.23 chicks	2.57 chicks	-20.6%		
Nest failure rate at egg stage	43	1968-2011	49	Linear decline	2.74% nests/day	0.20% nests/day	-92.7%		
Nest failure rate at chick stage	43	1968-2011	49	Linear decline	1.67% nests/day	0.10% nests/day	-94.0%		
Laying date	43	1968-2011	33	Linear decline	Apr 24	Mar 24	-31 days		


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

Causes of change

The number of fledglings per breeding attempt increased strongly until the 1990s but then stabilised, a pattern mirroring the population index, which suggests that changing breeding success has been an important driver of population change. There is little published evidence about the ecological drivers of change. Changes in control of Magpies could have played a role, but their generalist ecology means that they are able to prosper in suburban and intensively farmed landscapes, which is likely to have allowed populations to reach a historically high equilibrium level.

Change factor	Primary driver	Secondary driver
Demographic	Change in breeding success	
Ecological	Unknown	

Further information on causes of change

Although there is little evidence directly supporting this, it is likely that the stabilisation in Magpie numbers reflects the population reaching carrying capacity in the intensively farmed and modern suburban landscapes. The fact that recent stability or decline is associated with parallel trends in fledglings per breeding attempt supports this. Demographic data presented here show that the number of fledglings per breeding attempt increased dramatically up until the 1990s but then stabilised (see above). Although clutch and brood sizes have declined over the whole time series (1968-2009), there have also been decreases in the failure of nests at the egg and chick stages (see above). A strong trend towards earlier laying has also been identified and may be partly explained by recent climate change (Crick & Sparks 1999).

The historical increases in Magpies have occurred at the same time as falling levels of control by gamekeepers from the time of the First World War (Tapper 1992), but there is no direct evidence for a causal link. Since 1990, the widespread adoption of the Larsen trap for predator control has been responsible for a large increase in Magpie numbers killed on shooting estates ([GWCT data](#)), and this could have played a role in stabilising population growth in some areas, but is unlikely to explain population change in towns and cities.


Magpies have increased in farmland and woodland habitats, with the largest population growth on mixed and pastoral farms, and the smallest on arable land (Gregory & Marchant 1996). The remarkable adaptability of Magpies has enabled them to colonise many new urban and suburban localities since the 1960s.

Key facts

Conservation listings:	Europe: no SPEC category (not concentrated in Europe, conservation status favourable) (BiE04) UK: green (species level); amber (races hibernicus and rufitergum, >20% of European breeders) (BoCC3)
Long-term trend:	UK, England: fluctuating, with no long-term trend
Population size:	170,000 territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)

Status summary

The UK Jay population remained stable in the species' preferred woodland habitat until the late 1980s, after which the population began to decline. This decrease followed an earlier decline on farmland CBC plots (Gregory & Marchant 1996). With the losses since the 1980s now regained, long-term trends are stable overall. The BBS PECBMS 2013a).








Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	367	6	-13	34		Small CBC sample
	25	1986-2011	563	-3	-13	10		Small CBC sample
	10	2001-2011	885	11	4	18		
	5	2006-2011	1002	2	-5	8		
CBC/BBS England	44	1967-2011	324	-1	-19	26		Small CBC sample
	25	1986-2011	493	-9	-18	2		Small CBC sample

Source	Period (yrs)	2001-2011 Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
	5	2006-2011	875	4	-1	9		
BBS UK	16	1995-2011	749	15	7	23		
	10	2001-2011	885	11	5	19		
	5	2006-2011	1002	2	-5	9		
BBS England	16	1995-2011	646	7	0	16		
	10	2001-2011	760	12	6	20		
	5	2006-2011	865	4	-1	9		
BBS Wales	16	1995-2011	71	38	10	65		
	10	2001-2011	82	13	0	33		
	5	2006-2011	85	3	-8	17		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.






Demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Brood size	43	1968-2011	11	Linear increase	3.41 chicks	3.93 chicks	15.3%		Small sample

For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts


Conservation listings:	Europe: no SPEC category (concentrated in Europe, conservation status favourable) (BiE04) UK: green (BoCC3)
Long-term trend:	UK, England: rapid increase
Population size:	1.4 (1.2-1.5) million pairs in 2009 (APEP13: distance sampling estimate for 2006 (Newson et al. 2008) updated using BBS trend)

Migrant status:	Resident
Nesting habitat:	Cavity nester
Primary breeding habitat:	Farmland
Secondary breeding habitat:	
Breeding diet:	Animal
Winter diet:	Animal

Status summary



Jackdaws have increased in abundance since the 1960s (Gregory & Marchant 1996), and more recent BBS data suggest that the increase is continuing in all UK countries. The BBS PECBMS 2013a).






CBC/BBS UK 1966–2012 Jackdaw




Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	690	123	38	235		
	25	1986-2011	1163	56	19	90		
	10	2001-2011	1952	27	21	33		
	5	2006-2011	2219	19	13	24		
CBC/BBS England	44	1967-2011	554	108	36	262		
	25	1986-2011	932	49	21	88		
	10	2001-2011	1569	30	23	37		
	5	2006-2011	1807	19	15	25		
BBS UK	16	1995-2011	1665	51	40	65		
	10	2001-2011	1918	29	22	35		
	5	2006-2011	2161	20	15	25		
BBS England	16	1995-2011	1333	53	44	62		
	10	2001-2011	1542	31	24	38		
	5	2006-2011	1763	19	14	23		
BBS Scotland	16	1995-2011	113	39	14	79		
	10	2001-2011	126	26	3	60		
	5	2006-2011	142	34	12	56		
BBS Wales	16	1995-2011	138	35	-4	96		
	10	2001-2011	156	10	-9	25		
	5	2006-2011	157	2	-20	20		
BBS N.Ireland	16	1995-2011	75	107	51	148		
	10	2001-2011	90	42	21	66		
	5	2006-2011	96	39	18	56		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.






Fledglings per breeding attempt Jackdaw 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Jackdaw




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	61	Linear increase	1.73 fledglings	2.53 fledglings	46.5%		
Clutch size	43	1968-2011	54	None					
Brood size	43	1968-2011	132	None					
Nest failure rate at egg stage	43	1968-2011	69	Linear decline	0.70% nests/day	0.17% nests/day	-75.7%		
Nest failure rate at chick stage	43	1968-2011	66	Linear decline	1.10% nests/day	0.22% nests/day	-80.0%		
Laying date	43	1968-2011	30	Curvilinear	Apr 24	Apr 16	-8 days		


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

Causes of change

There is no evidence available regarding the ecological causes of increase for this species but changes have been associated with improvements in breeding performance, probably due to increased food availability.

Change factor	Primary driver	Secondary driver
Demographic	Increased breeding success	
Ecological	Unknown	

Further information on causes of change

As with [Magpie](#), [Rook](#) and [Carrion Crow](#), the increase has been associated with improvements in breeding performance and probably reflects the species' generalist feeding habits, which allow it to exploit diverse and ephemeral food resources, although direct evidence for this is limited. There have been substantial declines in nest failure rates during the egg and chick stages, and the number of fledglings per breeding attempt has improved steadily. Laying dates have advanced by more than a week.

Typically in this species, the younger chicks of a brood perish quickly if food becomes limited. Henderson & Hart (1993) provided evidence that increases in fledging success are likely to be due to improved provisioning by the parents. Most of the variation in annual reproductive output was caused by nestling mortality rather than clutch size or hatching success. Soler & Soler (1996) used data from Spain to show that additional food advanced the laying date, increased the clutch size, independently of laying date, and increased fledging success.


Changes in the landscape may have also benefited this species. Gregory & Marchant (1996) found an increase in Jackdaw numbers in agricultural habitats, particularly in the south-west, but an overall decrease in forests. These increases were associated with trends in cultivation and population gains have been most pronounced on grazing farms and in the north and south-west where such farms predominate. A similar pattern was found in Sweden by Andren (1992), who provided evidence that the density of Jackdaws increased as forest became fragmented and intermixed with agricultural land.

Key facts

Conservation listings:	Europe: no SPEC category (not concentrated in Europe, conservation status favourable) (BiE04) UK: green (BoCC3)
Long-term trend:	UK: increase
Population size:	1.1 (1.0-1.2) million pairs in 2009 (APEP13: 1996 estimate (Marchant & Gregory 1999) updated using BBS trend)

Status summary

Relatively few rookeries fell within CBC plots, but an index calculated from the available CBC nest counts showed a shallow, long-term increase (Wilson et al. 1998). The trend is confirmed by the results of the most recent BTO rookeries survey, which identified a 40% increase in abundance between 1975 and 1996 (Marchant & Gregory 1999). This increase probably reflects the species' considerable adaptability in the face of agricultural change. BBS indices, which are drawn from sightings during transect walks and not from BBS's nest counts, suggest that some decrease has occurred subsequently, especially in Scotland and Northern Ireland since around 2000. The BBS PECBMS 2013a).




Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	16	1995-2011	1281	-17	-24	-9		
	10	2001-2011	1415	-21	-27	-13		
	5	2006-2011	1571	-11	-17	-4		
BBS England	16	1995-2011	1017	-10	-19	-2		

Source	10 Period (yrs)	2001-2011 Years 2006-2011	1126 Plots (2009)	-9 Change (%)	-17 Lower limit	-2 Upper limit	Alert	Comment
BBS Scotland	16	1995-2011	109	-34	-48	-11	>25	
	10	2001-2011	114	-37	-51	-19	>25	
	5	2006-2011	125	-20	-34	-1		
BBS Wales	16	1995-2011	78	-26	-46	4		
	10	2001-2011	85	-17	-40	12		
	5	2006-2011	86	-6	-23	16		
BBS N.Ireland	16	1995-2011	74	-1	-29	36		
	10	2001-2011	88	-40	-51	-24	>25	
	5	2006-2011	91	-8	-28	15		


Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




BBS UK graph




BBS England graph



BBS Scotland graph



BBS Wales graph




BBS N.Ireland graph

Demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Clutch size	43	1968-2011	11	None					Small sample
Brood size	43	1968-2011	79	Curvilinear	2.22 chicks	2.02 chicks	-9.0%		
Nest failure rate at egg stage	43	1968-2011	29	None					Small sample
Nest failure rate at chick stage	43	1968-2011	46	None					
Laying date	43	1968-2011	13	None			0 days		Small sample

For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend


This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts

Conservation listings:	Europe (C. corone/cornix): no SPEC category (not concentrated in Europe, conservation status favourable) (BiE04) UK (C. corone/cornix): green (BoCC3)
Long-term trend:	England: rapid increase
Population size:	1.0 million territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend for England)
Migrant status:	Resident
Nesting habitat:	Above-ground nester
Primary breeding habitat:	?
Secondary breeding habitat:	
Breeding diet:	Animal
Winter diet:	Animal


Status summary

Carrion Crows increased steadily since the 1960s (Gregory & Marchant 1996) and reached a plateau around the turn of the century. The BBS [map](#) of change in relative density between 1994-96 and 2007-09 indicates that, despite strong increase in southeastern England and much of Scotland, there have been sharp decreases in south-western England, Wales, upland England and northeastern Scotland. Since 1968, breeding productivity has increased steeply and laying date has advanced by ten days.




Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	16	1995-2011	2301	13	6	21		
	10	2001-2011	2602	-3	-8	3		
	5	2006-2011	2908	-1	-5	3		
BBS England	16	1995-2011	1901	20	12	30		
	10	2001-2011	2158	5	0	11		
	5	2006-2011	2435	2	-2	6		
BBS Scotland	16	1995-2011	185	-8	-24	10		
	10	2001-2011	202	-19	-34	-4		
	5	2006-2011	228	-9	-21	3		
BBS Wales	16	1995-2011	201	7	-8	20		
	10	2001-2011	227	-9	-20	4		
	5	2006-2011	228	-6	-20	7		


Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




BBS UK graph



BBS England graph




BBS Scotland graph



BBS Wales graph


Fledglings per breeding attempt Carrion Crow 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Carrión Crow




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	40	Curvilinear	1.73 fledglings	2.35 fledglings	36.4%		
Clutch size	43	1968-2011	31	Curvilinear	4.05 eggs	4.17 eggs	3.1%		
Brood size	43	1968-2011	80	Curvilinear	2.91 chicks	2.49 chicks	-14.6%		
Nest failure rate at egg stage	43	1968-2011	48	Linear decline	1.65% nests/day	0.15% nests/day	-90.9%		
Nest failure rate at chick stage	43	1968-2011	42	Linear decline	0.78% nests/day	0.09% nests/day	-88.5%		
Laying date	43	1968-2011	30	Linear decline	Apr 18	Apr 8	-10 days		Small sample


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

Causes of change

There are few specific studies providing evidence for the causes of the increase in this species, although evidence presented here shows that increases in breeding success have been important. Ecological causes of this could be increases in food availability and the increasing suitability of urban areas (driving the species' expansion there), although specific evidence supporting these hypotheses is limited.

Change factor	Primary driver	Secondary driver
Demographic	Increased breeding success	
Ecological	Other	

Further information on causes of change

The demographic trends shown here reveal that there has been a strong increase in the number of fledglings produced per breeding attempt between 1968 and 2008, reflecting a linear decline in daily failure rate of nests at the egg and chick stages. Clutch size has increased, but brood size has decreased. This suggests that the increase in Carrion Crow numbers is related to increases in breeding success although, as there are no estimates of survival, it is not possible to say what part this has played.

This species is omnivorous and highly adaptable and is thus able to exploit changing habitats and the ephemeral food resources in intensive agriculture, from ploughed fields to grazed pasture, allowing breeding pairs to hold territories year-round. It is also able to exploit the varied food sources found in towns and cities. Richner (1992) provided good evidence that food-supplemented pairs had a higher nesting success and produced more and heavier fledglings, demonstrating that food limitation can cause low fitness for individuals and thus could potentially restrict population-level reproductive success. In a local study, Yom-Tov (1974) showed that provision of excess food improved chick survival, and concluded that the distribution pattern of food was the ultimate factor limiting breeding success, perhaps because this affects levels of intraspecific nest predation. Although the impact on population size was not considered in these studies, it is possible that food availability for Carrion Crows has increased and so helped to support the population increase. O'Connor & Shrubbs (1986) suggest that the general increase in density of sheep in upland areas, and the increase in carrion resulting from this, may be responsible for the expansion of Carrion Crow populations, although evidence for this was not given and this is clearly not relevant to lowland areas (where sheep numbers have decreased).

A second hypothesis to explain this species' increase is that control by gamekeepers has reduced, but evidence supporting this is limited. Tharme et al. (2001) stated that the control of Carrion Crows by gamekeepers was the most probable cause of the low densities on grouse moors, although they found no significant relationship between the number of gamekeepers and Carrion Crow density. Furthermore, bag returns have shown no overall change in the number of Carrion Crows killed since 1961 (Tapper 1992, Tapper & France 1992).

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglinton, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>


Key facts

Conservation listings:	Europe (<i>C. corone/cornix</i>): no SPEC category (not concentrated in Europe, conservation status favourable) (BiE04) UK: green (BoCC3)
Long-term trend:	UK: uncertain
Population size:	260,000 territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)

Status summary

The BOU Records Committee took the decision in 2002 to treat Hooded Crow and Parkin et al. 2003). This split is not yet recognised in European conservation listings. In the UK, Hooded Crows occur in Northern Ireland, the Isle of Man, and in Scotland, mainly west and north of the Great Glen. Retrospective analysis of BBS trends is simple because observers have always recorded Hooded Crows (coded HC) separately from Carrion Crows and from intermediates (coded HB). Intermediate forms between Carrion and Hooded, which predominate in a band across western Scotland and occur less frequently elsewhere in the UK, are not included in either BBS index. BBS data suggest that some decrease in Hooded Crows may have occurred in Scotland, but that this has been countered by increase in Northern Ireland. Hooded Crows have increased markedly in Ireland since 1924 (Hutchinson 1989).

BBS UK 1994–2012
Hooded Crow




Smoothed population index, relative to an arbitrary 100 in the year given, with 85% confidence limits in green

Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	16	1995-2011	134	6	-18	36		
	10	2001-2011	147	10	-12	35		
	5	2006-2011	151	10	-9	34		
BBS Scotland	16	1995-2011	50	-23	-47	10		

Source	Period (yrs)	2001-2011 Years	#Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
	5	2006-2011	49	5	-18	41		
BBS N.Ireland	16	1995-2011	81	127	77	189		
	10	2001-2011	97	3	-8	18		
	5	2006-2011	100	15	1	31		


Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.



BBS UK graph



BBS Scotland graph



BBS N.Ireland graph

Demographic trends

Productivity and survival trends for this species are not currently produced by BTO


This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglinton, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts

Conservation listings:	Europe: no SPEC category (not concentrated in Europe, conservation status favourable) (BiE04) UK: green (BoCC3)
Long-term trend:	UK: increase
Population size:	7,400 pairs in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)

Status summary

Between the 1968-72 and 1988-91 atlas periods, the Raven's range contracted from some areas of Scotland and northern England. Declines in southern Scotland and northern England were associated with large-scale afforestation (Marquiss et al. 1978), while closer sheep husbandry and conversion of pasture to arable were also implicated (Mearns 1983). A thorough survey of northwest Wales during 1998 to 2005 found at least 69% more nesting pairs than a previous survey of the same area during 1978-85 and evidence of an increase of 173% since around 1950, at a rate that accelerated after 1990 (Driver 2006). Ravens have increased along the English-Welsh border and colonised extensive new areas of the south coast, western and midland England and southern Scotland since 1988-91 (Cross 2002, Balmer et al. 2013). BBS indicates steep increase in England, Scotland and Wales since 1994. Nesting success appears to have improved, but brood size has fallen. No trend is evident in the number of fledglings per breeding attempt. Ravens have increased markedly across Europe since 1980, though with little change since 1990 (PECBMS 2013a): increases are evident in all regions but are weakest in the south and west, including UK (PECBMS 2009).







Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	16	1995-2011	281	5	-34	94		
	10	2001-2011	346	-4	-23	35		


Source	5 Period (yrs)	2006-2011 Years	409 Plots	-16 Change (%)	-35 Lower limit	12 Upper limit	Alert	Comment
BBS England		1995-2011	421	(%)	60	277		
	10	2001-2011	165	16	-21	118		
	5	2006-2011	214	-10	-33	43		
BBS Scotland	16	1995-2011	44	35	-8	97		
	10	2001-2011	46	26	-14	74		
	5	2006-2011	54	-8	-44	43		
BBS Wales	16	1995-2011	88	26	-13	95		
	10	2001-2011	101	-10	-31	7		
	5	2006-2011	103	-17	-35	1		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.



Fledglings per breeding attempt


Raven 1966-2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Raven




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	22	None					
Clutch size	43	1968-2011	14	None					Small sample
Brood size	43	1968-2011	70	Linear decline	3.15 chicks	2.91 chicks	-7.5%		
Nest failure rate at egg stage	43	1968-2011	22	Curvilinear	0.26% nests/day	0.03% nests/day	-88.5%		Small sample
Nest failure rate at chick stage	43	1968-2011	31	None					
Laying date	43	1968-2011	11	None			0 days		Small sample


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend


Key facts

Conservation listings:	Europe: no SPEC category (concentrated in Europe, conservation status favourable) (BiE04) UK: green (BoCC3)
Long-term trend:	England: fluctuating, with no long-term trend
Population size:	610,000 territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)

Status summary

Goldcrest abundance is unusually severely affected by winter weather, and the strong increase in the species' CBC/BBS index up to the mid 1970s can be interpreted as recovery from the cold winters of the early 1960s. The subsequent decline temporarily moved the species to the amber list, but its status has now been restored to green. The long-term trend looks very much like a series of damped oscillations following recovery from the 1962/63 winter. The high amplitude of year-to-year change reflects the species high breeding potential, and its sensitivity to cold winter weather. CBC had relatively poor coverage of conifer plantations, in which Goldcrests occur at increasing densities as the trees mature. A general increase in the area of prime habitat has therefore been poorly reflected in the long-term trend. BBS has recorded substantial decreases in Wales, and all UK countries show decline in the recent five-year period. The BBS PECBMS 2013a).

CBC/BBS England 1966-2012 Goldcrest



Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	44	1967-2011	262	29	-26	168		
	25	1986-2011	397	21	-12	44		
	10	2001-2011	641	-11	-19	1		
	5	2006-2011	698	-9	-14	-1		

BBS UK Source	Period (yrs)	1995-2011 Years	POBs (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
	10	2001-2011	888	-34	-40	-25	>25	
	5	2006-2011	960	-28	-34	-20	>25	
BBS England	16	1995-2011	536	15	3	33		
	10	2001-2011	629	-11	-18	0		
	5	2006-2011	679	-9	-14	-1		
BBS Scotland	16	1995-2011	93	-5	-27	33		
	10	2001-2011	104	-50	-61	-32	>25	
	5	2006-2011	119	-43	-54	-28	>25	
BBS Wales	16	1995-2011	80	-49	-63	-21	>25	
	10	2001-2011	86	-43	-52	-30	>25	
	5	2006-2011	83	-17	-30	3		
BBS N.Ireland	16	1995-2011	44	1	-32	17		
	10	2001-2011	52	-46	-53	-27	>25	
	5	2006-2011	56	-51	-56	-38	>50	

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




CBC/BBS England 1966–2012
Goldcrest




CBC/BBS England graph

BBS UK 1994–2012
Goldcrest




BBS UK graph

BBS England 1994–2012
Goldcrest




BBS England graph


BBS Scotland 1994–2012
Goldcrest



BBS Scotland graph



BBS Wales graph



BBS N.Ireland graph

Demographic trends

Productivity and survival trends for this species are not currently produced by BTO


This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts

Conservation listings:	Europe: no SPEC category (concentrated in Europe, conservation status favourable) (BiE04) UK: green (species level); amber (race obscurus, >20% of European breeders) (BoCC3)
Long-term trend:	UK, England: shallow increase
Population size:	3.6 million territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)

Status summary

Blue Tit populations have increased in abundance, in parallel with those of PECBMS 2013a).






Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	1028	37	23	53		
	25	1986-2011	1642	9	3	15		
	10	2001-2011	2611	4	1	6		
	5	2006-2011	2922	-2	-4	0		
CBC/BBS England	44	1967-2011	844	37	22	53		
	25	1986-2011	1342	7	1	13		
	10	2001-2011	2115	3	0	6		

Source	Period (yrs)	2006-2011 Years	2389 Plots (n)	1 Change (%)	3 Lower limit	2 Upper limit	Alert	Comment
CES adults	27	1984-2011	99	14	-1	33		
	25	1986-2011	104	23	9	42		
	10	2001-2011	106	12	3	24		
	5	2006-2011	106	3	-5	12		
CES juveniles	27	1984-2011	99	-17	-40	22		
	25	1986-2011	104	-10	-33	21		
	10	2001-2011	107	-7	-19	8		
	5	2006-2011	105	20	3	35		
BBS UK	16	1995-2011	2297	7	3	10		
	10	2001-2011	2611	3	0	6		
	5	2006-2011	2922	-2	-4	0		
BBS England	16	1995-2011	1850	6	1	9		
	10	2001-2011	2092	3	0	6		
	5	2006-2011	2350	-1	-3	1		
BBS Scotland	16	1995-2011	167	8	-3	18		
	10	2001-2011	192	7	-7	20		
	5	2006-2011	224	-6	-15	5		
BBS Wales	16	1995-2011	176	17	6	29		
	10	2001-2011	200	10	1	22		
	5	2006-2011	200	-1	-8	10		
BBS N.Ireland	16	1995-2011	77	3	-28	25		
	10	2001-2011	92	-11	-20	-1		
	5	2006-2011	95	-21	-29	-16		


Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




CES adults graph




CES juveniles graph




BBS UK graph




BBS England graph



BBS Scotland graph




BBS Wales graph




Demographic trends

Fledglings per breeding attempt Blue Tit 1966–2012



Laying date 1966–2012


Blue Tit




More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	509	None					
Clutch size	43	1968-2011	422	None					
Brood size	43	1968-2011	800	Linear decline	8.13 chicks	7.34 chicks	-9.6%		
Nest failure rate at egg stage	43	1968-2011	717	Linear decline	0.36% nests/day	0.19% nests/day	-47.2%		
Nest failure rate at chick stage	43	1968-2011	510	Curvilinear	0.60% nests/day	0.67% nests/day	11.7%		
Laying date	43	1968-2011	519	Curvilinear	Apr 30	Apr 21	-9 days		
Juvenile to Adult ratio (CES)	27	1984-2011	103	Smoothed trend	221 Index value	100 Index value	-55%	>50	
Juvenile to Adult ratio (CES)	25	1986-2011	108	Smoothed trend	202 Index value	100 Index value	-51%	>50	
Juvenile to Adult ratio (CES)	10	2001-2011	110	Smoothed trend	130 Index value	100 Index value	-23%		
Juvenile to Adult ratio (CES)	5	2006-2011	109	Smoothed trend	95 Index value	100 Index value	5%		


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend




Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend




Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Smoothed long-term trend in ratio of juvenile:adult birds caught - green lines indicate 85% confidence limits



Proportion of adult birds surviving to following year - green bars represent 95% confidence limits

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Great Tit

Parus major

Key facts


Conservation listings:	Europe: no SPEC category (not concentrated in Europe, conservation status favourable) (BiE04) UK: green (species level, race major); amber (race newtoni, >20% of European breeders) (BoCC3)
Long-term trend:	UK: rapid increase England: moderate increase
Population size:	2.6 million territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)
Migrant status:	Resident
Nesting habitat:	Cavity nester
Primary breeding habitat:	Woodland
Secondary breeding habitat:	
Breeding diet:	Animal
Winter diet:	Animal

Status summary

Great Tit numbers have increased fairly steadily since the 1960s, with the exception of two or three brief periods of stability or shallow decline. The BBS Lawson et al. 2012a). Laying dates have advanced by ten days since 1968. Numbers have shown widespread moderate increase across Europe since 1980 (PECBMS 2013a).



CBC/BBS UK 1966–2012

Great Tit




Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	974	110	82	140		
	25	1986-2011	1554	42	32	49		
	10	2001-2011	2498	23	19	27		
	5	2006-2011	2817	1	-2	3		
CBC/BBS England	44	1967-2011	801	98	72	130		
	25	1986-2011	1271	36	28	46		
	10	2001-2011	2026	18	15	22		
	5	2006-2011	2303	-2	-4	0		
CES adults	27	1984-2011	94	29	3	52		
	25	1986-2011	98	45	16	75		
	10	2001-2011	103	28	16	43		
	5	2006-2011	102	-10	-21	2		
CES juveniles	27	1984-2011	96	2	-25	41		
	25	1986-2011	101	11	-20	60		
	10	2001-2011	105	5	-8	21		
	5	2006-2011	104	-10	-20	7		
BBS UK	16	1995-2011	2169	45	39	50		
	10	2001-2011	2498	23	19	27		
	5	2006-2011	2817	1	-2	3		
BBS England	16	1995-2011	1748	38	34	43		
	10	2001-2011	2003	18	15	21		
	5	2006-2011	2265	-2	-4	0		
BBS Scotland	16	1995-2011	152	51	26	81		
	10	2001-2011	176	34	19	53		
	5	2006-2011	206	4	-8	15		
BBS Wales	16	1995-2011	170	56	38	78		
	10	2001-2011	194	33	20	50		
	5	2006-2011	198	9	0	21		
BBS N.Ireland	16	1995-2011	72	176	110	204		
	10	2001-2011	88	48	32	69		
	5	2006-2011	94	17	5	30		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.



CES adult abundance 1983–2012


Great Tit



CES adults graph

CES juvenile abundance 1983–2012


Great Tit



CES juveniles graph

BBS UK 1994–2012


Great Tit



BBS UK graph

BBS England 1994–2012


Great Tit



BBS England graph

BBS Scotland 1994–2012


Great Tit




BBS Scotland graph

BBS Wales 1994–2012


Great Tit





Demographic trends


Fledglings per breeding attempt Great Tit 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Great Tit




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	429	None					
Clutch size	43	1968-2011	333	Linear decline	8.02 eggs	7.49 eggs	-6.6%		
Brood size	43	1968-2011	700	Linear decline	7.19 chicks	6.31 chicks	-12.3%		
Nest failure rate at egg stage	43	1968-2011	619	Linear decline	0.53% nests/day	0.23% nests/day	-56.6%		
Nest failure rate at chick stage	43	1968-2011	429	Curvilinear	0.58% nests/day	0.63% nests/day	8.6%		
Laying date	43	1968-2011	379	Curvilinear	May 1	Apr 21	-10 days		
Juvenile to Adult ratio (CES)	27	1984-2011	102	Smoothed trend	152 Index value	100 Index value	-34%	>25	
Juvenile to Adult ratio (CES)	25	1986-2011	106	Smoothed trend	151 Index value	100 Index value	-34%	>25	
Juvenile to Adult ratio (CES)	10	2001-2011	110	Smoothed trend	121 Index value	100 Index value	-17%		
Juvenile to Adult ratio (CES)	5	2006-2011	108	Smoothed trend	100 Index value	100 Index value	0%		


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend




Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend




Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Smoothed long-term trend in ratio of juvenile:adult birds caught - green lines indicate 85% confidence limits



Proportion of adult birds surviving to following year - green bars represent 95% confidence limits

Causes of change

Demographic trends in breeding parameters do not suggest that increases in this species are due to improvements in breeding performance. There is some evidence, albeit limited, that improvements in survival rates, due to amelioration in wintering conditions, may have been responsible. Evidence for ecological drivers of the population increase is limited but increased provisioning in gardens and milder winters may have played a role.

Change factor	Primary driver	Secondary driver
Demographic	Improved survival	
Ecological	Other	Climate change

Further information on causes of change

The number of fledglings per breeding attempt has been unchanged but clutch and brood sizes have decreased (see above). Daily failure rates at the egg stage have also decreased but daily failure rates at the chick stage have increased. Thus, there is no evidence from this that improvements in breeding parameters have driven the population increase.

Increases in survival rates, due to more widespread food provision in gardens during winter is one possible explanation for the increase. Horak & Lebreton (1998) found that survival rates in Estonia were higher in urban populations than rural ones and suggested that this was partly due to supplementary feeding in gardens. Increasing winter temperature may have also played a role. Ahola et al. (2009) suggested that, for their study population in Sweden, increasingly favourable conditions in winters have enhanced the survival rates of Great Tit and resulted in the observed increase in Great Tit breeding density.

Other factors may also have influenced survival rates. There is some evidence that the beech crop production may be influential and it has been shown that survival rates can be related to beech crop (Verhulst 1992, Perdeck et al. 2000), although there is no evidence that beechmast production has gone up. Perdeck et al. (2000) provided further evidence for this as supplemental food increased survival of both juveniles and adults, supporting the winter-food limitation hypothesis. In a Finnish population, Orell (1989) reported that the high survival rates of resident juveniles after a warm August may be attributable to food availability during the time when the birds undergo their post-juvenile moult. Great Tits have advanced their laying date (see above), in line with climatic change. This has been found by several studies (e.g. Sanz 2002, Visser et al. 2009, Bauer et al. 2010).

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>


Key facts

Conservation listings:	Europe: no SPEC category (not concentrated in Europe, conservation status favourable) (BiE04) UK: green (species level, race hibernicus); amber (race britannicus, >20% of European breeders) (BoCC3)
Long-term trend:	UK, England: moderate increase
Population size:	760,000 territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)

Status summary

While other common tit species have increased, the UK Coal Tit population has been rather stable since the mid 1970s, following earlier rapid increase. The ratios of Coal Tit to Perrins 2003), however, although in these figures population change may be confounded to some degree with changes in behaviour among birds and bird ringers. The BBS PECBMS 2013a).

CBC/BBS UK 1966–2012 Coal Tit






Smoothed population index, relative to an arbitrary 100 in the year given, with 85% confidence limits in green


Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	386	79	1	188		
	25	1986-2011	594	34	12	58		
	10	2001-2011	949	13	4	23		
	5	2006-2011	1078	13	5	21		
CBC/BBS England	44	1967-2011	269	74	-2	227		
	25	1986-2011	406	35	10	64		


Source	Period (yrs)	2001-2011 Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
	5	2006-2011	741	13	3	24		
BBS UK	16	1995-2011	802	20	8	33		
	10	2001-2011	949	13	3	23		
	5	2006-2011	1078	13	5	20		
BBS England	16	1995-2011	526	29	13	45		
	10	2001-2011	625	16	8	25		
	5	2006-2011	714	14	4	25		
BBS Scotland	16	1995-2011	130	8	-6	26		
	10	2001-2011	148	10	-9	32		
	5	2006-2011	175	11	-3	27		
BBS Wales	16	1995-2011	72	2	-25	35		
	10	2001-2011	82	20	-5	39		
	5	2006-2011	82	35	14	56		
BBS N.Ireland	16	1995-2011	63	93	39	138		
	10	2001-2011	76	9	-9	34		
	5	2006-2011	80	10	-6	23		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.







BBS England graph



BBS Scotland graph



BBS Wales graph




BBS N.Ireland graph

Demographic trends

Fledglings per breeding attempt


Coal Tit 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Coal Tit




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	54	Curvilinear	6.70 fledglings	6.26 fledglings	-6.5%		
Clutch size	43	1968-2011	40	None					
Brood size	43	1968-2011	75	Curvilinear	7.35 chicks	6.87 chicks	-6.5%		
Nest failure rate at egg stage	43	1968-2011	56	Linear decline	0.50% nests/day	0.14% nests/day	-72.0%		
Nest failure rate at chick stage	43	1968-2011	59	Linear increase	0.18% nests/day	0.42% nests/day	133.3%		
Laying date	43	1968-2011	45	Linear decline	May 3	Apr 19	-14 days		


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

Key facts


Conservation listings:	Europe: no SPEC category (not concentrated in Europe, conservation status favourable) (BiE04) UK: red (>50% population decline) (BoCC3); an RBBP species UK Biodiversity Action Plan: priority species
Long-term trend:	UK, England: rapid decline
Population size:	3,400 pairs in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)
Migrant status:	Resident
Nesting habitat:	Cavity nester
Primary breeding habitat:	Woodland
Secondary breeding habitat:	
Breeding diet:	Animal
Winter diet:	Vegetation

Status summary

Willow Tits have been in decline since the mid 1970s, and have become locally extinct in an ever-growing number of former haunts. The UK conservation listing was upgraded from amber to red in 2002. Atlas surveys during 2008-11 found that the species had virtually disappeared from the southeastern part of its English range since 1988-91 (Balmer et al. 2013). The continuing decline in the CBC/BBS index through the 1990s, following a brief period of stability during the 1980s, is replicated in the CES abundance trend. All UK breeding records since 2010 should be forwarded to the Rare Breeding Birds Panel, who have developed specific PECBMS 2007, 2013a).

CBC/BBS UK 1966–2012

Willow Tit









Smoothed population index, relative to an arbitrary 100 in the year given, with 85% confidence limits in green

Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	43	-92	-96	-86	>50	
	25	1986-2011	46	-90	-94	-84	>50	Small CBC sample
	10	2001-2011	45	-64	-74	-57	>50	
	5	2006-2011	46	-47	-59	-37	>25	
CBC/BBS England	44	1967-2011	40	-91	-97	-85	>50	
	25	1986-2011	41	-89	-94	-84	>50	Small CBC sample
	10	2001-2011	40	-62	-72	-55	>50	
	5	2006-2011	42	-43	-59	-32	>25	
CES adults	27	1984-2011	18	-57	-85	-21	>50	Small sample
	25	1986-2011	18	-63	-87	-38	>50	Small sample
CES juveniles	27	1984-2011	26	-62	-83	-26	>50	
	25	1986-2011	27	-59	-81	-16	>50	
	10	2001-2011	14	-33	-62	11		Small sample
	5	2006-2011	13	-16	-48	32		Small sample
BBS UK	16	1995-2011	50	-82	-87	-76	>50	
	10	2001-2011	45	-64	-73	-57	>50	
	5	2006-2011	45	-46	-59	-35	>25	
BBS England	16	1995-2011	45	-81	-87	-74	>50	
	10	2001-2011	40	-63	-72	-51	>50	
	5	2006-2011	42	-43	-55	-29	>25	

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.






Demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Juvenile to Adult ratio (CES)	27	1984-2011	29	Smoothed trend	213 Index value	100 Index value	-53%		
Juvenile to Adult ratio (CES)	25	1986-2011	30	Smoothed trend	143 Index value	100 Index value	-30%		
Juvenile to Adult ratio (CES)	10	2001-2011	17	Smoothed trend	187 Index value	100 Index value	-47%		
Juvenile to Adult ratio (CES)	5	2006-2011	15	Smoothed trend	132 Index value	100 Index value	-24%		

For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).



Smoothed long-term trend in ratio of juvenile:adult birds caught - green lines indicate 85% confidence limits

Causes of change

Willow Tits have declined in woodland, probably because of habitat degradation. How this relates to demographic trends is unclear.

Change factor	Primary driver	Secondary driver
Demographic	Unknown	
Ecological	Changes in woodland	

Further information on causes of change

Little evidence is available regarding changes in the demography of this species but CES trends suggest a decline in productivity since 1983 (see above). Lampila et al. (2006) found that adult survival was the main driver of Willow Tit populations in northern Finland, although this was in a study in boreal forests, so the processes may not be the same as for the British population. The British subspecies shows very different habitat preferences to the Fennoscandian one, preferring wet woodland rather than conifers, emphasising that Continental studies may not be very relevant to population change in the UK.

There are several hypotheses that have been put forward to explain the cause of population declines of Willow Tit. One is that deterioration in the quality of woodland as feeding habitat for this species through canopy closure and increased browsing by deer (Perrins 2003, Siriwardena 2004, Fuller et al. 2005) has been important. The area of wet woodland and scrub is also thought to have declined as a result of drainage and the occurrence of increasingly dry summers (Vanhinsbergh et al. 2003). A field study based on former CBC sites and other woods that were known to have held the species in the past provided good evidence that the sites still holding Willow Tit tended to be wetter, so drying out of woodlands may have been a factor (Lewis et al. 2007, 2009a, 2009b). Siriwardena (2004) analysed long-term CBC trends and found that, although population trends have been stable in their preferred, wet habitats, Willow Tit have declined in woodland, probably because of habitat degradation.

A second hypothesis is that nest predation pressure, from Jays, Great Spotted Woodpeckers and grey squirrel, for example, has increased, both because some of these predators have grown more abundant (Harris et al. 1995, this report) and because restrictions in nest-site availability are likely to have forced more birds into suboptimal, more vulnerable sites. In the study mentioned above, Siriwardena (2004) found increases in Green Woodpecker abundance on CBC plots at the same time as declines in Willow Tit abundance, but this is unlikely to reflect a causal link - this woodpecker being unrecorded as a nest predator. A negative relationship between Great Spotted Woodpecker and Willow Tit abundance on farmland plots is more likely to reflect a real population effect, but farmland is only a minor habitat for the species, so it is unlikely that such a relationship has biological significance for Willow Tits nationally. There were no significant associations with other avian potential nest predators. Supporting this result, Lewis et al. (2007, 2009a, 2009b) found that sites that were known to have held the species in the past and that were still holding Willow Tits did not differ in the density of potential nest predators.

Thirdly, increases in the local populations of behaviourally dominant, sympatric species such as Blue Tit, Great Tit, Marsh Tit and Nuthatch could have led to increased competition, especially for nest-holes. There is little direct evidence specifically concerning foraging interactions involving Willow Tit in the UK but it is possible that increases in other tit species have placed extra pressure on Willow Tit populations through competition for food or nest sites (Vanhinsbergh et al. 2003). In Lanarkshire, central Scotland, Great and Blue Tits were found commonly to take over the nest sites of Willow Tit (Maxwell 2002, 2003) but it is unclear how widespread this phenomenon is. In the analysis of long-term CBC trends carried out by Siriwardena (2004), no negative relationships were found between Willow Tit and its potential competitors. Again, this was supported by field data from Lewis et al. (2007, 2009a, 2009b), who found that sites that were known to have held the species in the past and that that were still holding Willow Tits did not differ in the density of avian competitors.

Overall, therefore, habitat deterioration is the strongest candidate as the cause of Willow Tit decline nationally. As well as increasing woodland drainage, degradation has been hypothesised to have occurred via a reduction in nest-site availability resulting from falls in the amount of dead wood and number of dead trees in woodland reducing nesting opportunities (Vanhinsbergh et al. 2003). This has yet to be tested formally, however, probably because historical data on quantities of dead wood are not available.

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts


Conservation listings:	Europe: SPEC category 3 (declining) (BiE04) UK: red (>50% population decline) (BoCC3) UK Biodiversity Action Plan: priority species
Long-term trend:	UK, England: rapid decline
Population size:	41,000 territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)
Migrant status:	Resident
Nesting habitat:	Cavity nester
Primary breeding habitat:	Woodland
Secondary breeding habitat:	
Breeding diet:	Animal
Winter diet:	Animal

Status summary

Marsh Tit abundance has declined almost continuously since BTO monitoring began. The species' UK conservation listing has recently been upgraded from amber to red. Numbers have shown widespread moderate decline across Europe since 1980, though with little change since 1990 (PECBMS 2013a) and the European status of this species is no longer considered "secure" (BirdLife International 2004).

CBC/BBS UK 1966–2012




Marsh Tit




Smoothed population index, relative to an arbitrary 100 in the year given, with 85% confidence limits in green

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	98	-73	-81	-64	>50	
	25	1986-2011	129	-41	-55	-22	>25	
	10	2001-2011	163	-28	-39	-15	>25	
	5	2006-2011	173	-14	-26	-1		
CBC/BBS England	44	1967-2011	90	-72	-80	-62	>50	
	25	1986-2011	118	-40	-53	-21	>25	
	10	2001-2011	147	-25	-37	-13	>25	
	5	2006-2011	157	-9	-20	4		
BBS UK	16	1995-2011	146	-24	-36	-11		
	10	2001-2011	159	-29	-40	-18	>25	
	5	2006-2011	166	-12	-23	-2		
BBS England	16	1995-2011	133	-26	-37	-12	>25	
	10	2001-2011	145	-26	-36	-12	>25	
	5	2006-2011	154	-8	-21	5		


Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.





Demographic trends


Fledglings per breeding attempt Marsh Tit 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Marsh Tit




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	19	None					
Clutch size	43	1968-2011	14	None					Small sample
Brood size	43	1968-2011	24	None					Small sample
Nest failure rate at egg stage	43	1968-2011	21	Linear decline	0.74% nests/day	0.11% nests/day	-85.1%		Small sample
Nest failure rate at chick stage	43	1968-2011	20	None					Small sample
Laying date	43	1968-2011	14	Linear decline	Apr 28	Apr 17	-11 days		Small sample


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

Causes of change

There is good evidence that changes in the habitat quality of woodlands, particularly a loss of understorey, have been responsible for the decline in Marsh Tits. Analysis of the BTO's ring-recovery archive provides evidence that there has been a significant negative trend in annual survival rates during the period of decline, although this is based on a small sample size.

Change factor	Primary driver	Secondary driver
Demographic	Reduced survival	
Ecological	Changes in woodland	

Further information on causes of change

Analysis of the BTO's ring-recovery archive provides evidence that there has been a significant negative trend in annual survival rates during the period of decline, although this is based on a small sample size. The absence of any reduction in breeding performance as the population has declined supports a reduction in annual survival as the demographic mechanism (Siriwardena 2006). Nest failure rates have fallen during the period of decline, but no trend is evident in the number of fledglings per breeding attempt (see above).

One hypothesis relating to the causes of decline is that changes in woodland understorey have reduced habitat quality, due to increased browsing by deer (Perrins 2003, Fuller et al. 2005). Carpenter (2008) and Carpenter et al. (2010) conducted a detailed study providing good evidence that Marsh Tits were more likely to locate their territories in sections of woodland with more understorey cover. Carpenter found that birds in territories with more understorey raised more and heavier young than did birds in territories with less understorey, although this was based on only one year of data. The same study reported that understorey and low canopy sections were also important during winter while Hinsley et al. (2007) provide further evidence that this was important, showing that that Marsh Tits were selecting the understorey and habitat lower down in the woodland canopy. Another field study conducted by Broughton et al. (2006), however, did not find any difference in the amount of shrub layer in Marsh Tit territories compared to pseudo-territories, although this was from just one site and the authors noted that the understorey there was unusually healthy and complete, perhaps explaining this result.

A reduction in habitat quality through fragmentation is another possible factor that has contributed to declines, although there has been little fragmentation of woodland in a gross sense in recent years. Nevertheless, Hinsley et al. (1995) found that Marsh Tits need a minimum wood size of 0.5 ha and it's possible that habitat deterioration has reduced effective habitat patch size.

Another hypothesis concerning causes of decline relates to competition and nest predation. Marsh Tit is subdominant to both 2006) found no evidence for population effects of the Marsh Tit being outcompeted for natural nest cavities. Similarly, the same study found no evidence that avian nest predation is a major factor in the long-term

decline as Marsh Tit abundance was not significantly related to abundance in the previous year of any of the nest predators considered (Siriwardena 2006). Amar et al. (2006) found no association between population change and grey squirrel abundance and adding to this, Smart et al. (2007) conducted an initial analysis and showed that Marsh Tit declines were also unlikely to be caused by predation by grey squirrel, as presence and abundance of Marsh Tit was positively related to squirrel density.

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Woodlark

Lullula arborea

Key facts

Conservation listings:	Europe: SPEC category 2 (depleted) (BiE04) UK: amber (European status, long-term UK range contraction, localised UK breeding) (BoCC3); an RBBP species UK Biodiversity Action Plan: click here , priority species
Long-term trend:	UK: increase
Population size:	3,100 (2,500-3,700) pairs in 2006 (APEP13: Conway et al. 2009)

Status summary

This species is too rare and restricted in range for population changes to be monitored annually by BTO volunteer surveys. A 62% reduction occurred in the number of 10-km squares occupied between 1968-72 and 1988-91; the species had ceased to breed in Wales and in several southern English counties over this period (Gibbons et al. 1993). Sitters et al. (1996) report that the UK population increased from c.250 pairs in 1986 to c.600 pairs in 1993, probably helped by mild winters and increased habitat availability due to storm damage in plantations, forest restocking, and heathland management. A repeat national survey in 1997 showed that the population had increased further, accompanied by expansion of the range into new areas (Wotton & Gillings 2000; for more information, [click here](#)). A further repeat in 2006 recorded an increase since 1997 of 88% accompanied by major range expansion, with a pair breeding in Wales for the first time since 1981 (Conway et al. 2009; also Balmer et al. 2013).


Farmland setaside, especially close to forest, was valuable additional habitat for the expanding population, although clutch sizes may be lower there than in more traditional habitats (Wright et al. 2007). Climate change may benefit Woodlark, because it is able to make more nesting attempts in warmer years (Wright et al. 2009). The cold 2009/10 winter may, however, have brought about the small reduction in numbers reported to RBBP for 2010 (Holling & RBBP 2012). The small NRS sample suggests that nest failure rates have become less frequent at the egg stage. There has been no trend, however, in the number of fledglings per breeding attempt. Human disturbance at heathland sites apparently reduces population density, but the effects are partly offset by higher breeding productivity at lower densities (Mallord et al. 2007). The species' partial recovery in numbers and range resulted in a move from the red to the amber list at the 2009 review (Eaton et al. 2009).

Population changes in detail

Annual breeding population changes for this species are not currently monitored by BTO

Demographic trends


Fledglings per breeding attempt Woodlark 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Woodlark




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	22	None					
Clutch size	43	1968-2011	20	None					Small sample
Brood size	43	1968-2011	32	None					
Nest failure rate at egg stage	43	1968-2011	23	Curvilinear	5.31% nests/day	2.10% nests/day	-60.5%		Small sample
Nest failure rate at chick stage	43	1968-2011	34	None					
Laying date	43	1968-2011	22	None			0 days		Small sample


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend




Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

Key facts

Conservation listings:	Europe: SPEC category 3 (depleted) (BiE04) UK: red (species level, race arvensis); amber (race scotica, >20% of European breeders) (BoCC3) UK Biodiversity Action Plan: click here , priority species
Long-term trend:	England: rapid decline
Population size:	1.5 million territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend for England)
Migrant status:	Resident
Nesting habitat:	Ground nester
Primary breeding habitat:	Farmland
Secondary breeding habitat:	
Breeding diet:	Animal
Winter diet:	Animal

Status summary

The Skylark declined rapidly from the mid 1970s until the mid 1980s, when the rate of decline slowed. BBS data show further decline, recently extending to Scotland. The BBS PECBMS 2013a).




Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	44	1967-2011	601	-62	-70	-55	>50	
	25	1986-2011	970	-32	-39	-22	>25	
	10	2001-2011	1524	-10	-13	-5		
	5	2006-2011	1721	-8	-11	-5		
BBS UK	16	1995-2011	1687	-22	-27	-17		
	10	2001-2011	1828	-15	-19	-10		
	5	2006-2011	2013	-14	-17	-11		
BBS England	16	1995-2011	1345	-23	-26	-18		
	10	2001-2011	1473	-10	-13	-5		
	5	2006-2011	1638	-9	-11	-5		
BBS Scotland	16	1995-2011	204	-19	-30	-5		
	10	2001-2011	208	-17	-26	-6		
	5	2006-2011	229	-22	-27	-14		
BBS Wales	16	1995-2011	103	-14	-30	4		
	10	2001-2011	111	-15	-27	4		
	5	2006-2011	112	-9	-17	2		
BBS N.Ireland	16	1995-2011	34	-50	-60	-41	>50	
	10	2001-2011	35	-57	-68	-50	>50	
	5	2006-2011	33	-33	-42	-22	>25	

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




CBC/BBS England 1966–2012
Skylark




CBC/BBS England graph

BBS UK 1994–2012
Skylark




BBS UK graph


BBS England 1994–2012
Skylark




BBS England graph



BBS Scotland graph




BBS Wales graph



BBS N.Ireland graph

Demographic trends


Fledglings per breeding attempt Skylark 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Skylark




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	42	Linear increase	1.06 fledglings	1.57 fledglings	48.3%		
Clutch size	43	1968-2011	35	Linear increase	3.37 eggs	3.70 eggs	9.8%		
Brood size	43	1968-2011	65	Linear increase	3.18 chicks	3.45 chicks	8.7%		
Nest failure rate at egg stage	43	1968-2011	44	None					
Nest failure rate at chick stage	43	1968-2011	53	Linear decline	4.77% nests/day	3.07% nests/day	-35.6%		
Laying date	43	1968-2011	19	Curvilinear	May 24	May 30	6 days		Small sample


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

Causes of change

There is good evidence to indicate that the most likely cause of declines in Skylark is agricultural intensification, specifically the change from spring to autumn sowing of cereals, which reduces the number of breeding attempts possible and may also reduce overwinter survival due to loss of winter stubbles.

Change factor	Primary driver	Secondary driver
Demographic	Reduced breeding succes	
Ecological	Agricultural intensification	

Further information on causes of change

Demographic trends presented here show that there has been a general increase in the number of fledglings per breeding attempt, because clutch size and brood size have increased while the daily failure rate of nests at the chick stage has gone down. Chamberlain & Crick (1999) and Siriwardena et al. (2000b) found that breeding success per nesting attempt increased during the steepest period of decline, suggesting that these demographic changes have not contributed to the causes of population decline. The available data do not allow tests for effects of survival. Conversely, it is easy to test for effects on breeding success, especially locally and with respect to contemporary as opposed to historical land use. This creates a big imbalance in the amounts of evidence available.

Agricultural intensification has been put forward as the ultimate cause of Skylark declines. The relevant changes in agriculture have been decreases in preferred crops (spring cereals and cereal stubble) and an increase in unfavourable habitats (winter cereals, oilseed rape and intensively managed or grazed grass) (Chamberlain & Siriwardena 2000). There is good evidence that the most likely cause of the decline is the change from spring to autumn sowing of cereals. This practice restricts opportunities for late-season nesting attempts, because the crop is by then too tall. Chamberlain et al. (2000a) used habitat data from CBC surveys to show that the occurrence of autumn-sown, winter cereals increased from 33% to 78% between 1965 and 1995. Evans et al. (1995) and Wilson et al. (1997) all found that Skylarks deserted areas of autumn-sown crops as soon the sward reached a critical height, which occurred before the end of the breeding season. Jenny (1990), Chamberlain et al. (1999, 2000a, 2000b) and Donald & Vickery (2000) all recorded low and seasonally declining densities of Skylarks in cereals and suggested that this was at least partly due to the effects of changing vegetation structure. As well as preventing nesting, crop development also influences the positioning of the nests that are produced and hence their productivity: as the crop develops the birds are forced to nest closer to tramlines with a consequent increase in nest predation rate (Donald & Vickery 2000, Morris & Gilroy 2008). Analyses by Chamberlain & Crick (1999) provided detailed evidence from both regional and habitat-based analyses that the greatest declines in Skylark numbers were associated with agricultural habitat, although their evidence suggests that different patterns of decline were unlikely to be due to differences in breeding success per attempt between habitats. However, Siriwardena et al. (2001) showed that the population trend can be explained by national changes in crop areas, together with a cold winter in 1981/82.

There is also some evidence that the increase in autumn sowing may depress overwinter survival by reducing the area of stubbles (Wilson et al. 1997, Donald & Vickery

2000, 2001). Donald & Vickery (2001) used data from BTO and RSPB studies to show that, in winter, cereal stubbles were strongly selected by Skylarks, probably owing to the presence of spilt grain and regenerating weeds, and go on to state that the area of stubbles has declined greatly in recent years. Gillings et al. (2005) identified better population performance in areas with extensive winter stubble, presumably because overwinter survival is relatively high. Note, however, that definitive evidence about Skylark survival rates and what may have influenced them is not available because the species is rarely ringed and ring-recovery sample sizes are extremely small.

Use of pesticides and associated declines as declines in weed populations and weed-seed abundance have been suggested as another factor in the decline of Skylarks (Wilson 2001). Wilson et al. (1997) found higher densities of Skylarks in organic systems. Chamberlain & Crick (1999) suggest that the use of toxic pesticides mediated through effects on food supplies may be responsible for declines in invertebrate food, due to non-target insects being killed by insecticide and insect food-plants being killed by herbicide. However, since this would in theory affect breeding success, it doesn't seem to have been a problem. Donald et al. (2001) state that, although recent agricultural changes have affected diet and possibly body condition of nestlings, these effects are unlikely to have been an important factor in recent population declines. There may also be implications for overwinter survival, as herbicides reduce weeds, and hence seeds for the winter, making stubbles and uncropped land less valuable as a food resource. However, the increases in pesticide use have happened at the same time as the switch to autumn sowing, so is hard to detect this as a specific effect.

There is some evidence to suggest that high densities of raptors may reduce the abundance of local Skylark populations (Amaet al. 2008b). Chamberlain & Crick (1999) state that recovery of [Sparrowhawk](#) numbers has been most evident in the most intensively farmed areas, and that this is correlated with the declines in Skylark numbers across habitats and regions. However, this apparent link cannot be taken as evidence of a causal relationship as there have been many other broad-scale changes in the countryside that are at least as well correlated with Skylark changes. They state that it is doubtful whether predation alone could account for the decreases in Skylark numbers.

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts


Conservation listings:	Europe: SPEC category 3 (depleted) (BiE04) UK: amber (European status) (BoCC3)
Long-term trend:	UK: fluctuating, with no long-term trend
Population size:	54,000-174,000 nests in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)

Status summary

This species is unusually difficult to monitor, because active and inactive nest holes are difficult to distinguish, and because whole colonies frequently disperse or shift to new locations as suitable sand cliffs are created and destroyed. WBS counts were of apparently occupied nest holes along riverbanks but BBS and WBBS record birds seen. WBS/WBBS suggests a stable or shallowly increasing population, with wide fluctuations, although the ongoing decrease since the late 1990s has been steep enough to raise BTO alerts. BBS counts show clearly that large year-to-year changes occur, but do not yet reveal a clear long-term trend. Nest-record samples are small, but indicate that nest failure rates have decreased enormously since the 1960s; brood size has fallen and no trend can be detected in the numbers of fledglings per breeding attempt. Rainfall in the species' trans-Saharan wintering grounds prior to the birds' arrival promotes annual survival and thus abundance in the following breeding season (Szep 1995). Annual survival rates from RAS sites in the UK for 1990-2004 were correlated positively with minimum monthly rainfall during the wet season in West Africa (Robinson et al. 2008). Mark-recapture in Cheshire during 1981-2003 found that, after allowing for the effects of African rainfall, some demographic measures were density dependent, with adult survival low when wintering densities (measured as the size of the western European population) were high and recruitment low when the local Cheshire population was high (Norman & Peach 2013). This study did not replicate an earlier finding (Cowley & Siriwardena 2005) that summer rainfall on the breeding grounds has a negative influence on survival rates through the following winter.

WBS/WBBS 1977-2012

Sand Martin




Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
WBS/WBBS waterways	33	1978-2011	46	26	-20	151		

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
	10	2001-2011	94	10	-21	53		
	5	2006-2011	90	27	2	63		
BBS UK	16	1995-2011	127	39	-5	119		
	10	2001-2011	140	44	10	76		
	5	2006-2011	158	10	-12	37		
BBS England	16	1995-2011	82	11	-22	35		
	10	2001-2011	89	-1	-25	23		
	5	2006-2011	98	-9	-28	13		


Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.



WBS/WBBS waterways graph




BBS UK graph



BBS England graph


Fledglings per breeding attempt Sand Martin 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012


Sand Martin




More on demographic trends


Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	36	None					
Clutch size	43	1968-2011	41	None					
Brood size	43	1968-2011	58	Curvilinear	3.22 chicks	2.35 chicks	-27.1%		
Nest failure rate at egg stage	43	1968-2011	37	Curvilinear	2.04% nests/day	0.07% nests/day	-96.6%		
Nest failure rate at chick stage	43	1968-2011	61	Linear decline	1.75% nests/day	0.04% nests/day	-97.7%		
Laying date	43	1968-2011	42	None			0 days		

For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).







Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of adult birds surviving to following year - error bars represent 95% confidence limits

Key facts


Conservation listings:	Europe: SPEC category 3 (depleted) (BiE04) UK: amber (European status) (BoCC3)
Long-term trend:	England: possible shallow increase
Population size:	860,000 territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend for England)

Status summary

Swallow was originally amber-listed partly on the strength of a decline on CBC plots in the early 1980s, but continues to qualify through its widespread decline across the European continent (BirdLife International 2004). Moderate decrease is also evident across Europe in the period since 1980 (PECBMS 2013a). Later modelling of UK population change from CBC gave evidence of fluctuations but not for long-term decline (Robinson et al. 2003). BBS data suggest increases in England, Scotland and Wales since 1994. The BBS Baillie & Peach 1992). More particularly, population change has been shown to be correlated with rainfall in the western Sahel prior to the birds' spring passage through West Africa, but with neither cattle numbers nor nest-site availability in the UK (Robinson et al. 2003). Annual survival rates from RAS sites in the UK for 1998-2004 were correlated positively with mean monthly rainfall during the early austral summer in southern Africa (Robinson et al. 2008). It is likely that, in eastern parts of the UK, the loss of livestock farming and grazed grassland, together with arable intensification, has caused the Swallow population to decline, while an increase in the area of pasture in the west and north has promoted a population increase which apparently has more than compensated for declines elsewhere (Evans & Robinson 2004). A link between regional changes in the availability of preferred feeding habitats and the regional patterns of UK population change again suggests that habitat change on the breeding grounds may explain population trend, at least partly (Henderson et al. 2007). Brood size increased up to the late 1980s, and may now be falling again, while nest losses have increased and the number of fledglings per breeding attempt shows no trend. Climatic warming is leading to both an earlier start and later finish to the breeding season for European Swallows, but there has been increased chick mortality in hot, dry summers and reduced post-fledging survival because of poor conditions for birds migrating through North Africa (Turner 2009).

CBC/BBS England 1966–2012

Swallow




Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	44	1967-2011	615	35	-3	83		
	25	1986-2011	1034	58	27	87		
	10	2001-2011	1715	27	20	34		
	5	2006-2011	1963	11	6	15		
BBS UK	16	1995-2011	1919	36	28	43		
	10	2001-2011	2173	14	7	19		
	5	2006-2011	2432	2	-3	5		
BBS England	16	1995-2011	1481	43	34	51		
	10	2001-2011	1678	27	20	34		
	5	2006-2011	1901	11	6	15		
BBS Scotland	16	1995-2011	170	34	17	58		
	10	2001-2011	188	11	-1	27		
	5	2006-2011	213	-1	-11	15		
BBS Wales	16	1995-2011	170	31	11	49		
	10	2001-2011	192	-3	-12	12		
	5	2006-2011	198	-11	-21	-1		
BBS N.Ireland	16	1995-2011	85	3	-19	31		
	10	2001-2011	100	-22	-32	-13		
	5	2006-2011	104	-24	-31	-15		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




CBC/BBS England 1966–2012
Swallow




CBC/BBS England graph

BBS UK 1994–2012
Swallow




BBS UK graph


BBS England 1994–2012
Swallow




BBS England graph



BBS Scotland graph




BBS Wales graph



BBS N.Ireland graph

Demographic trends


Fledglings per breeding attempt Swallow 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012


Swallow




More on demographic trends


Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	498	None					
Clutch size	43	1968-2011	471	None					
Brood size	43	1968-2011	820	Curvilinear	4.12 chicks	4.16 chicks	1.0%		
Nest failure rate at egg stage	43	1968-2011	581	None					
Nest failure rate at chick stage	43	1968-2011	499	Linear increase	0.31% nests/day	0.43% nests/day	38.7%		
Laying date	43	1968-2011	207	Curvilinear	Jun 19	Jun 11	-8 days		

For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).







Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of adult birds surviving to following year - error bars represent 95% confidence limits

Key facts


Conservation listings:	Europe: SPEC category 3 (declining) (BiE04) UK: amber (25-50% population decline) (BoCC3)
Long-term trend:	England: probable rapid decline
Population size:	510,000 (360,000-670,000) pairs in 2009 (APEP13: distance sampling estimate for 2006 (Newson et al. 2008) updated using BBS trend)

Status summary

The House Martin's loosely colonial nesting habits and its strong association with human settlements mean that it is extraordinarily difficult to monitor. Anecdotal evidence of decline is often unreliable, because demise of a colony may be balanced by single nests or small groups becoming established elsewhere. For these reasons, study areas should be large, covered thoroughly, and ideally randomly selected. The available long-term data suggest a rapid decline, although BBS's first decade or so showed an increase. The BBS BirdLife International 2004). There has been widespread moderate decrease across Europe since 1980, though with little change since 1990 (PECBMS 2013a). Annual survival rates from RAS sites in the UK for 1994-2004 were correlated positively with maximum monthly rainfall in West Africa; some decline in survival rate is apparent over this period but does not correspond to the population decline (Robinson et al. 2008).

CBC/BBS England 1966-2012

House Martin



Smoothed population index, relative to an arbitrary 100 in the year given, with 85% confidence limits in green

Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	44	1967-2011	292	-65	-91	0		Small CBC sample
	25	1986-2011	499	-55	-83	40		Small CBC sample
	10	2001-2011	811	-18	-24	-11		
	5	2006-2011	876	-19	-25	-14		

BBS UK Source	Period (yrs)	1995-2011 Years	2011 (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
	10	2001-2011	1022	-16	-23	-9		
	5	2006-2011	1092	-15	-22	-9		
BBS England	16	1995-2011	730	-17	-24	-7		
	10	2001-2011	801	-18	-25	-10		
	5	2006-2011	860	-20	-26	-14		
BBS Scotland	16	1995-2011	62	114	45	216		
	10	2001-2011	72	-12	-34	35		
	5	2006-2011	82	-6	-26	29		
BBS Wales	16	1995-2011	87	16	-12	50		
	10	2001-2011	93	-16	-35	7		
	5	2006-2011	92	-7	-29	21		
BBS N.Ireland	16	1995-2011	42	69	2	171		
	10	2001-2011	51	7	-23	47		
	5	2006-2011	53	10	-13	33		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




CBC/BBS England 1966–2012
House Martin




CBC/BBS England graph

BBS UK 1994–2012
House Martin




BBS UK graph

BBS England 1994–2012
House Martin




BBS England graph


BBS Scotland 1994–2012
House Martin



BBS Scotland graph




BBS Wales graph



BBS N.Ireland graph

Demographic trends

Productivity and survival trends for this species are not currently produced by BTO



Proportion of adult birds surviving to following year - error bars represent 95% confidence limits

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>


Key facts

Conservation listings:	Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe) (BiE04) UK: green (BoCC3); an RBBP species
Long-term trend:	England, Wales: increase
Population size:	2,000 males in 2006-10 (APEP13: RBBP data)

Status summary

Cetti's Warblers were first recorded in Britain as recently as 1961, as part of their range expansion across northwest Europe (Bonham & Robertson 1975). Colonisation, which began in Kent in 1972 or 1973, continues to be monitored annually by Holling & RBBP 2013. Numbers and breeding range increased spectacularly during the first 12 years, with Norfolk and Dorset gradually overtaking Kent as the main host counties (Gibbons et al. 1993, Wotton et al. 1998). Severe winters after 1978 led to the temporary extinction of the Kent population in 1988. Populations in milder regions continued to grow, but overall the UK population fell by over a third between 1984 and 1986. In the absence of severe winters during 1986-2009, increase and range expansion continued. The first breeding records north of the Humber were made in 2006 (Holling & RBBP 2009). Much constant-effort ringing takes place in prime Cetti's Warbler habitat; despite the comparative rarity of this species, therefore, CES population and productivity indices are already available (Robinson et al. 2007). CES data confirm the species' sensitivity to cold winters, which appears to have become more evident as the breeding range has expanded into more testing climates. The index has fallen owing to cold winters since 2010. Numbers have shown a remarkably steep rate of increase across Europe since 1990, but no longer-term trend is available (PECBMS 2013a).

CES adult abundance 1986-2012
Cetti's Warbler



Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CES adults	10	2001-2011	14	73	5	315		Small sample
	5	2006-2011	18	7	-18	52		Small sample

CES juveniles Source	Period (yrs)	2001-2011 Years	#ots (n)	Change (%)	Lower limit	Upper limit	Alert	Small sample Comment
	5	2006-2011	18	3	-36	74		Small sample

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




CES adult abundance 1986–2012
Cetti's Warbler



CES adults graph

CES juvenile abundance 1986–2012
Cetti's Warbler



CES juveniles graph

Demographic trends

Productivity and survival trends for this species are not currently produced by BTO

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>


Key facts

Conservation listings:	Europe: no SPEC category (not concentrated in Europe, conservation status favourable) (BiE04) UK: green (species level); amber (race <i>rosaceus</i> , >20% of European breeders) (BoCC3)
Long-term trend:	England: rapid increase
Population size:	340,000 territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend for England)

Status summary

This species undergoes wide fluctuations in numbers between breeding seasons, suffering heavy mortality when winters are severe, but is able to recover quickly by virtue of its high breeding potential. Numbers were low after the severe winters of the early 1960s and again during a series of relatively cold winters beginning in the late 1970s, but have not fallen after the cold winters since 2010. The starting years of the 25-year and longest monitoring periods coincided with troughs in population, thus exaggerating the long-term trend. CBC/BBS index trends show progressive increases in Long-tailed Tit abundance beginning in the early 1980s. The BBS Crick & Sparks 1999). Numbers across Europe have shown little change since 1980 (PECBMS 2013a).

CBC/BBS England 1966-2012 Long-tailed Tit



Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	44	1967-2011	401	122	63	240		
	25	1986-2011	626	105	76	135		
	10	2001-2011	986	21	13	29		
	5	2006-2011	1145	20	14	26		

CES adults Source	27 Period (yrs)	1984-2011 Years	81 Plots	35 Change (%)	4 Lower limit	82 Upper limit	Alert	Comment
	10	2001-2011	91	-3	-13	9		
	5	2006-2011	91	6	-6	17		
CES juveniles	27	1984-2011	75	56	9	135		
	25	1986-2011	80	103	47	192		
	10	2001-2011	87	-17	-31	-2		
	5	2006-2011	88	-1	-16	13		
BBS UK	16	1995-2011	942	32	22	44		
	10	2001-2011	1110	22	13	33		
	5	2006-2011	1283	23	15	31		
BBS England	16	1995-2011	830	24	15	34		
	10	2001-2011	980	21	14	30		
	5	2006-2011	1135	20	15	27		
BBS Wales	16	1995-2011	59	37	-1	108		
	10	2001-2011	65	31	8	85		
	5	2006-2011	68	19	2	61		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




CBC/BBS England 1966-2012
Long-tailed Tit




CBC/BBS England graph

CES adult abundance 1983-2012
Long-tailed Tit






CES adults graph

CES juvenile abundance 1983-2012
Long-tailed Tit




CES juveniles graph



Demographic trends

Fledglings per breeding attempt


Long-tailed Tit 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Long-tailed Tit




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	33	None					
Clutch size	43	1968-2011	42	Curvilinear	7.75 eggs	7.01 eggs	-9.5%		
Brood size	43	1968-2011	33	None					
Nest failure rate at egg stage	43	1968-2011	59	Linear decline	3.22% nests/day	0.85% nests/day	-73.6%		
Nest failure rate at chick stage	43	1968-2011	40	Linear increase	0.75% nests/day	1.94% nests/day	158.7%		
Laying date	43	1968-2011	53	Linear decline	Apr 21	Apr 5	-16 days		
Juvenile to Adult ratio (CES)	27	1984-2011	87	Smoothed trend	115 Index value	100 Index value	-13%		
Juvenile to Adult ratio (CES)	25	1986-2011	92	Smoothed trend	78 Index value	100 Index value	28%		
Juvenile to Adult ratio (CES)	10	2001-2011	99	Smoothed trend	115 Index value	100 Index value	-13%		
Juvenile to Adult ratio (CES)	5	2006-2011	99	Smoothed trend	100 Index value	100 Index value	0%		


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend




Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend




Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Smoothed long-term trend in ratio of juvenile:adult birds caught - green lines indicate 85% confidence limits



Proportion of adult birds surviving to following year - green bars represent 95% confidence limits

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts


Conservation listings:	Europe: SPEC category 2 (declining) (BiE04) UK: red (breeding decline, European status) (BoCC3) UK Biodiversity Action Plan: priority species
Long-term trend:	UK: decline
Population size:	6,500 (5,900-7,000) males in 2009 (APEP13: 1984-85 estimate (Bibby 1989) updated using CBC/BBS trend)
Migrant status:	Long-distance migrant
Nesting habitat:	Ground nester
Primary breeding habitat:	Woodland
Secondary breeding habitat:	
Breeding diet:	Animal
Winter diet:	Animal

Status summary

Wood Warblers, which have a westerly distribution in Britain, were monitored relatively poorly until BBS began. Little change was evident at the few CBC plots on which the species occurred (Marchant et al. 1990). The species' breeding range varied little between the first two atlas periods (Gibbonset al. 1993), but has subsequently withdrawn from large areas of lowland England (Balmer et al. 2013). BBS shows a rapid and significant decline since 1994, and accordingly the species was moved from the green to the amber list in 2002; the continued decline warranted a further shift to the red list in 2009. With declines evident across northern and western Europe, this previously 'secure' species is now provisionally categorised as 'declining' (BirdLife International 2004). Numbers have shown widespread moderate decrease across Europe since 1980 (PECBMS 2013a).

BBS UK 1994–2012


Wood Warbler



Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	16	1995-2011	52	-69	-78	-54	>50	
	10	2001-2011	49	-40	-57	-18	>25	
	5	2006-2011	53	-32	-49	-10	>25	

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See here for more information.




Demographic trends

Fledglings per breeding attempt Wood Warbler 1966–2012



Laying date 1966–2012

Wood Warbler




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	24	None					
Clutch size	43	1968-2011	19	None					Small sample
Brood size	43	1968-2011	39	None					
Nest failure rate at egg stage	43	1968-2011	24	Linear decline	1.81% nests/day	0.75% nests/day	-58.6%		Small sample
Nest failure rate at chick stage	43	1968-2011	31	Curvilinear	2.47% nests/day	4.06% nests/day	64.4%		
Laying date	43	1968-2011	35	Curvilinear	May 24	May 17	-7 days		


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

Causes of change

There is little evidence explaining either the demographic or ecological drivers of the decline in this species and the causes are largely unknown.

Change factor	Primary driver	Secondary driver
Demographic	Unknown	
Ecological	Unknown	

Further information on causes of change

There is little evidence regarding any demographic causes of the decline of this species. Nest failures now seem more likely to occur at the chick stage (see above), although nest record samples are small. There has been no trend in the number of fledglings per breeding attempt.

Bibby (1989) postulated that soils, climate, competition or predator numbers have probably had an effect on Wood Warbler numbers but provided no evidence in support. Smart et al. (2007) state that the loss of oak trees, the decrease in canopy cover, and the large increases in understorey cover could have been particularly detrimental for Wood Warbler, but again, direct evidence to validate this is largely lacking. Smart et al. (2007) and Amar et al. (2006) did find that Wood Warblers have tended to decrease more in woods with fewer dead limbs on trees and at sites surrounded by more woodland, which suggests that changes in dead wood could be important or that dead limbs could be a surrogate for other changes in habitat, although Smart et al. (2007) found an overall increase in the amount of dead wood, which should have been beneficial for this species. In another Welsh study, Mallord et al. (2012b) found that Wood Warblers were associated with a number of structural features of the study woods, which could relate to their past management; they suggest that management should aim to restore habitat quality through introducing a moderate grazing regime.

Studies in Poland, where an average of over 70% of nests were lost and predators were responsible for over 80% of the losses, have reported that varying predation rates were a main factor responsible for variation in production between years and habitats (Wesolowski 1985). Wesolowski & Maziarz (2009) provided further evidence relating to this, finding that both Wood Warbler numbers and ratios of their change were significantly negatively correlated with rodent numbers. However, the authors state that, since Wood Warblers simply don't settle in areas with high rodent outbreaks, the changes probably reflect changes in distribution rather than overall trends. In Wales, nest predators during 2009-11 were mainly avian and rates of predation did not appear to have changed since 1982-84 (Mallord et al. 2012a). This species is a long-distance migrant and therefore changes outside the breeding grounds cannot be ruled out.


Key facts

Conservation listings:	Europe: no SPEC category (not concentrated in Europe, conservation status favourable) (BiE04) UK: green (BoCC3)
Long-term trend:	UK, England: moderate increase
Population size:	1.2 million territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)

Status summary

Chiffchaff abundance declined in the late 1960s/early 1970s in common with that of other trans-Saharan warblers (Siriwardena et al. 1998a). After remaining stable for a decade, the population recovered strongly, and has continued to increase. This recovery is evident from both CBC/BBS and CES data. The BBS PECBMS 2013a).

CBC/BBS UK 1966-2012
Chiffchaff





Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	644	91	56	140		
	25	1986-2011	1043	238	191	277		
	10	2001-2011	1737	63	55	69		
	5	2006-2011	1970	48	44	53		
CBC/BBS England	44	1967-2011	549	94	55	155		
	25	1986-2011	885	236	195	280		

Source	10 Period (yrs)	2001-2011 Years 2006-2011	1463 Plots (1969-2011)	58 Change (%)	51 Lower limit	64 Upper limit	Alert	Comment
CES adults	27	1984-2011	73	261	136	501		
	25	1986-2011	77	235	130	461		
	10	2001-2011	89	60	40	91		
	5	2006-2011	90	45	34	59		
CES juveniles	27	1984-2011	84	199	101	394		
	25	1986-2011	88	157	78	317		
	10	2001-2011	96	47	27	79		
	5	2006-2011	97	53	34	80		
BBS UK	16	1995-2011	1453	93	85	104		
	10	2001-2011	1737	64	57	70		
	5	2006-2011	1970	47	43	52		
BBS England	16	1995-2011	1221	90	82	101		
	10	2001-2011	1453	58	51	62		
	5	2006-2011	1651	43	39	47		
BBS Scotland	16	1995-2011	47	414	257	716		
	10	2001-2011	62	118	67	193		
	5	2006-2011	77	69	41	111		
BBS Wales	16	1995-2011	137	75	48	109		
	10	2001-2011	160	80	59	99		
	5	2006-2011	165	55	40	74		
BBS N.Ireland	16	1995-2011	33	50	5	68		
	10	2001-2011	40	81	39	144		
	5	2006-2011	47	89	56	115		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




CES adult abundance 1983-2012
Chiffchaff




CES adults graph

CES juvenile abundance 1983-2012
Chiffchaff




CES juveniles graph

BBS UK 1994-2012
Chiffchaff




BBS UK graph

BBS England 1994-2012
Chiffchaff




BBS England graph

BBS Scotland 1994-2012
Chiffchaff




BBS Scotland graph

BBS Wales 1994-2012
Chiffchaff




BBS Wales graph



Demographic trends


Fledglings per breeding attempt Chiffchaff 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Chiffchaff




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	40	None					
Clutch size	43	1968-2011	39	None					
Brood size	43	1968-2011	44	Linear decline	5.10 chicks	4.70 chicks	-7.8%		
Nest failure rate at egg stage	43	1968-2011	50	None					
Nest failure rate at chick stage	43	1968-2011	43	None					
Laying date	43	1968-2011	58	Linear decline	May 16	May 2	-14 days		
Juvenile to Adult ratio (CES)	27	1984-2011	90	Smoothed trend	123 Index value	100 Index value	-19%		
Juvenile to Adult ratio (CES)	25	1986-2011	95	Smoothed trend	161 Index value	100 Index value	-38%	>25	
Juvenile to Adult ratio (CES)	10	2001-2011	102	Smoothed trend	143 Index value	100 Index value	-30%	>25	
Juvenile to Adult ratio (CES)	5	2006-2011	103	Smoothed trend	97 Index value	100 Index value	3%		


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend




Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend




Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Smoothed long-term trend in ratio of juvenile:adult birds caught - green lines indicate 85% confidence limits



Proportion of adult birds surviving to following year - green bars represent 95% confidence limits

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>


Key facts

Conservation listings:	Europe: no SPEC category (not concentrated in Europe, conservation status favourable) (BiE04) UK: amber (species level, 25-50% population decline; race trochilus, 25-50% population decline, European status) (BoCC3)
Long-term trend:	England: rapid decline
Population size:	2.4 million territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)

Status summary

Willow Warbler abundance has shown regionally different trends within the UK (Morrison et al. 2010, 2013, Massimino et al. 2013, Balmer et al. 2013). The overall CBC/BBS trend shows a rapid decline during the 1980s and early 1990s, after 20 years of relative stability, and, on the strength of a 31% decline on CBC plots between 1974 and 1999, the species was moved from the green to the amber list. This decline occurred mainly in southern Britain, however, accompanied by a fall in survival rates there (Peach et al. 1995a), with Scottish populations remaining unaffected. BBS figures since 1994 indicate a stark contrast between an upward trend in Scotland and in Northern Ireland, and continued severe decreases in England, with no overall trend in Wales. The BBS Fuller et al. 2005). The recent population decline is associated with a shallow decline in productivity as measured by CES and with a substantial increase in nest failure rates, which raises NRS concern (Leech & Barimore 2008). There is a small but significant decrease in the number of fledglings per breeding attempt. Average laying dates have shifted earlier by more than a week, perhaps in response to recent climatic warming (Crick & Sparks 1999). Numbers have shown widespread moderate decrease across Europe since 1980 (PECBMS 2013a, Lehtikoinen et al. 2014).

CBC/BBS England 1966–2012 Willow Warbler





Smoothed population index, relative to an arbitrary 100 in the year given, with 85% confidence limits in green

Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	44	1967-2011	477	-57	-70	-44	>50	
	25	1986-2011	713	-57	-66	-50	>50	

Source	Period (yrs)	2001-2011 Years	95% CIs (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
	5	2006-2011	1071	6	0	11		
CES adults	27	1984-2011	89	-66	-73	-57	>50	
	25	1986-2011	93	-62	-68	-52	>50	
	10	2001-2011	86	-17	-26	-4		
	5	2006-2011	86	-11	-20	-1		
CES juveniles	27	1984-2011	93	-36	-56	-2	>25	
	25	1986-2011	97	-33	-54	-3	>25	
	10	2001-2011	95	9	-10	35		
	5	2006-2011	95	21	-1	47		
BBS UK	16	1995-2011	1389	5	-2	11		
	10	2001-2011	1451	12	5	16		
	5	2006-2011	1611	22	16	27		
BBS England	16	1995-2011	913	-29	-34	-23	>25	
	10	2001-2011	918	-8	-14	-2		
	5	2006-2011	1012	5	0	11		
BBS Scotland	16	1995-2011	213	38	22	52		
	10	2001-2011	230	18	5	29		
	5	2006-2011	268	23	15	34		
BBS Wales	16	1995-2011	159	0	-15	15		
	10	2001-2011	171	21	7	32		
	5	2006-2011	173	33	21	46		
BBS N.Ireland	16	1995-2011	79	114	55	152		
	10	2001-2011	93	46	27	65		
	5	2006-2011	97	54	38	72		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




CES juvenile abundance 1983-2012
Willow Warbler




CES juveniles graph

BBS UK 1994-2012
Willow Warbler




BBS UK graph

BBS England 1994-2012
Willow Warbler




BBS England graph

BBS Scotland 1994-2012
Willow Warbler




BBS Scotland graph

BBS Wales 1994-2012
Willow Warbler




BBS Wales graph

BBS N. Ireland 1994-2012
Willow Warbler



BBS N.Ireland graph


Fledglings per breeding attempt Willow Warbler 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Willow Warbler




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	69	Linear decline	3.59 fledglings	3.16 fledglings	-12.1%		
Clutch size	43	1968-2011	49	None					
Brood size	43	1968-2011	149	Linear increase	5.12 chicks	5.38 chicks	5.0%		
Nest failure rate at egg stage	43	1968-2011	69	Linear increase	0.94% nests/day	1.76% nests/day	87.2%		
Nest failure rate at chick stage	43	1968-2011	128	Linear increase	1.54% nests/day	2.04% nests/day	32.5%		
Laying date	43	1968-2011	88	Curvilinear	May 19	May 10	-9 days		
Juvenile to Adult ratio (CES)	27	1984-2011	99	Smoothed trend	131 Index value	100 Index value	-24%		
Juvenile to Adult ratio (CES)	25	1986-2011	103	Smoothed trend	120 Index value	100 Index value	-17%		
Juvenile to Adult ratio (CES)	10	2001-2011	101	Smoothed trend	102 Index value	100 Index value	-2%		
Juvenile to Adult ratio (CES)	5	2006-2011	101	Smoothed trend	86 Index value	100 Index value	16%		


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend




Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend




Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Smoothed long-term trend in ratio of juvenile:adult birds caught - green lines indicate 85% confidence limits



Proportion of adult birds surviving to following year - green bars represent 95% confidence limits

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>


Key facts

Conservation listings:	Europe: no SPEC category (concentrated in Europe, conservation status favourable) (BiE04) UK: green (BoCC3)
Long-term trend:	UK, England: rapid increase
Population size:	1.2 million territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)
Migrant status:	Short-distance migrant
Nesting habitat:	Above-ground nester
Primary breeding habitat:	Woodland
Secondary breeding habitat:	
Breeding diet:	Animal
Winter diet:	Animal

Status summary

Blackcap abundance in the UK has increased consistently since the late 1970s, a trend common to all habitats and evident from both the CBC/BBS and the CES indices. Overall increase has occurred despite a reduction in habitat quality for Blackcap, and other species dependent on the understorey, brought about by deer browsing in young woodland (Holt et al. 2012d). The BBS Balmer et al. 2013), is indicated by the BBS trend. Numbers have shown widespread moderate increase across Europe since 1980 (PECBMS 2013a).



CBC/BBS UK 1966–2012 Blackcap




Smoothed population index, relative to an arbitrary 100 in the year given, with 85% confidence limits in green

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	697	272	218	398		
	25	1986-2011	1119	174	156	200		
	10	2001-2011	1815	65	61	72		
	5	2006-2011	2066	54	53	62		
CBC/BBS England	44	1967-2011	606	234	181	326		
	25	1986-2011	966	151	135	172		
	10	2001-2011	1547	53	50	60		
	5	2006-2011	1765	46	43	52		
CES adults	27	1984-2011	91	116	79	163		
	25	1986-2011	95	132	99	178		
	10	2001-2011	100	59	47	72		
	5	2006-2011	99	26	16	35		
CES juveniles	27	1984-2011	93	28	-10	73		
	25	1986-2011	97	56	23	106		
	10	2001-2011	102	20	-2	42		
	5	2006-2011	101	10	-5	24		
BBS UK	16	1995-2011	1545	133	124	151		
	10	2001-2011	1815	65	62	74		
	5	2006-2011	2066	54	53	62		
BBS England	16	1995-2011	1321	108	99	119		
	10	2001-2011	1537	53	51	60		
	5	2006-2011	1748	45	43	52		
BBS Scotland	16	1995-2011	57	357	215	534		
	10	2001-2011	74	110	59	177		
	5	2006-2011	94	76	53	119		
BBS Wales	16	1995-2011	120	163	118	211		
	10	2001-2011	139	82	60	111		
	5	2006-2011	145	70	56	97		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




CES adult abundance 1983-2012
Blackcap




CES adults graph

CES juvenile abundance 1983-2012
Blackcap




CES juveniles graph

BBS UK 1994-2012
Blackcap




BBS UK graph

BBS England 1994-2012
Blackcap




BBS England graph

BBS Scotland 1994-2012
Blackcap




BBS Scotland graph

BBS Wales 1994-2012
Blackcap



BBS Wales graph


Fledglings per breeding attempt Blackcap 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Blackcap




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	40	None					
Clutch size	43	1968-2011	40	None					
Brood size	43	1968-2011	47	None					
Nest failure rate at egg stage	43	1968-2011	52	None					
Nest failure rate at chick stage	43	1968-2011	40	None					
Laying date	43	1968-2011	42	Curvilinear	May 20	May 7	-13 days		
Juvenile to Adult ratio (CES)	27	1984-2011	98	Smoothed trend	174 Index value	100 Index value	-43%	>25	
Juvenile to Adult ratio (CES)	25	1986-2011	103	Smoothed trend	201 Index value	100 Index value	-50%	>50	
Juvenile to Adult ratio (CES)	10	2001-2011	106	Smoothed trend	142 Index value	100 Index value	-30%	>25	
Juvenile to Adult ratio (CES)	5	2006-2011	105	Smoothed trend	139 Index value	100 Index value	-28%	>25	


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend




Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend




Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Smoothed long-term trend in ratio of juvenile:adult birds caught - green lines indicate 85% confidence limits



Proportion of adult birds surviving to following year - green bars represent 95% confidence limits

Causes of change

The causes of the increase in this species remain unknown.

Change factor	Primary driver	Secondary driver
Demographic	Unknown	
Ecological	Unknown	

Further information on causes of change

According to CES, productivity has fluctuated markedly, obscuring any long-term trend in CES or NRS data. Survival rates have been stable. Using data from France, Julliard (2004) found that population growth rate was under the additive influence of survival and recruitment.

The trend towards earlier laying, amounting to an advance of almost two weeks since 1968, may be a response to recent climate change (Crick & Sparks 1999, Croxton et al. 2006). The more rapid increase in Scotland indicated by BBS suggests that climatic warming may be allowing this species to extend its range northwards (Hewson et al. 2007).


This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts

Conservation listings:	Europe: no SPEC category (concentrated in Europe, conservation status favourable) (BiE04) UK: green (BoCC3)
Long-term trend:	UK, England: shallow decline
Population size:	170,000 territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)

Status summary

Garden Warbler abundance has varied alongside that of other trans-Saharan migrant warblers (Siriwardena et al. 1998b), probably reflecting the influence of changes in their winter environment. Despite large short-term fluctuations in abundance, the CBC/BBS data suggest that the population may be in long-term decline. The BBS PECBMS 2013a).







Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	234	-13	-44	41		
	25	1986-2011	351	-11	-27	12		
	10	2001-2011	475	3	-8	14		
CBC/BBS England	5	2006-2011	527	9	-1	19		
	44	1967-2011	196	-23	-50	28		
	25	1986-2011	292	-23	-36	-1		


Source	Period (yrs)	2001-2011 Years	BBS (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
	5	2006-2011	435	0	-7	8		
CES adults	27	1984-2011	64	-5	-34	46		
	25	1986-2011	67	-9	-35	24		
	10	2001-2011	63	10	-5	24		
	5	2006-2011	63	2	-9	18		
CES juveniles	27	1984-2011	64	-50	-65	-10	>25	
	25	1986-2011	67	-38	-56	-12	>25	
	10	2001-2011	64	-25	-41	-8	>25	
	5	2006-2011	61	-32	-44	-17	>25	
BBS UK	16	1995-2011	446	-10	-21	1		
	10	2001-2011	475	4	-7	12		
	5	2006-2011	527	9	0	17		
BBS England	16	1995-2011	362	-21	-29	-12		
	10	2001-2011	383	-9	-18	-1		
	5	2006-2011	426	-2	-7	7		
BBS Wales	16	1995-2011	57	-11	-40	27		
	10	2001-2011	59	12	-14	41		
	5	2006-2011	61	21	-6	50		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.







CES juveniles graph



BBS UK graph




BBS England graph



BBS Wales graph

Fledglings per breeding attempt


Garden Warbler 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Garden Warbler




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	19	None					
Clutch size	43	1968-2011	17	None					Small sample
Brood size	43	1968-2011	25	None					Small sample
Nest failure rate at egg stage	43	1968-2011	23	None					Small sample
Nest failure rate at chick stage	43	1968-2011	20	Linear increase	1.10% nests/day	2.73% nests/day	148.2%		Small sample
Laying date	43	1968-2011	22	Linear decline	May 28	May 19	-9 days		Small sample
Juvenile to Adult ratio (CES)	27	1984-2011	78	Smoothed trend	264 Index value	100 Index value	-62%	>50	
Juvenile to Adult ratio (CES)	25	1986-2011	82	Smoothed trend	262 Index value	100 Index value	-62%	>50	
Juvenile to Adult ratio (CES)	10	2001-2011	79	Smoothed trend	149 Index value	100 Index value	-33%	>25	
Juvenile to Adult ratio (CES)	5	2006-2011	77	Smoothed trend	157 Index value	100 Index value	-36%	>25	


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend




Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend




Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Smoothed long-term trend in ratio of juvenile:adult birds caught - green lines indicate 85% confidence limits



Proportion of adult birds surviving to following year - green bars represent 95% confidence limits

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts


Conservation listings:	Europe: no SPEC category (not concentrated in Europe, conservation status favourable) (BiE04) UK: green (BoCC3)
Long-term trend:	UK, England: fluctuating, with no long-term trend
Population size:	74,000 territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)

Status summary

Lesser Whitethroat abundance was roughly stable (albeit with short-term fluctuations) from the 1960s until the late 1980s, but the CBC/BBS and CES trends provide evidence for a subsequent moderate decline that lasted into the late 1990s. These changes were statistically significant, and large enough over the relevant periods to trigger BTO alerts. BBS has subsequently shown a significant sharp upturn, but this contrasts strongly with the continued decrease recorded by CES ringers. Wide fluctuations in survival and productivity have been recorded by CES ringers, and may be influencing population change, but pressures during migration and in winter are the most likely causes of any decline (Fuller et al. 2005). There has been little long-term change across Europe since 1980 (PECBMS 2013a).

CBC/BBS UK 1966-2012

Lesser Whitethroat






Smoothed population index, relative to an arbitrary 100 in the year given, with 85% confidence limits in green


Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	144	4	-28	49		
	25	1986-2011	212	-16	-31	0		
	10	2001-2011	306	28	14	43		
	5	2006-2011	353	15	4	23		


CBC/BBS England Source	44 Period (yrs)	1967-2011 Years	138 Plots	-4 Change (%)	-33 Lower limit	44 Upper limit	Alert	Comment
	10	2001-2011	292	26	13	40		
	5	2006-2011	338	15	5	25		
CES adults	27	1984-2011	38	-64	-82	-49	>50	
	25	1986-2011	39	-67	-79	-55	>50	
	10	2001-2011	33	-22	-41	0		
	5	2006-2011	32	-18	-36	5		
CES juveniles	27	1984-2011	44	-61	-78	-30	>50	
	25	1986-2011	46	-65	-78	-37	>50	
	10	2001-2011	42	-35	-57	-3	>25	
	5	2006-2011	39	-24	-48	13		
BBS UK	16	1995-2011	267	4	-12	18		
	10	2001-2011	304	35	17	53		
	5	2006-2011	350	17	6	27		
BBS England	16	1995-2011	255	0	-15	15		
	10	2001-2011	291	30	15	44		
	5	2006-2011	336	15	6	25		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.






CES juveniles graph



BBS UK graph




BBS England graph


Demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Juvenile to Adult ratio (CES)	27	1984-2011	53	Smoothed trend	110 Index value	100 Index value	-9%		
Juvenile to Adult ratio (CES)	25	1986-2011	56	Smoothed trend	131 Index value	100 Index value	-24%		
Juvenile to Adult ratio (CES)	10	2001-2011	50	Smoothed trend	135 Index value	100 Index value	-26%		
Juvenile to Adult ratio (CES)	5	2006-2011	48	Smoothed trend	126 Index value	100 Index value	-21%		

For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).



Smoothed long-term trend in ratio of juvenile:adult birds caught - green lines indicate 85% confidence limits



Proportion of adult birds surviving to following year - green bars represent 95% confidence limits

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts


Conservation listings:	Europe: no SPEC category (concentrated in Europe, conservation status favourable) (BiE04) UK: amber (25-50% decline, 1969-2006) (BoCC3)
Long-term trend:	UK, England: rapid decline (followed by increase)
Population size:	1.1 million territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)

Migrant status:	Long-distance migrant
Nesting habitat:	Above-ground nester
Primary breeding habitat:	Farmland
Secondary breeding habitat:	
Breeding diet:	Animal
Winter diet:	Animal

Status summary

Whitethroat numbers had been stable for a few years up to 1968 but, despite a normal departure for their West African wintering grounds in autumn 1968, crashed by around 70% between the 1968 and 1969 breeding seasons (Winstanley et al. 1974). They fluctuated around their lower level until the mid 1980s, since when the population has sustained a consistent shallow recovery. Recovery of the UK population has been most apparent along linear waterways. The BBS PECBMS 2013a). The limited extent of UK recovery, coupled with change in the BoCC criteria, has resulted in the species moving from the green to the amber list at the latest review (Eaton et al. 2009).

CBC/BBS UK 1966–2012
Whitethroat



Smoothed population index, relative to an arbitrary 100 in the year given, with 85% confidence limits in green


Population changes in detail







Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	596	-56	-68	-42	>50	
	25	1986-2011	955	133	101	166		
	10	2001-2011	1530	25	22	32		
	5	2006-2011	1732	22	20	28		
CBC/BBS England	44	1967-2011	516	-56	-69	-38	>50	
	25	1986-2011	825	137	103	179		
	10	2001-2011	1314	24	21	30		
	5	2006-2011	1491	22	20	27		
WBS/WBBS waterways	36	1975-2011	80	160	5	344		
	25	1986-2011	104	291	179	462		
	10	2001-2011	158	20	6	33		
	5	2006-2011	151	17	8	30		
CES adults	27	1984-2011	62	-18	-45	3		
	25	1986-2011	65	-2	-29	30		
	10	2001-2011	69	28	6	54		
	5	2006-2011	68	41	27	63		
CES juveniles	27	1984-2011	67	42	-22	176		
	25	1986-2011	71	56	-24	253		
	10	2001-2011	73	31	-2	74		
	5	2006-2011	74	47	15	101		
BBS UK	16	1995-2011	1338	40	34	50		
	10	2001-2011	1520	26	21	32		
	5	2006-2011	1714	21	20	28		
BBS England	16	1995-2011	1156	38	33	49		
	10	2001-2011	1309	24	21	31		
	5	2006-2011	1483	22	20	27		
BBS Scotland	16	1995-2011	79	112	63	196		
	10	2001-2011	91	36	18	75		
	5	2006-2011	104	12	3	31		
BBS Wales	16	1995-2011	83	-4	-18	16		
	10	2001-2011	94	18	7	42		
	5	2006-2011	100	27	18	48		


Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB








BBS Scotland graph



BBS Wales graph

Demographic trends


Fledglings per breeding attempt Whitethroat 1966-2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Whitethroat




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	43	None					
Clutch size	43	1968-2011	31	None					
Brood size	43	1968-2011	68	None					
Nest failure rate at egg stage	43	1968-2011	44	Curvilinear	1.02% nests/day	1.66% nests/day	62.7%		
Nest failure rate at chick stage	43	1968-2011	51	None					
Laying date	43	1968-2011	20	Curvilinear	May 27	May 17	-10 days		Small sample
Juvenile to Adult ratio (CES)	27	1984-2011	77	Smoothed trend	94 Index value	100 Index value	6%		
Juvenile to Adult ratio (CES)	25	1986-2011	81	Smoothed trend	167 Index value	100 Index value	-40%	>25	
Juvenile to Adult ratio (CES)	10	2001-2011	83	Smoothed trend	155 Index value	100 Index value	-36%	>25	
Juvenile to Adult ratio (CES)	5	2006-2011	83	Smoothed trend	102 Index value	100 Index value	-2%		


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend




Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend




Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Smoothed long-term trend in ratio of juvenile:adult birds caught - green lines indicate 85% confidence limits



Proportion of adult birds surviving to following year - green bars represent 95% confidence limits

Causes of change

There is good evidence that the major changes in the population of this species have been driven by conditions on its wintering grounds and so are related to overwinter survival.

Change factor	Primary driver	Secondary driver
Demographic	Decreased survival	
Ecological	Changes on wintering grounds	

Further information on causes of change

In a pioneering study, Winstanley et al. (1974) provided good evidence to link the 1969 crash to drought in the Whitethroat's wintering grounds in the western Sahel, just south of the Sahara Desert. Correspondingly, Baillie & Peach (1992) found that breeding performance was poorly correlated with population changes. They found that fluctuations in losses of adult birds were correlated with conditions on the wintering grounds, and were correlated with Sahel rainfall. Thus, the population appears to be limited by food resources on the wintering grounds, because rainfall in the dry Sahelian landscape promotes greater invertebrate abundance. There has been no long-term trend in the number of fledglings per breeding attempt (see above). Productivity, as measured by CES, rose during the 1980s and has since fluctuated and fallen back.

More recent work has provided good evidence that the density of Whitethroats wintering in the Sahel is correlated with the number and size of trees, and that the increase in overall density of trees was related to an increase in Whitethroats in the area (Stevens et al. 2010). Wilson & Cresswell (2006) found that Whitethroats were most common in areas with intermediate tree heights. They suggest that Whitethroats appear to be able to survive in extremely degraded habitats, yet may be vulnerable to the disappearance of *Salvadora* trees, the fruit of which assists pre-migratory fattening. This is likely to be a separate mechanism to the earlier rainfall mechanism contributing to the population decline and is probably linked to the more recent gradual increase.

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Grasshopper Warbler

Locustella naevia

Key facts


Conservation listings:	Europe: no SPEC category (concentrated in Europe, conservation status favourable) (BiE04) UK: red (>50% population decline) (BoCC3) UK Biodiversity Action Plan: priority species
Long-term trend:	UK: decline
Population size:	16,000 territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)
Migrant status:	Long-distance migrant
Nesting habitat:	Above-ground nester
Primary breeding habitat:	Wetland
Secondary breeding habitat:	
Breeding diet:	Animal
Winter diet:	Animal

Status summary

Grasshopper Warbler was previously amber-listed because of a contraction in range during the period preceding the 1988-91 Atlas (Gibbons et al. 1993). The CBC index suffered from small and severely dwindling sample sizes, but the available data indicate a rapid population decline between the mid 1960s and mid 1980s, when numbers became too small for annual monitoring (Marchant et al. 1990). On this basis, the species is now red-listed. The BBS shows wide fluctuations in abundance since 1994, but no clear trend. There has been little overall change across Europe since 1980 but a moderate decline is evident since 1990 (PECBMS 2013a).

BBS UK 1994–2012


Grasshopper Warbler




Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	16	1995-2011	81	24	2	74		
	10	2001-2011	92	25	15	70		
	5	2006-2011	110	12	10	50		
BBS England	16	1995-2011	37	-9	-32	43		
	10	2001-2011	42	-3	-18	35		
	5	2006-2011	53	15	-3	63		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See here for more information.



BBS UK graph



BBS England graph

Demographic trends

Productivity and survival trends for this species are not currently produced by BTO

Causes of change

The demographic and ecological causes of population change in this species are largely unknown.

Change factor	Primary driver	Secondary driver
Demographic	Unknown	
Ecological	Unknown	

Further information on causes of change

There are not enough data to carry out demographic analyses for this species and the causes of the decline, both demographic and ecological, are largely unknown.

Although there is no specific evidence available, as this species is a migrant, it is possible that it has suffered from changes in conditions in the African Sahel zone along with some other trans-Saharan migrants.

Another hypothesis, again lacking good evidence to support or refute it, is that the decline is related to a recent decrease in the amount of suitable scrub habitat preferred by breeding Grasshopper Warblers. There are strong pointers that structural aspects of preferred habitat are important, including heterogeneity, and it seems likely that breeding habitat is limited, at least in some parts of Britain (Gilbert 2012). However, the Grasshopper Warbler's decline has been fairly steep and perhaps too rapid for gradual changes in scrub habitat availability or post-afforestation decline to be major factors (Riddiford 1983).


This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglinton, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts

Conservation listings:	Europe: no SPEC category (concentrated in Europe, conservation status favourable) (BiE04) UK: green (BoCC3)
Long-term trend:	UK: shallow decline England: moderate decline
Population size:	290,000 territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)

Status summary

The trend in England is apparently of moderate decline, but this is uncertain because the long-term changes are partly obscured by shorter fluctuations in numbers. Detailed analysis of BTO data sets has shown that much of the year-to-year variation in population size is driven by changes in adult survival rates which, in turn, are related to changes in rainfall on their wintering grounds, which lie just south of the Sahara Desert, in the West African Sahel (Peach et al. 1991). The smoothed CBC/BBS and WBS/WBBS trends show four troughs in population, related to years of poor West African rainfall, with a low point in 1984-85. The CES, which provides the biggest Sedge Warbler sample, shows the most recent three of the same troughs. Daily nest failure rates at the egg stage have increased slightly but the number of fledglings per breeding attempt has shown no change. CES productivity data show a sustained decrease since the late 1980s. There has been little long-term change across Europe since 1980 (PECBMS 2013a).




Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	147	-19	-51	14		
	25	1986-2011	222	49	12	80		
	10	2001-2011	326	-6	-20	6		

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	44	1967-2011	99	-30	-65	2		
	25	1986-2011	144	36	6	81		
	10	2001-2011	209	-7	-18	7		
	5	2006-2011	232	13	2	27		
WBS/WBBS waterways	36	1975-2011	71	-37	-57	-9	>25	
	25	1986-2011	87	-17	-36	6		
	10	2001-2011	118	-21	-34	-14		
	5	2006-2011	107	3	-10	14		
CES adults	27	1984-2011	66	-42	-59	-26	>25	
	25	1986-2011	70	-50	-62	-38	>25	
	10	2001-2011	71	-35	-43	-27	>25	
	5	2006-2011	71	-13	-24	-3		
CES juveniles	27	1984-2011	64	59	-3	413		
	25	1986-2011	68	23	-26	111		
	10	2001-2011	69	-5	-29	30		
	5	2006-2011	69	8	-13	38		
BBS UK	16	1995-2011	298	11	-6	30		
	10	2001-2011	321	-4	-17	7		
	5	2006-2011	354	9	-1	19		
BBS England	16	1995-2011	190	1	-13	22		
	10	2001-2011	207	-5	-15	7		
	5	2006-2011	228	13	3	27		
BBS Scotland	16	1995-2011	54	35	-3	97		
	10	2001-2011	56	-1	-29	22		
	5	2006-2011	63	9	-9	33		







Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




CBC/BBS UK graph



CBC/BBS England graph




Fledglings per breeding attempt Sedge Warbler 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012


Sedge Warbler




More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	39	None					
Clutch size	43	1968-2011	35	None					
Brood size	43	1968-2011	55	None					
Nest failure rate at egg stage	43	1968-2011	41	Curvilinear	1.47% nests/day	1.67% nests/day	13.6%		
Nest failure rate at chick stage	43	1968-2011	47	None					
Laying date	43	1968-2011	47	Curvilinear	May 29	May 20	-9 days		
Juvenile to Adult ratio (CES)	27	1984-2011	72	Smoothed trend	250 Index value	100 Index value	-60%	>50	
Juvenile to Adult ratio (CES)	25	1986-2011	76	Smoothed trend	191 Index value	100 Index value	-48%	>25	
Juvenile to Adult ratio (CES)	10	2001-2011	78	Smoothed trend	119 Index value	100 Index value	-16%		
Juvenile to Adult ratio (CES)	5	2006-2011	79	Smoothed trend	115 Index value	100 Index value	-13%		


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend




Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend




Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Smoothed long-term trend in ratio of juvenile:adult birds caught - green lines indicate 85% confidence limits



Proportion of adult birds surviving to following year - green bars represent 95% confidence limits

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts


Conservation listings:	Europe: no SPEC category (concentrated in Europe, conservation status favourable) (BiE04) UK: green (BoCC3)
Long-term trend:	UK: rapid increase England: moderate increase
Population size:	130,000 (100,000-160,000) pairs in 2009 (APEP13: distance sampling estimate for 2006 (Newson et al. 2008) updated using BBS trend)
Migrant status:	Long-distance migrant
Nesting habitat:	Above-ground nester
Primary breeding habitat:	Wetland
Secondary breeding habitat:	
Breeding diet:	Animal
Winter diet:	Animal

Status summary

This species has an unusually clumped distribution, with very high breeding concentrations in Phragmites reedbeds, where numbers are very hard to census. CES, which has many sites in reedbeds, ought perhaps to be a better measure of population change than either CBC/BBS or WBS/WBBS, where the species is encountered mainly at low density or in linear habitats. Both CBC/BBS and WBS/WBBS show progressive strong increases. CES, however, shows a decline from 1983 until the early 1990s, followed by a partial recovery, and another more recent decline. Population increase, as indicated by the census work, accords with the remarkable range expansion the species has achieved since the 1960s, as recorded by atlas projects. West Wales, northwest and northeast England were colonised, as was the east coast of Ireland, between 1968-72 and 1988-91 (Gibbons et al. 1993), and the species is now regular as far north as the Tay reedbeds (Robertson 2003, Balmer et al. 2013). There has been little long-term change across Europe since 1980 (PECBMS 2013a).

CBC/BBS UK 1966–2012



Reed Warbler




Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	66	109	24	333		
	25	1986-2011	99	62	22	146		
	10	2001-2011	149	8	-7	25		
	5	2006-2011	172	2	-11	14		
CBC/BBS England	44	1967-2011	63	87	20	234		
	25	1986-2011	94	52	24	103		
	10	2001-2011	141	10	-1	27		
	5	2006-2011	162	7	-4	22		
WBS/WBBS waterways	30	1981-2011	43	88	3	281		
	25	1986-2011	48	85	18	237		
	10	2001-2011	71	0	-15	16		
	5	2006-2011	65	2	-10	13		
CES adults	27	1984-2011	56	-29	-45	-9	>25	
	25	1986-2011	59	-23	-42	-5		
	10	2001-2011	60	-13	-25	-3		
	5	2006-2011	62	5	-6	15		
CES juveniles	27	1984-2011	58	37	-32	183		
	25	1986-2011	61	26	-21	99		
	10	2001-2011	63	-23	-36	-3		
	5	2006-2011	65	1	-14	24		
BBS UK	16	1995-2011	128	31	8	52		
	10	2001-2011	149	12	-4	26		
	5	2006-2011	172	3	-10	16		
BBS England	16	1995-2011	122	30	13	55		
	10	2001-2011	141	13	-1	28		
	5	2006-2011	162	7	-5	19		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.







WBS/WBBS waterways graph




CES adults graph



CES juveniles graph




BBS UK graph



BBS England graph


Fledglings per breeding attempt Reed Warbler 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012


Reed Warbler




More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	141	Linear increase	2.34 fledglings	2.84 fledglings	21.7%		
Clutch size	43	1968-2011	148	None					
Brood size	43	1968-2011	165	None					
Nest failure rate at egg stage	43	1968-2011	183	Linear decline	1.68% nests/day	1.28% nests/day	-23.8%		
Nest failure rate at chick stage	43	1968-2011	142	Linear decline	1.96% nests/day	0.62% nests/day	-68.4%		
Laying date	43	1968-2011	208	Curvilinear	Jun 17	Jun 6	-11 days		
Juvenile to Adult ratio (CES)	27	1984-2011	62	Smoothed trend	77 Index value	100 Index value	29%		
Juvenile to Adult ratio (CES)	25	1986-2011	66	Smoothed trend	84 Index value	100 Index value	19%		
Juvenile to Adult ratio (CES)	10	2001-2011	67	Smoothed trend	115 Index value	100 Index value	-13%		
Juvenile to Adult ratio (CES)	5	2006-2011	69	Smoothed trend	107 Index value	100 Index value	-6%		


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend




Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend




Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Smoothed long-term trend in ratio of juvenile:adult birds caught - green lines indicate 85% confidence limits



Proportion of adult birds surviving to following year - green bars represent 95% confidence limits

Causes of change

Breeding performance has increased, with some suggestion that this may be related to warming climate or improved habitat management, although the evidence for this is sparse.

Change factor	Primary driver	Secondary driver
Demographic	Increased breeding success	
Ecological	Climate change	

Further information on causes of change

There is some evidence to suggest that this species has benefited from warmer climates. Reed Warblers have shown a trend towards earlier laying (see above), which can be partly explained by recent climate change (Crick & Sparks 1999, Halupka et al. 2008). Halupka et al. (2008) analysed changes in breeding parameters of Polish Reed Warblers, studied during 12 breeding seasons between 1970 and 2006, and found that the onset of breeding advanced with warming temperatures, although the end of breeding did not change, thus resulting in an extension of the breeding season. The lengthening of the laying period by about three weeks meant that more birds were able to rear second broods. Furthermore, mean temperature during May-July correlated negatively with the proportion of nests that failed and there was some evidence of a positive relationship with the number of fledglings. The data show a linear increase in the numbers of fledglings per breeding attempt, and a small improvement is apparent in CES productivity, although there is no available evidence to suggest that this is related to changing climate. Breeding performance as measured by brood size has also improved slightly.

Both CBC/BBS and WBS/WBBS trends show progressive moderate increases perhaps linked to increasingly sensitive management of small and linear wetland sites. Thaxter et al. (2006) analysed data from two sites and found indirect evidence linking good habitat management to local abundance and survival.

As this species is a migrant it is possible that factors operating outside the breeding season may be responsible for changes in population in the UK. Thaxter et al. (2006) found that, unlike in the 2004) found that the French Reed Warbler population appears to be strongly regulated and that population growth rate was more influenced by survival rate than by recruitment.

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts


Conservation listings:	Europe: no SPEC category (not concentrated in Europe, conservation status favourable) (BiE04) UK: green (BoCC3)
Long-term trend:	UK, England: rapid increase
Population size:	220,000 territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)
Migrant status:	Resident
Nesting habitat:	Cavity nester
Primary breeding habitat:	Woodland
Secondary breeding habitat:	
Breeding diet:	Animal
Winter diet:	Vegetation

Status summary

Nuthatch abundance in the UK has increased rapidly since the mid 1970s. Despite minor setbacks during the 1990s, there is no indication yet of a halt to the upward trend. This increase has been accompanied by a range expansion into northern England and southern Scotland (Balmer et al. 2013). The BBS PECBMS 2013a).




CBC/BBS UK 1966–2012


Nuthatch




Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	226	251	162	365		
	25	1986-2011	355	121	90	165		
	10	2001-2011	577	57	47	67		
	5	2006-2011	659	29	23	37		
CBC/BBS England	44	1967-2011	195	263	163	418		
	25	1986-2011	304	124	86	163		
	10	2001-2011	493	62	53	75		
	5	2006-2011	573	28	22	36		
BBS UK	16	1995-2011	469	88	65	105		
	10	2001-2011	570	57	45	69		
	5	2006-2011	648	27	21	35		
BBS England	16	1995-2011	397	92	72	113		
	10	2001-2011	488	62	51	74		
	5	2006-2011	564	27	21	35		
BBS Wales	16	1995-2011	70	49	20	83		
	10	2001-2011	81	25	6	49		
	5	2006-2011	81	15	2	33		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.






BBS England graph



BBS Wales graph

Demographic trends


Fledglings per breeding attempt Nuthatch 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Nuthatch




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	54	Linear increase	3.71 fledglings	5.28 fledglings	42.5%		
Clutch size	43	1968-2011	31	None					
Brood size	43	1968-2011	76	Linear increase	4.98 chicks	5.61 chicks	12.5%		
Nest failure rate at egg stage	43	1968-2011	54	Linear decline	0.94% nests/day	0.20% nests/day	-78.7%		
Nest failure rate at chick stage	43	1968-2011	62	Linear decline	0.47% nests/day	0.19% nests/day	-59.6%		
Laying date	43	1968-2011	31	Linear decline	May 2	Apr 20	-12 days		


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

Causes of change

The demographic causes of the population increase appear to be an increase in the number of fledglings per breeding attempt, larger brood sizes and a decrease in daily failure rates. However, it is unclear what the ecological drivers of these changes are.

Change factor	Primary driver	Secondary driver
Demographic	Increased breeding success	
Ecological	Unknown	

Further information on causes of change

The number of fledglings per breeding attempt has increased strongly, through an increase in brood size and a fall in nest failure rates.

There is little evidence relating to Nuthatch population change in the UK. However, studies from Europe provide evidence that mild winters are likely to have helped this species. Kallander (1997) used a long-term data set (1977-91) to provide good evidence that Nuthatches in a Swedish national park had a population size in spring which co-varied positively with winter temperatures and suggest that increases in population size may be associated with increasing mean winter temperature. Nilsson (1982, 1987) also found that mortality was concentrated in winter and that starvation was probably the major cause. However, a long-term study in Poland from 1975 to 1990 found that bird numbers in spring were not significantly correlated with the severity of the preceding winter, though winter survival was higher in the unusually mild winter of 1989/90, which had a rich supply of hornbeam seeds (Wesolowski & Stawarczyk 1991). It is not possible to say whether such factors have also operated in the UK, as the climate here is considerably less extreme.

Several studies have also reported a link between population size and the size of food availability in the autumn. A study of two Nuthatch populations in Sweden provided good evidence that autumn population size was correlated with the size of the hazelnut crop, suggesting food supplies play a role, although beechmast crop was not correlated with overwinter survival and nor was autumn population size correlated with the population density in spring (Enoksson & Nilsson 1983, Enoksson 1990). In the studies by Nilsson mentioned above, the main density-dependent factor, recruitment of young of the year to the autumn population, was positively related to the current beechmast supply and negatively to the density of adults (Nilsson 1982, 1987). A long-term study in Poland from 1975 to 1990 also found that Nuthatch numbers seemed to be influenced by autumn seed supply and also availability of caterpillars in the preceding spring (Wesolowski & Stawarczyk 1991). Another continental study in Europe found that local survival in autumn was higher in beechmast years for juveniles, but not for adults and that local winter survival was not higher in years with than in years without beechmast (Matthysen 1989). Thus there is some evidence that increases in population size are linked to food supplies, but again, this has not been directly tested for UK birds.

Although there is no direct evidence available, Nuthatches are known to favour dead wood, and so it is possible that they may have benefited from the increase in dead

wood in the UK (Amar et al. 2010).

In Belgium, competition for nest sites with the non-native, invasive Strubbe & Matthysen 2009). However, there is evidence showing that this is not a problem in the UK at present (Newson et al. 2011).

The reasons for the poorer performance of Nuthatches in Wales are unknown.


This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts

Conservation listings:	Europe: no SPEC category (not concentrated in Europe, conservation status favourable) (BiE04) UK: green (species level); amber (race britannica, >20% of European breeders) (BoCC3)
Long-term trend:	UK, England: fluctuating, with no long-term trend
Population size:	200,000 territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)

Status summary

The UK Treecreeper population peaked in the mid 1970s, but has been roughly stable since about 1980. Intensive study has shown that Treecreeper numbers and survival rates are reduced by wet winter weather (Peach et al. 1995b). The influence of cold weather is also evident in the low start to the index, following the severe winter of 1962/63, and the trough around 1980. Census data suggest a minor decline has occurred since the early 1980s, but CES adult captures have increased for much of this period. Productivity, calculated using CES data, shows fluctuations around a long-term shallow increase. There has been a significant fall in nest failure rates at the egg stage and a small increase in overall nest success. The trend towards earlier laying can be partly explained by recent climate change (Crick & Sparks 1999). There has been little long-term change across Europe since 1980, but moderate decrease since 1990 (PECBMS 2013a).






Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	206	14	-14	49		
	25	1986-2011	287	-2	-18	17		
	10	2001-2011	381	4	-8	18		
	5	2006-2011	421	3	-4	13		

CBC/BBS England Source	Period (yrs)	1967-2011 Years	Points (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
	25	1986-2011	221	-2	-18	15		
	10	2001-2011	284	7	-3	18		
	5	2006-2011	318	12	3	22		
CES adults	27	1984-2011	37	39	-12	107		
	25	1986-2011	39	91	35	159		
	10	2001-2011	39	29	7	52		
	5	2006-2011	36	-3	-18	17		
CES juveniles	27	1984-2011	61	59	7	149		
	25	1986-2011	64	92	31	177		
	10	2001-2011	68	7	-10	25		
	5	2006-2011	64	22	-1	41		
BBS UK	16	1995-2011	342	4	-8	20		
	10	2001-2011	381	3	-8	20		
	5	2006-2011	421	3	-4	14		
BBS England	16	1995-2011	252	1	-10	16		
	10	2001-2011	280	7	-4	19		
	5	2006-2011	311	11	3	22		
BBS Scotland	16	1995-2011	37	0	-29	37		
	10	2001-2011	43	25	-24	68		
	5	2006-2011	52	-12	-28	6		
BBS Wales	16	1995-2011	40	3	-30	45		
	10	2001-2011	42	-31	-49	-7	>25	
	5	2006-2011	40	-6	-25	11		


Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




CES adults graph




CES juveniles graph




BBS UK graph



BBS England graph




BBS Scotland graph



BBS Wales graph

Fledglings per breeding attempt


Treecreeper 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Treecreeper




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	20	Curvilinear	2.69 fledglings	2.81 fledglings	4.2%		
Clutch size	43	1968-2011	14	None					Small sample
Brood size	43	1968-2011	28	None					Small sample
Nest failure rate at egg stage	43	1968-2011	22	Curvilinear	2.34% nests/day	1.53% nests/day	-34.6%		Small sample
Nest failure rate at chick stage	43	1968-2011	22	None					Small sample
Laying date	43	1968-2011	13	Linear decline	May 7	Apr 25	-12 days		Small sample
Juvenile to Adult ratio (CES)	27	1984-2011	69	Smoothed trend	97 Index value	100 Index value	3%		
Juvenile to Adult ratio (CES)	25	1986-2011	72	Smoothed trend	122 Index value	100 Index value	-18%		
Juvenile to Adult ratio (CES)	10	2001-2011	76	Smoothed trend	136 Index value	100 Index value	-27%	>25	
Juvenile to Adult ratio (CES)	5	2006-2011	72	Smoothed trend	71 Index value	100 Index value	41%		


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend




Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Smoothed long-term trend in ratio of juvenile:adult birds caught - green lines indicate 85% confidence limits

Key facts

Conservation listings:	Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe) (BiE04) UK: green (species level); amber (race <i>indigenus</i> , >20% of European breeders; races <i>hebridensis</i> and <i>zetlandicus</i> , >20% of European breeders, European status); red (races <i>fridiariensis</i> and <i>hirtensis</i> , rare breeders of global importance) (BoCC3) UK Biodiversity Action Plan: priority species (Fair Isle & St Kilda races)
Long-term trend:	UK, England: moderate increase
Population size:	8.6 million territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)


Migrant status:	Resident
Nesting habitat:	Above-ground nester
Primary breeding habitat:	Woodland
Secondary breeding habitat:	
Breeding diet:	Animal
Winter diet:	Animal

Status summary

The Wren's current UK population estimate is the highest for any species and, on the latest figures, one in ten of our breeding birds is a Wren (APEP13). Abundance can vary sharply from year to year, however. Wren numbers in the UK were greatly depleted by the cold winter of 1962/63 (Marchant et al. 1990). Following a rapid recovery up to the mid 1970s, abundance fell again in response to a further series of cold winters only to return to its previous high level. Following recent severe winters, numbers are depleted once more, especially in Scotland and Northern Ireland. The BBS PECBMS 2013a).

CBC/BBS UK 1966–2012


Wren











Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	1080	54	35	76		Small CBC sample
	25	1986-2011	1731	16	7	23		Small CBC sample
	10	2001-2011	2757	-22	-23	-20		
	5	2006-2011	3080	-23	-24	-20		
CBC/BBS England	44	1967-2011	859	63	43	85		Small CBC sample
	25	1986-2011	1366	23	13	29		Small CBC sample
	10	2001-2011	2166	-16	-18	-13		
	5	2006-2011	2449	-14	-15	-12		
CES adults	27	1984-2011	99	18	1	37		
	25	1986-2011	104	22	4	44		
	10	2001-2011	107	-23	-28	-16		
	5	2006-2011	106	-22	-27	-16		
CES juveniles	27	1984-2011	99	19	-9	59		
	25	1986-2011	104	17	-8	54		
	10	2001-2011	108	-20	-29	-9		
	5	2006-2011	106	-14	-24	-3		
BBS UK	16	1995-2011	2425	-4	-8	-1		
	10	2001-2011	2757	-22	-23	-19		
	5	2006-2011	3080	-23	-24	-20		
BBS England	16	1995-2011	1869	-2	-6	0		
	10	2001-2011	2116	-16	-18	-14		
	5	2006-2011	2364	-14	-15	-12		
BBS Scotland	16	1995-2011	225	-6	-14	8		
	10	2001-2011	248	-37	-42	-27	>25	
	5	2006-2011	284	-46	-49	-39	>25	
BBS Wales	16	1995-2011	195	-8	-21	-1		
	10	2001-2011	219	-24	-31	-17		
	5	2006-2011	220	-21	-25	-14		
BBS N.Ireland	16	1995-2011	91	12	-11	48		
	10	2001-2011	108	-34	-37	-23	>25	
	5	2006-2011	111	-35	-37	-26	>25	

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.








Demographic trends


Fledglings per breeding attempt Wren 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Wren




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	96	Curvilinear	2.37 fledglings	2.86 fledglings	20.6%		
Clutch size	43	1968-2011	95	None					
Brood size	43	1968-2011	127	Curvilinear	3.61 chicks	4.21 chicks	16.6%		
Nest failure rate at egg stage	43	1968-2011	139	Linear decline	1.87% nests/day	1.20% nests/day	-35.8%		
Nest failure rate at chick stage	43	1968-2011	96	None					
Laying date	43	1968-2011	87	Linear decline	May 14	May 8	-6 days		


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend




Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend




Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Smoothed long-term trend in ratio of juvenile:adult birds caught - green lines indicate 85% confidence limits



Proportion of adult birds surviving to following year - green bars represent 95% confidence limits

Causes of change

There is good evidence that mortality rates are severely affected by cold winter weather. Thus, a warming climate may have benefited this species, although there is only circumstantial evidence for this.

Change factor	Primary driver	Secondary driver
Demographic	Increased survival	
Ecological	Climate change	

Further information on causes of change

There has been a reduction in the failure rate of nests at the egg stage, reflected in larger brood sizes and an increase in fledglings per breeding attempt, but the effects of productivity are overshadowed by the strong influence of winter weather on this species.

There is good evidence that annual numbers are influenced by mortality rates and that mortality may be very high in severe winters (Peach et al. 1995). Wren survival rates were negatively correlated with the number of snow days in winter (Peach et al. 1995). Robinson et al. (2007b) showed that survival is related to the strength of the North Atlantic Oscillation, an ocean-scale weather pattern that has a strong influence on UK weather. First-year survival was more influenced by weather than that of adult birds, although adult survival was also affected. These observations suggest that a warming climate may benefit this species.

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts

Conservation listings:	Europe: SPEC category 3 (declining) (BiE04) UK: red (species level, race <i>vulgaris</i>); amber (race <i>zetlandicus</i> , >20% of European breeders) (BoCC3) UK Biodiversity Action Plan: priority species
Long-term trend:	England: rapid decline
Population size:	1,900,000 (1,700,000-2,200,000) pairs in 2009 (APEP13: distance sampling estimate for 2006 (Newson et al. 2008) updated using BBS trend)


Migrant status:	Resident
Nesting habitat:	Cavity nester
Primary breeding habitat:	Farmland
Secondary breeding habitat:	
Breeding diet:	Animal
Winter diet:	Vegetation

Status summary

The abundance of breeding Starlings in the UK has fallen rapidly, particularly since the early 1980s and especially in woodland (Robinson et al. 2002, 2005a), and continues to be strongly downward. The BBS BirdLife International 2004). Overall, there has been widespread moderate decrease across Europe since 1980 (PECBMS 2013a).

CBC/BBS England 1966–2012

Starling




Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	44	1967-2011	626	-88	-92	-84	>50	
	25	1986-2011	1013	-80	-84	-77	>50	
	10	2001-2011	1539	-46	-50	-43	>25	
	5	2006-2011	1669	-32	-36	-29	>25	
BBS UK	16	1995-2011	1745	-53	-56	-50	>50	
	10	2001-2011	1866	-46	-50	-42	>25	
	5	2006-2011	1996	-33	-39	-27	>25	
BBS England	16	1995-2011	1429	-58	-61	-55	>50	
	10	2001-2011	1520	-47	-50	-43	>25	
	5	2006-2011	1638	-33	-36	-29	>25	
BBS Scotland	16	1995-2011	146	-40	-53	-27	>25	
	10	2001-2011	158	-44	-54	-34	>25	
	5	2006-2011	171	-35	-50	-16	>25	
BBS Wales	16	1995-2011	80	-70	-80	-60	>50	
	10	2001-2011	81	-52	-61	-43	>50	
	5	2006-2011	77	-40	-51	-29	>25	
BBS N.Ireland	16	1995-2011	79	25	-6	59		
	10	2001-2011	94	-41	-51	-31	>25	
	5	2006-2011	98	-21	-39	-6		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




CBC/BBS England 1966–2012
Starling




CBC/BBS England graph

BBS UK 1994–2012
Starling




BBS UK graph


BBS England 1994–2012
Starling




BBS England graph



BBS Scotland graph



BBS Wales graph




BBS N.Ireland graph

Demographic trends

Fledglings per breeding attempt


Starling 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Starling




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	120	Linear increase	2.60 fledglings	3.32 fledglings	27.6%		
Clutch size	43	1968-2011	77	Linear increase	4.44 eggs	4.97 eggs	11.8%		
Brood size	43	1968-2011	247	None					
Nest failure rate at egg stage	43	1968-2011	125	Linear decline	1.12% nests/day	0.23% nests/day	-79.5%		
Nest failure rate at chick stage	43	1968-2011	146	Linear decline	0.59% nests/day	0.16% nests/day	-72.9%		
Laying date	43	1968-2011	86	Curvilinear	Apr 27	Apr 20	-7 days		


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

Causes of change

There is good evidence that changes in first-year overwinter survival rates best account for observed population change. Although the ecological drivers of Starling decline are poorly understood, changes in the management of pastoral farmland are thought to be largely responsible.

Change factor	Primary driver	Secondary driver
Demographic	Decreased juvenile survival	
Ecological	Unknown	

Further information on causes of change

As the population has dropped, the numbers of fledglings per breeding attempt has increased markedly (see above); clutches are now larger, and rates of nest loss at the egg and chick stage have fallen. These improvements in breeding performance suggest that decreasing survival rates are likely to be responsible for the decline. Evidence for this is provided by Freeman et al. (2007b), who conducted a population modelling exercise and found that changes in first-year overwinter survival rates could best account for observed population change, and were sufficient, on their own, to explain the broad pattern of decline. The decline in survival rates nationwide coincided with the major period of population decline. MacLeod et al. (2008) also provide evidence linking Starling declines to the environmental conditions outside the breeding season, suggesting that the species' population status is dependent on interactive or synergistic effects of food availability and predation.

There is little direct evidence from studies analysing the ecological drivers of the declines. However, changes in pastoral farming practices are likely to account for at least some of the decline in the wider countryside, probably related to changes in food resources, though these are largely unquantified (Robinson et al. 2005). Loss of permanent pasture, which is the species' preferred feeding habitat, and general intensification of livestock rearing are likely to be having adverse effects on rural populations, but other causes should be sought in urban areas (Robinson et al. 2002, 2005). Whilst the number of cattle has declined, sheep numbers have increased, producing a different sward structure (Chamberlain et al. 2000b, Fuller & Gough 1999) and patterns of stock rearing have changed. These may have reduced foraging opportunities for Starling (Robinson et al. 2002, 2005). Also the use of insecticides on grassland, though low, is targeted partly at tipulids, which may have reduced foraging opportunities further (Vickery et al. 2001). Although there is little published evidence that the density of tipulids has changed over time (Wilson et al. 1999), the area of permanent pasture has declined and the use of insecticides on them has increased. Drainage of grasslands is also thought to have reduced the quality of foraging conditions (Newton 2004). Even after considerable decline among farmland Starlings, tipulids remain important to them for provisioning young (Rhymer et al. 2012).

Further research into urban Starling population dynamics is to be encouraged if we are to understand the causes of decline of this charismatic species more fully.


Key facts

Conservation listings:	Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe) (BiE04) UK: green (species level); amber (race gularis, >20% of European breeders; race hibernicus, >20% of European breeders, European status) (BoCC3)
Long-term trend:	UK waterways: moderate decline
Population size:	6,200-18,700 pairs in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)

Status summary

The WBS/WBBS shows that Dipper populations have fluctuated over the last thirty years, but with an overall downward trend. The species is unusually sensitive to acidity and other water-borne pollution (Ormerod & Tyler 1989, 1990), with lower breeding densities and productivity on acidic than on more neutral streams (Ormerod et al. 1991, Vickery 1991, 1992). Breeding performance has improved strongly over time, and laying dates have shifted earlier, perhaps because of climate change (Crick & Sparks 1999). Broods now average larger than in the late 1960s and 1970s, and there has been substantial reduction in failure rates of nests at the egg stage, leading to sustained increase in the number of fledglings per breeding attempt. In a river system in southern Norway, climate variables including winter temperature explained 84% of the variation in population level during 1978-2008 (Nilsson et al. 2011).

WBS/WBBS 1974-2012
Dipper





Smoothed population index, relative to an arbitrary 100 in the year given, with 85% confidence limits in green

Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
WBS/WBBS waterways	36	1975-2011	64	-30	-47	-8	>25	
	25	1986-2011	76	-26	-40	-7	>25	
	10	2001-2011	113	-18	-28	-2		
	5	2006-2011	106	-6	-16	2		

BBS UK Source	Period (yrs)	1995-2011 Years	Stats (n)	Change (%)	Lower limit	Upper limit	>25 Alert	Comment
	10	2001-2011	64	-39	-57	-18	>25	
	5	2006-2011	74	-29	-45	-5	>25	


Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.



Demographic trends

Fledglings per breeding attempt


Dipper 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Dipper




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	82	Curvilinear	2.05 fledglings	3.06 fledglings	49.5%		
Clutch size	43	1968-2011	78	None					
Brood size	43	1968-2011	150	Curvilinear	3.45 chicks	3.77 chicks	9.3%		
Nest failure rate at egg stage	43	1968-2011	108	Curvilinear	2.80% nests/day	0.36% nests/day	-87.1%		
Nest failure rate at chick stage	43	1968-2011	82	Curvilinear	0.66% nests/day	0.54% nests/day	-18.2%		
Laying date	43	1968-2011	67	Linear decline	Apr 19	Apr 7	-12 days		


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend




Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of adult birds surviving to following year - error bars represent 95% confidence limits

Ring Ouzel

Turdus torquatus

Key facts

Conservation listings:	Europe: no SPEC category (concentrated in Europe, conservation status favourable) (BiE04) UK: red (>50% population decline) (BoCC3) UK Biodiversity Action Plan: priority species
Long-term trend:	UK: decline
Population size:	6,200-7,500 pairs in 1999 (APEP13: Wotton et al. 2002a)

Status summary

The first two breeding atlases showed a decline of 27% in the number of 10-km squares occupied between 1968-72 and 1988-91 (Gibbons et al. 1993), and the extent of population decline was later established by a special survey: a 58% population decline was estimated for the period between 1988-91 and 1999, warranting red listing for this species (Gregory et al. 2002). By 2008-11, the number of occupied 10-km squares had fallen by 43% since 1968-72 (Balmeret al. 2013). Long-term surveys coordinated by the Sim et al. 2010). British & Irish bird observatory data show a decline in spring passage Ring Ouzels at western locations during 1970-98 that matches the estimated UK breeding decline, but no decline at eastern observatories where most birds are of Fennoscandian origin (Burfield & Brooke 2005). These authors infer that, since these populations winter together, the reasons for decline among UK breeders must lie on the breeding grounds or on passage: they also point out that UK birds are more exposed to hunting pressures, particularly in southwest France. It has proved difficult to establish any reasons for decline that are linked to the breeding grounds (Buchanan et al. 2003). In southeast Scotland, however, the breeding sites that are still occupied tend to be those at higher altitude and that have retained an extensive cover of heather (Sim et al. 2007b). In the same study, it was shown that declines were greatest in years following warm summers on the breeding grounds and also greater two years after high spring rainfall in Morocco: these results suggest that the population decline could be linked to reduced food supplies, and consequently higher rates of natural mortality, in autumn and winter (Beale et al. 2006). Large areas of apparently suitable juniper scrub, with abundant berries but no wintering Ring Ouzels, exist in the Atlas Mountains, however (Green et al. 2012). Low survival between breeding seasons is apparently a major national cause of decline (Simet al. 2010). Within Glen Clunie, however, Sim et al. (2011) found that varying combinations of demographic factors produced each year-to-year decline.


Population changes in detail

Annual breeding population changes for this species are not currently monitored by BTO

Demographic trends

Fledglings per breeding attempt


Ring Ouzel 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Ring Ouzel




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	11	None					
Brood size	43	1968-2011	22	None					Small sample
Nest failure rate at egg stage	43	1968-2011	11	None					Small sample
Nest failure rate at chick stage	43	1968-2011	15	None					Small sample
Laying date	43	1968-2011	22	Linear decline	May 15	May 7	-8 days		Small sample


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

Key facts


Conservation listings:	Europe: no SPEC category (concentrated in Europe, conservation status favourable) (BiE04) UK: green (BoCC3)
Long-term trend:	UK, England: shallow decline
Population size:	5.1 (4.9-5.3) million pairs in 2009 (APEP13: distance sampling estimate for 2006 (Newson et al. 2008) updated using BBS trend)

Status summary

Both CBC/BBS and CES data show long-term declines in Blackbird abundance up to about the mid 1990s followed by a strong but partial recovery, which currently has stalled. The BBS Siriwardena et al. 1998a), although there has been little overall change in survival as recorded by CES since 1983. Annual population changes correlate best with adult survival, but population processes appear to differ between eastern and western Britain (Robinson et al. 2012). Fledgling numbers per breeding attempt increased during the population decline and are now decreasing again. Agricultural intensification is likely to have contributed to the population decline (Fuller et al. 1995) but, since numbers fell in woodland as well as farmland, additional factors probably operated. There has been widespread moderate increase across Europe since 1980 (PECBMS 2013a).

CBC/BBS UK 1966–2012

Blackbird





Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	1093	-16	-23	-9		
	25	1986-2011	1748	9	3	15		
	10	2001-2011	2780	4	2	6		
	5	2006-2011	3115	-1	-3	1		

CBC/BBS England Source	Period (yrs)	1967-2011 Years	Obs (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
	25	1986-2011	1405	6	0	12		
	10	2001-2011	2218	3	1	5		
	5	2006-2011	2512	-1	-3	0		
CES adults	27	1984-2011	99	-15	-26	-3		
	25	1986-2011	104	-13	-24	-3		
	10	2001-2011	106	-13	-18	-4		
	5	2006-2011	104	-1	-7	6		
CES juveniles	27	1984-2011	90	-43	-59	-8	>25	
	25	1986-2011	94	-19	-39	11		
	10	2001-2011	96	-15	-29	6		
	5	2006-2011	93	-28	-39	-10	>25	
BBS UK	16	1995-2011	2449	22	19	26		
	10	2001-2011	2780	4	2	6		
	5	2006-2011	3115	-1	-2	0		
BBS England	16	1995-2011	1937	19	15	23		
	10	2001-2011	2185	3	1	5		
	5	2006-2011	2457	-1	-3	0		
BBS Scotland	16	1995-2011	197	30	14	51		
	10	2001-2011	224	12	1	25		
	5	2006-2011	258	2	-5	10		
BBS Wales	16	1995-2011	196	42	30	56		
	10	2001-2011	222	18	10	27		
	5	2006-2011	225	1	-4	7		
BBS N.Ireland	16	1995-2011	86	30	0	50		
	10	2001-2011	101	-21	-28	-14		
	5	2006-2011	103	-11	-17	-6		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




CES adult abundance 1983-2012
Blackbird




CES adults graph

CES juvenile abundance 1983-2012
Blackbird




CES juveniles graph

BBS UK 1994-2012
Blackbird




BBS UK graph

BBS England 1994-2012
Blackbird




BBS England graph

BBS Scotland 1994-2012
Blackbird




BBS Scotland graph

BBS Wales 1994-2012
Blackbird




BBS Wales graph



Demographic trends


Fledglings per breeding attempt Blackbird 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Blackbird




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	233	Curvilinear	1.47 fledglings	1.54 fledglings	4.4%		
Clutch size	43	1968-2011	195	None					
Brood size	43	1968-2011	255	Curvilinear	3.35 chicks	3.28 chicks	-2.0%		
Nest failure rate at egg stage	43	1968-2011	285	Curvilinear	2.60% nests/day	3.78% nests/day	45.4%		
Nest failure rate at chick stage	43	1968-2011	233	Linear decline	2.87% nests/day	1.89% nests/day	-34.1%		
Laying date	43	1968-2011	231	Curvilinear	Apr 23	Apr 25	2 days		
Juvenile to Adult ratio (CES)	27	1984-2011	101	Smoothed trend	159 Index value	100 Index value	-37%	>25	
Juvenile to Adult ratio (CES)	25	1986-2011	106	Smoothed trend	145 Index value	100 Index value	-31%	>25	
Juvenile to Adult ratio (CES)	10	2001-2011	108	Smoothed trend	109 Index value	100 Index value	-8%		
Juvenile to Adult ratio (CES)	5	2006-2011	106	Smoothed trend	135 Index value	100 Index value	-26%	>25	


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend




Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend




Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Smoothed long-term trend in ratio of juvenile:adult birds caught - green lines indicate 85% confidence limits



Proportion of adult birds surviving to following year - green bars represent 95% confidence limits

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts


Conservation listings:	Europe: no SPEC category (concentrated in Europe, conservation status favourable) (BiE04) UK: red (species level, races <i>clarkei</i> and <i>hebridensis</i>) (BoCC3) UK Biodiversity Action Plan: click here , priority species (clarkei and hebridensis)
Long-term trend:	UK, England: rapid decline
Population size:	1.2 million territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)
Migrant status:	Short-distance migrant
Nesting habitat:	Above-ground nester
Primary breeding habitat:	Woodland
Secondary breeding habitat:	
Breeding diet:	Animal
Winter diet:	Animal

Status summary

CBC/BBS shows a rapid decline in Song Thrush abundance that began in the mid 1970s. The latter part of this decline can also be seen in the CES index. BBS data from all UK countries show increase from 1994 to 2008, followed by a sharp downturn, but population levels remained relatively low throughout. The BBS PECBMS 2013a).

CBC/BBS UK 1966–2012

Song Thrush




Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	895	-57	-63	-49	>50	
	25	1986-2011	1410	-4	-14	7		
	10	2001-2011	2277	-9	-13	-6		
	5	2006-2011	2555	-14	-16	-11		
CBC/BBS England	44	1967-2011	711	-55	-62	-44	>50	
	25	1986-2011	1110	0	-12	13		
	10	2001-2011	1785	0	-3	4		
	5	2006-2011	2025	-8	-10	-5		
CES adults	27	1984-2011	82	-33	-47	-17	>25	
	25	1986-2011	86	-23	-38	-7		
	10	2001-2011	90	-12	-24	1		
	5	2006-2011	87	-17	-26	-6		
CES juveniles	27	1984-2011	69	-54	-67	-32	>50	
	25	1986-2011	72	-23	-42	9		
	10	2001-2011	76	-12	-26	5		
	5	2006-2011	73	-19	-33	-4		
BBS UK	16	1995-2011	1969	7	3	14		
	10	2001-2011	2277	-9	-12	-5		
	5	2006-2011	2555	-14	-16	-11		
BBS England	16	1995-2011	1523	12	7	16		
	10	2001-2011	1759	0	-3	3		
	5	2006-2011	1982	-8	-10	-5		
BBS Scotland	16	1995-2011	176	-5	-19	11		
	10	2001-2011	199	-26	-37	-12	>25	
	5	2006-2011	229	-25	-32	-16	>25	
BBS Wales	16	1995-2011	167	10	-1	24		
	10	2001-2011	192	-17	-24	-9		
	5	2006-2011	196	-20	-23	-14		
BBS N.Ireland	16	1995-2011	77	23	-6	57		
	10	2001-2011	92	-22	-31	-6		
	5	2006-2011	95	-24	-30	-11		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




CBC/BBS UK 1966–2012
Song Thrush




CBC/BBS UK graph

CBC/BBS England 1966–2012
Song Thrush




CES adult abundance 1983–2012
Song Thrush




CES adults graph

CES juvenile abundance 1983–2012
Song Thrush




CES juveniles graph

BBS UK 1994–2012
Song Thrush




BBS UK graph

BBS England 1994–2012
Song Thrush




BBS England graph

BBS Scotland 1994–2012
Song Thrush




BBS Scotland graph

BBS Wales 1994–2012
Song Thrush




BBS Wales graph



BBS N.Ireland graph

Demographic trends


Fledglings per breeding attempt Song Thrush 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Song Thrush




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	30	1981-2011	227	Curvilinear	1.23 fledglings	1.28 fledglings	3.6%		
Clutch size	43	1968-2011	172	None					
Brood size	43	1968-2011	190	None					
Nest failure rate at egg stage	30	1981-2011	309	Curvilinear	4.40% nests/day	4.65% nests/day	5.7%		
Nest failure rate at chick stage	30	1981-2011	227	Curvilinear	2.56% nests/day	2.21% nests/day	-13.7%		
Laying date	43	1968-2011	197	None			0 days		
Juvenile to Adult ratio (CES)	27	1984-2011	90	Smoothed trend	143 Index value	100 Index value	-30%	>25	
Juvenile to Adult ratio (CES)	25	1986-2011	94	Smoothed trend	104 Index value	100 Index value	-4%		
Juvenile to Adult ratio (CES)	10	2001-2011	99	Smoothed trend	96 Index value	100 Index value	4%		
Juvenile to Adult ratio (CES)	5	2006-2011	96	Smoothed trend	92 Index value	100 Index value	9%		


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend




Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend




Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Smoothed long-term trend in ratio of juvenile:adult birds caught - green lines indicate 85% confidence limits



Proportion of adult birds surviving to following year - green bars represent 95% confidence limits

Causes of change

Changes in survival in the first winter, and perhaps also the post-fledging period, are sufficient to have caused the population decline. The environmental causes of this are unknown but are likely to include changes in farming practices, particularly land drainage and possibly increased pesticide usage.

Change factor	Primary driver	Secondary driver
Demographic	Decreased juvenile survival	
Ecological	Unknown	

Further information on causes of change

CES productivity shows an initial decrease, followed by some partial recovery, and the number of fledglings per breeding attempt increased during the 1980s and 1990s (see above). There is good evidence to show that changes in survival in the first winter, and perhaps also the post-fledging period, are sufficient to have caused the population decline (Thomson et al. 1997, Siriwardena et al. 1998, Robinson et al. 2004).

Peach et al. (2004) suggested that loss of hedgerows, scrub and permanent grassland with livestock and the widespread installation of field drainage systems, all of which would act to reduce the availability of good quality foraging areas, have probably contributed to the decline of the Song Thrush in the UK. Similarly, it has been suggested that the species is unable to survive the winter in woodland, due to a lack of food, and a reduction of food supply in other habitat types has also been reported (Simms 1989). It is likely that a reduction in food supply would adversely affect the survival of juvenile birds to a greater extent than adult birds, as appears to be the case (Robinson et al. 2004). Furthermore survival is reduced during periods of long drought or cold weather when food is likely to be less available (Robinsoret al. 2007).

In woodland, drainage of damp ground and the depletion of woodland shrub layers through canopy closure and deer browsing may also be implicated (Fuller et al. 2005). There is also some concern of the impact of overgrazing by deer (e.g. Gill & Beardall 2001) and canopy closure (Mason 2007), due to changes in woodland management (Hopkins & Kirby 2007) on the low woodland layers, although good evidence from the UK is sparse (but there are some experimental studies in America on different species which demonstrate this effect, e.g. McShea & Rappole 2000). Several papers (e.g. Gosler 1990, Perrins & Overall 2001, Perrins 2003) state that the understorey has declined in Britain, but few data are available to support this on a national scale. However, Amar et al. (2006) found a 27% increase in understorey in the RSPB sites used in the Repeat Woodland Bird Survey.

Robinson et al. (2004) suggested that predation was a candidate cause of reduced survival but there is conflicting evidence on the role of predators in Song Thrush decline, and further research is needed. Newson et al. (2010b) found no evidence of effects of avian predators or grey squirrels on Song Thrushes.


This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts

Conservation listings:	Europe: no SPEC category (concentrated in Europe, conservation status favourable) (BiE04) UK: amber (25-50% population decline) (BoCC3)
Long-term trend:	UK, England: rapid decline
Population size:	170,000 territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)

Status summary

Like those of Siriwardena et al. 1998). Numbers have shown moderate decline across Europe since 1980, though with little change since 1990 (PECBMS 2013a).







Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	552	-59	-65	-51	>50	
	25	1986-2011	855	-44	-50	-36	>25	
	10	2001-2011	1285	-36	-39	-31	>25	
	5	2006-2011	1363	-26	-30	-22	>25	
CBC/BBS England	44	1967-2011	449	-61	-68	-54	>50	
	25	1986-2011	689	-49	-55	-42	>25	
	10	2001-2011	1015	-38	-42	-34	>25	


Source	Period (yrs)	2006-2011 Years	1083 Plots (n)	-25 Change (%)	-29 Lower limit	-22 Upper limit	Alert	Comment
BBS UK	16	1995-2011	1176	-31	-37	-24	>25	
	10	2001-2011	1285	-36	-40	-32	>25	
	5	2006-2011	1363	-26	-30	-22	>25	
BBS England	16	1995-2011	922	-38	-42	-33	>25	
	10	2001-2011	992	-38	-41	-35	>25	
	5	2006-2011	1044	-25	-28	-22	>25	
BBS Scotland	16	1995-2011	77	-12	-33	31		
	10	2001-2011	87	-42	-54	-20	>25	
	5	2006-2011	99	-37	-49	-20	>25	
BBS Wales	16	1995-2011	100	-4	-25	16		
	10	2001-2011	111	-11	-25	4		
	5	2006-2011	110	-7	-20	7		
BBS N.Ireland	16	1995-2011	59	-16	-61	50		
	10	2001-2011	70	-38	-52	-20	>25	
	5	2006-2011	70	-33	-43	-25	>25	

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.







BBS England graph



BBS Scotland graph



BBS Wales graph




BBS N.Ireland graph

Demographic trends

Fledglings per breeding attempt


Mistle Thrush 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Mistle Thrush




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	51	None					
Clutch size	43	1968-2011	32	Linear increase	3.89 eggs	4.04 eggs	3.8%		
Brood size	43	1968-2011	65	None					
Nest failure rate at egg stage	43	1968-2011	53	None					
Nest failure rate at chick stage	43	1968-2011	57	None					
Laying date	43	1968-2011	27	None			0 days		Small sample


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

Spotted Flycatcher

Muscicapa striata

Key facts


Conservation listings:	Europe: SPEC category 3 (declining) (BiE04) UK: red (>50% population decline) (BoCC3) UK Biodiversity Action Plan: click here , priority species
Long-term trend:	UK, England: rapid decline
Population size:	36,000 territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)
Migrant status:	Long-distance migrant
Nesting habitat:	Above-ground nester
Primary breeding habitat:	Woodland
Secondary breeding habitat:	
Breeding diet:	Animal
Winter diet:	Animal

Status summary

Spotted Flycatchers have declined rapidly and consistently since the 1960s. The Repeat Woodland Bird Survey, however, using a set of CBC woodland and RSPB sites, detected a significant increase between the 1980s and 2003-04 in southwest England (Amar et al. 2006, Hewson et al. 2007), suggesting that change has not been uniform across Britain. Gaps are starting to appear in the 10-km distribution map, especially in urban areas and close to the east coast (Balmer et al. 2013). Numbers have shown widespread moderate decrease across Europe since 1980, though with little change since 1990 (PECBMS 2013a).

CBC/BBS UK 1966–2012

Spotted Flycatcher



Smoothed population index, relative to an arbitrary 100 in the year given, with 85% confidence limits in green


Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	127	-89	-93	-85	>50	
	25	1986-2011	160	-81	-86	-75	>50	
	10	2001-2011	186	-35	-49	-21	>25	
	5	2006-2011	185	-23	-39	-12		
CBC/BBS England	44	1967-2011	97	-91	-94	-88	>50	
	25	1986-2011	117	-84	-89	-78	>50	
	10	2001-2011	131	-47	-56	-36	>25	
	5	2006-2011	132	-31	-44	-20	>25	
BBS UK	16	1995-2011	194	-49	-59	-38	>25	
	10	2001-2011	186	-34	-49	-17	>25	
	5	2006-2011	185	-23	-37	-12		
BBS England	16	1995-2011	135	-56	-64	-45	>50	
	10	2001-2011	126	-45	-54	-33	>25	
	5	2006-2011	124	-30	-41	-17	>25	

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




CBC/BBS UK 1966–2012
Spotted Flycatcher




CBC/BBS UK graph

CBC/BBS England 1966–2012
Spotted Flycatcher




CBC/BBS England graph

BBS UK 1994–2012
Spotted Flycatcher




BBS UK graph



Demographic trends


Fledglings per breeding attempt Spotted Flycatcher 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Spotted Flycatcher




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	103	None					
Clutch size	43	1968-2011	78	None					
Brood size	43	1968-2011	128	Curvilinear	3.63 chicks	3.67 chicks	1.1%		
Nest failure rate at egg stage	43	1968-2011	115	Curvilinear	1.79% nests/day	1.56% nests/day	-12.8%		
Nest failure rate at chick stage	43	1968-2011	104	None					
Laying date	43	1968-2011	70	None			0 days		


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

Causes of change

Demographic modelling provides evidence that a decrease in the annual survival rates of birds in their first year may have driven the decline. The ecological causes of the decline are uncertain as good-quality, direct evidence is sparse.

Change factor	Primary driver	Secondary driver
Demographic	Decreased survival	
Ecological	Unknown	

Further information on causes of change

Nest failure rates have decreased but the number of fledglings per breeding attempt shows no trend. Though samples are too small to continue presenting a trend, there was a decrease overall in the ratio of juveniles to adults in CES captures. However, demographic modelling shows that decreases in the annual survival rates of birds in their first year of life are more likely to have driven the population decline than breeding parameters (Freeman & Crick 2003, Stevens et al. 2007). This effect on survival may operate in the pre-migration period, during migration or in the wintering quarters. The number of adult Spotted Flycatchers caught at CES ringing sites was found to have declined drastically, providing further evidence that post-fledging and overwinter survival may be important factors in the population decline (Peach et al. 1998).

Evidence for the ecological causes of the decline is sparse. Fuller et al. (2005) hypothesise that declines in large flying insects that are food to the flycatcher, or conditions either on the wintering grounds or along migration routes may be involved. However, there is little detailed evidence to directly support any of these ideas.

Data from the Repeat Woodland Bird Survey (Amaret al. 2006) showed that Spotted Flycatchers were more likely to have declined at sites with very open or very closed foliage conditions. Smart et al. (2007) also suggest this. However, overall, Amar et al. (2006) did not find that changes in habitat were significant in explaining population declines for this species. Stevens et al. (2007) found that nests in gardens were more successful than those in farmland or woodland and nests in gardens fledged twice as many chicks as those in either woodland or farmland. The proximate cause of lower success in farmland and woodland was higher nest predation rates. In terms of nesting success, farmland and woodland appear to be suboptimal when compared with gardens, providing evidence of a problem on the breeding grounds for this species, at least in these two habitats (Stevens et al. 2007).

In Leicestershire, Stoate & Szczur (2006) found that the removal of nest predators prompted an increase in Spotted Flycatcher breeding success, especially in woodland, where nest success was lower overall than in gardens. However, Carpenter et al. (2009) found no link between presence/absence, abundance and population change of the species and avian predator abundance.

Key facts


Conservation listings:	Europe: no SPEC category (concentrated in Europe, conservation status favourable) (BiE04) UK: green (BoCC3)
Long-term trend:	UK, England: shallow increase
Population size:	6.7 million territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)

Status summary

Robins have increased markedly since the mid 1980s, according to both CBC/BBS and CES results, having been set back earlier by a succession of cold winters. Steep improvements have occurred concurrently in the numbers of fledglings per breeding attempt, as measured by nest record data, with a reduction in nest failure rates at the egg stage, although CES productivity measures have been relatively unchanged. Survival rates, as measured by CES, appear stable. The CES and BBS data show that marked and significant annual fluctuations occur in numbers, perhaps in response to winter weather, although these are not evident in the smoothed trends: numbers have dropped sharply in the last few seasons, but remain relatively high. The BBS PECBMS 2013a).

CBC/BBS UK 1966-2012

Robin





Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	1049	27	14	42		
	25	1986-2011	1677	42	34	53		
	10	2001-2011	2673	-8	-10	-6		
	5	2006-2011	2995	-11	-13	-9		

CBC/BBS England Source	44 Period (yrs)	1967-2011 Years 1986-2011	841 Plots (1937)	42 Change (%)	27 Lower limit	59 Upper limit	Alert	Comment
	10	2001-2011	2119	-2	-5	0		
	5	2006-2011	2398	-7	-8	-5		
CES adults	27	1984-2011	93	14	-3	31		
	25	1986-2011	98	25	9	45		
	10	2001-2011	102	-20	-26	-13		
	5	2006-2011	99	-25	-30	-20	>25	
CES juveniles	27	1984-2011	99	9	-20	51		
	25	1986-2011	103	30	-1	73		
	10	2001-2011	107	-10	-21	2		
	5	2006-2011	106	-11	-20	0		
BBS UK	16	1995-2011	2349	6	3	10		
	10	2001-2011	2673	-8	-10	-6		
	5	2006-2011	2995	-11	-13	-9		
BBS England	16	1995-2011	1841	13	9	16		
	10	2001-2011	2088	-2	-4	1		
	5	2006-2011	2347	-7	-8	-5		
BBS Scotland	16	1995-2011	198	4	-6	13		
	10	2001-2011	221	-11	-19	-1		
	5	2006-2011	258	-18	-24	-11		
BBS Wales	16	1995-2011	192	-16	-22	-6		
	10	2001-2011	216	-26	-31	-19	>25	
	5	2006-2011	216	-25	-29	-19	>25	
BBS N.Ireland	16	1995-2011	88	1	-15	19		
	10	2001-2011	103	-21	-28	-13		
	5	2006-2011	107	-14	-20	-7		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




CES adult abundance 1983-2012




CES adults graph

CES juvenile abundance 1983-2012




CES juveniles graph

BBS UK 1994-2012




BBS UK graph

BBS England 1994-2012




BBS England graph

BBS Scotland 1994-2012




BBS Scotland graph

BBS Wales 1994-2012




BBS Wales graph



Demographic trends


Fledglings per breeding attempt Robin 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Robin




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	207	Linear increase	2.42 fledglings	2.85 fledglings	17.7%		
Clutch size	43	1968-2011	154	None					
Brood size	43	1968-2011	231	Curvilinear	4.40 chicks	4.31 chicks	-2.2%		
Nest failure rate at egg stage	43	1968-2011	227	Curvilinear	2.41% nests/day	1.31% nests/day	-45.6%		
Nest failure rate at chick stage	43	1968-2011	209	None					
Laying date	43	1968-2011	149	Linear decline	Apr 28	Apr 20	-8 days		
Juvenile to Adult ratio (CES)	27	1984-2011	102	Smoothed trend	100 Index value	100 Index value	0%		
Juvenile to Adult ratio (CES)	25	1986-2011	107	Smoothed trend	116 Index value	100 Index value	-14%		
Juvenile to Adult ratio (CES)	10	2001-2011	110	Smoothed trend	96 Index value	100 Index value	4%		
Juvenile to Adult ratio (CES)	5	2006-2011	109	Smoothed trend	93 Index value	100 Index value	7%		


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend




Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend




Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Smoothed long-term trend in ratio of juvenile:adult birds caught - green lines indicate 85% confidence limits



Proportion of adult birds surviving to following year - green bars represent 95% confidence limits

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Nightingale

Luscinia megarhynchos

Key facts

Conservation listings:	Europe: no SPEC category (concentrated in Europe, conservation status favourable) (BiE04) UK: amber (25-50% distribution decline) (BoCC3)
Long-term trend:	England: decline
Population size:	6,700 (5,600-9,400) males in 1999 (APEP13: Wilson et al. 2002)

Status summary

In 1999, the BTO organised a national survey of Nightingales, which showed a marked range contraction since the previous survey in 1980, but only an 8% overall population decline (Wilson et al. 2002; for more details Balmer et al. 2013). Fuller et al. (2005) suggest the likely causes of Nightingale decline relate to pressures on migration and in winter, perhaps compounded by habitat loss in Britain. The increasing intensity of browsing by deer is known to be reducing habitat quality for this species (Gill & Fuller 2007, Holt et al. 2010). Though samples are too small to continue presenting a trend, CES suggested a sharp decline in productivity during the 1980s, perhaps because Nightingale nesting success may be adversely affected by cold and wet springs. Holt et al. (2012b) emphasise the value for Nightingale conservation of promoting habitat quality and researching breeding biology and migration/wintering areas. Woodland-scrub mosaics appear to be important breeding habitats, with implications for conservation practice at such sites (Holt et al. 2012c). Nightingale has been in moderate decline across Europe since 1980, though with little change since 1990 (PECBMS 2013a); this overall trend masks a marked contrast between severe decreases in southern and western Europe and increases in the east of the range (PECBMS 2007).



Smoothed population index, relative to an arbitrary 100 in 2013, with 85% confidence limits in green



Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	16	1995-2011	31	-46	-63	-14	>25	
	10	2001-2011	32	-34	-51	0		
	5	2006-2011	35	-4	-25	45		
BBS England	16	1995-2011	31	-42	-58	0		
	10	2001-2011	32	-32	-47	7		
	5	2006-2011	35	-2	-19	43		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See here for more information.



No CBC/BBS UK trend is available for this species. Smoothed CBC/BBS England trend graph



Demographic trends

Productivity and survival trends for this species are not currently produced by BTO

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglinton, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Pied Flycatcher

Ficedula hypoleuca

Key facts

Conservation listings:	Europe: no SPEC category (concentrated in Europe, conservation status favourable) (BiE04) UK: amber (25-50% decline) (BoCC3)
Long-term trend:	UK: decline
Population size:	17,000-20,000 pairs in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)


Migrant status:	Long-distance migrant
Nesting habitat:	Above-ground nester
Primary breeding habitat:	Woodland
Secondary breeding habitat:	
Breeding diet:	Animal
Winter diet:	Animal

Status summary

Pied Flycatchers are restricted to upland deciduous woods in parts of western and northern Britain. The proportions of CBC plots occupied rose during the 1980s, but the species was never numerous enough for trends to be estimated (Marchant et al. 1990). The 1988-91 breeding atlas revealed a small expansion in range from 1968-72, aided by the provision of nest boxes in new areas (Gibbons et al. 1993). BBS indicates, however, that abundance has decreased steeply since 1994, prompting the species' recent move from the green to the amber list. Nest-box occupancy rates have also fallen over a similar period at a number of sites monitored as RAS projects. Numbers have shown widespread moderate decline across Europe since 1980 (PECBMS 2013a).

BBS UK 1994–2012


Pied Flycatcher



Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	16	1995-2011	40	-52	-67	-35	>50	

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See here for more information.



Demographic trends


Fledglings per breeding attempt Pied Flycatcher 1966-2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Pied Flycatcher




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	356	None					
Clutch size	43	1968-2011	357	Linear increase	6.55 eggs	6.89 eggs	5.3%		
Brood size	43	1968-2011	391	None					
Nest failure rate at egg stage	43	1968-2011	436	Curvilinear	0.57% nests/day	0.26% nests/day	-54.4%		
Nest failure rate at chick stage	43	1968-2011	358	Linear increase	0.38% nests/day	0.66% nests/day	73.7%		
Laying date	43	1968-2011	441	Linear decline	May 21	May 9	-12 days		


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend




Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of adult birds surviving to following year - error bars represent 95% confidence limits

Causes of change

The reasons for this decline are unknown, but there is good evidence that they lie at least partly outside the breeding season and are thought to be linked to changing conditions on wintering grounds and migration.

Change factor	Primary driver	Secondary driver
Demographic	Overwinter survival	
Ecological	Changes on wintering grounds	

Further information on causes of change

The reasons for this decline are unknown, but there is good evidence that they lie at least partly outside the breeding season (Goodenough et al. 2009). No trends are evident in the number of fledglings per breeding attempt (see above). There has been a linear increase in clutch size but although the failure rate at the egg stage has shown a decrease, failure rate at the chick stage has increased.

There is good evidence that declines are related to conditions outside the breeding season. Goodenough et al. (2009) found that decreasing breeding performance is contributing to decline, but that non-breeding factors are more important. Winter NAO index is a strong predictor of breeding population, probably because the North Atlantic oscillation influences food abundance in Africa and at migratory stopover points. Long-term autumn bird monitoring data from Russia were related to monthly mean temperatures on the West African wintering grounds; the positive relationship suggests that increasing bird numbers are explained by increasing mean November

temperatures. Precipitation and European autumn, spring and breeding-range temperatures did not show a strong relationship (Chernetsov & Huettmann 2005). Thingstad et al. (2006) found that weather conditions at the flycatcher's wintering areas in western Africa were suspected to be responsible for the decrease in Scandinavia, although the breeding success of the sink populations was significantly correlated to June temperatures.

In the Netherlands, climate change may have brought about decline in Pied Flycatchers by advancing the peak period of food availability for this species in deciduous forests - the birds being unable to compensate for the change in food supply by breeding earlier (Both 2002, Both et al. 2006). A more recent paper found that timing of spring migration has responded flexibly to climate change as recovery dates during spring migration in North Africa advanced by ten days between 1980 and 2002, which was explained by improving Sahel rainfall and a phenotypic effect of birth date. However, there was no advance in arrival dates on the breeding grounds, most likely due to environmental constraints during migration (Both 2010). Furthermore, declines were found to be stronger in forests, as these were more seasonal habitats whereas less seasonal marshes showed less steep declines (Both et al. 2009).

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts


Conservation listings:	Europe: SPEC category 2 (depleted) (BiE04) UK: amber (European status) (BoCC3)
Long-term trend:	UK, England: fluctuating, with no long-term trend
Population size:	100,000 (70,000-130,000) pairs in 2009 (APEP13: distance sampling estimate for 2006 (Newson et al. 2008) updated using BBS trend)

Status summary

A sharp decline in the late 1960s and early 1970s was thought to be due to severe drought conditions in the Sahel wintering area in Africa (Marchant et al. 1990). There was a loss of range of 20% in Britain between 1968-72 and 1988-91, in terms of the numbers of occupied 10-km squares (Gibbons et al. 1993). A recovery in population size began in the mid 1970s and appears to have been sustained subsequently, although with some setbacks. This increase has been associated with steeply improving numbers of fledglings per breeding attempt and progressively earlier laying dates. The trend towards earlier laying can be partly explained by recent climate change (Crick & Sparks 1999). Range, meanwhile, has contracted further, especially in the lowlands (Balmer et al. 2013). There has been widespread moderate increase across Europe since 1980 (PECBMS 2013a).

CBC/BBS UK 1966–2012

Redstart



Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	78	6	-29	62		
	25	1986-2011	120	45	23	76		
	10	2001-2011	171	26	13	44		
	5	2006-2011	194	35	21	55		

CBC/BBS England Source	Period (yrs)	1967-2011 Years	#Bots (n)	Change (%)	Lower limit	Upper limit	Alert	Small CBC sample Comment
	25	1986-2011	71	26	0	65		
	10	2001-2011	99	9	-4	22		
	5	2006-2011	117	23	9	39		
BBS UK	16	1995-2011	160	31	14	49		
	10	2001-2011	171	26	13	40		
	5	2006-2011	194	35	24	55		
BBS England	16	1995-2011	90	12	-10	32		
	10	2001-2011	99	10	-5	28		
	5	2006-2011	117	23	9	41		
BBS Wales	16	1995-2011	56	39	16	66		
	10	2001-2011	59	38	14	67		
	5	2006-2011	62	48	30	83		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




CBC/BBS UK 1966-2012
Redstart




CBC/BBS UK graph

CBC/BBS England 1966-2012
Redstart




CBC/BBS England graph

BBS UK 1994-2012
Redstart




BBS UK graph

BBS England 1994-2012
Redstart




BBS England graph



Demographic trends


Fledglings per breeding attempt Redstart 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Redstart




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	54	Curvilinear	3.52 fledglings	5.01 fledglings	42.4%		
Clutch size	43	1968-2011	50	Curvilinear	5.91 eggs	6.24 eggs	5.6%		
Brood size	43	1968-2011	90	Curvilinear	5.15 chicks	5.62 chicks	9.1%		
Nest failure rate at egg stage	43	1968-2011	76	Curvilinear	1.36% nests/day	0.52% nests/day	-61.8%		
Nest failure rate at chick stage	43	1968-2011	54	Linear decline	1.20% nests/day	0.34% nests/day	-71.7%		
Laying date	43	1968-2011	64	Curvilinear	May 21	May 7	-14 days		


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

Key facts

Conservation listings:	Europe: no SPEC category (concentrated in Europe, conservation status favourable) (BiE04) UK: amber (25-50% decline) (BoCC3)
Long-term trend:	UK: decline
Population size:	47,000 (19,000-75,000) pairs in 2009 (APEP13: distance sampling estimate for 2006 (Newson et al. 2008) updated using BBS trend)


Status summary

Whinchats were not monitored until the BBS began in 1994. By then, however, Gibbons et al. (1993) had already identified a major range contraction, mainly from lowland England, that was probably at least partly due to the loss of marginal farmland habitats (Marchant et al. 1990). Further extinctions have occurred since then among the remaining pockets of lowland breeders (Balmer et al. 2013). In the uplands, Whinchat habitat is now somewhat restricted, being sandwiched between intensive agriculture at lower levels and higher land unsuitable for breeding, and limited also by aspect (Calladine & Bray 2012). In a study focused on upland grasslands, a 95% decline was noted between 1968-80 and 1999-2000 (Henderson et al. 2004). BBS data indicate that strong population decline has taken place since 1994, raising BTO alerts for the UK as a whole as well as for England. Nest-record samples are small, but indicate a substantial recent rise in nest losses at the egg stage, which is of NRS concern (Leech & Barimore 2008). Whinchats have shown moderate decline across Europe since 1980, though with little change since 1990 (PECBMS 2013a). On the strength of its UK decline, Whinchat was moved from the green to the amber list of conservation concern in 2009 (Eaton et al. 2009).

In 2012, BTO conducted a [Wales Chat Survey](#) for Whinchats, Stonechats and Wheatears on over 300 sample Welsh 1-km squares. The survey will estimate breeding numbers and distribution and record habitat choice by territorial males.

BBS UK 1994–2012

Whinchat




Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	16	1995-2011	75	-60	-71	-48	>50	

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
	5	2006-2011	76	-34	-47	-18	>25	
BBS England	16	1995-2011	32	-48	-66	-27	>25	
	10	2001-2011	32	-37	-53	-21	>25	
	5	2006-2011	37	-15	-31	1		


Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.



Demographic trends

Fledglings per breeding attempt


Whinchat 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Whinchat




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	15	None					
Clutch size	43	1968-2011	13	None					Small sample
Brood size	43	1968-2011	39	None					
Nest failure rate at egg stage	43	1968-2011	16	Linear increase	0.65% nests/day	2.47% nests/day	280.0%		Small sample
Nest failure rate at chick stage	43	1968-2011	27	None					Small sample
Laying date	43	1968-2011	29	Linear decline	May 30	May 25	-5 days		Small sample


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

Key facts


Conservation listings:	Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe) (BiE04) UK: green (BoCC3)
Long-term trend:	UK: probably fluctuating, with no long-term trend
Population size:	59,000 (39,000-79,000) pairs in 2009 (APEP13: distance sampling estimate for 2006 (Newson et al. 2008) updated using BBS trend)

Status summary

Trends were poorly quantified before the start of the BBS, but a long-term decline is suspected in that period: severe winter weather, and loss and fragmentation of suitable breeding habitat in many inland regions, are believed to have reduced the population from the 1940s onward (Marchant et al. 1990). Breeding atlas data showed a substantial contraction in the Stonechat's range between 1968-72 and 1988-91 (Gibbons et al. 1993). Against this background, the strongly increasing BBS trend to 2006 represents substantial and possibly even complete recovery. By 2008-11, the earlier range losses had been almost entirely reversed (Balmer et al. 2013). Nest failure rates have fallen markedly over the long term. Following increases widely across Europe, the species is now provisionally categorised as 'secure' (BirdLife International 2004) and consequently has recently been moved from the amber to the green list in the UK (Eaton et al. 2009). UK data now indicate a sharp decrease, however, partly in response to recent snowy winters, and an accompanying decrease in nest productivity.

In 2012, BTO conducted a Wales Chat Survey for Whinchats, Stonechats and Wheatears on over 300 sample Welsh 1-km squares. The survey will estimate breeding numbers and distribution and record habitat choice by territorial males.

BBS UK 1994-2012
Stonechat




Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	16	1995-2011	157	3	-15	31		

Source	10 Period (yrs)	2001-2011 Years 2006-2011	203 Plots 231	-49 Change (%)	-56 Lower limit	-37 Upper limit	>25 Alert >50	Comment
BBS England	16	1995-2011	71	7	-31	68		
	10	2001-2011	95	-44	-56	-30	>25	
	5	2006-2011	117	-56	-64	-44	>50	
BBS Scotland	16	1995-2011	38	-20	-43	31		
	10	2001-2011	48	-56	-65	-36	>50	
	5	2006-2011	56	-73	-78	-58	>50	
BBS Wales	16	1995-2011	36	82	18	179		
	10	2001-2011	45	-20	-43	4		
	5	2006-2011	44	-36	-50	-24	>25	


Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




BBS UK graph



BBS England graph




BBS Scotland graph



BBS Wales graph

Fledglings per breeding attempt


Stonechat 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Stonechat




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	38	None					
Clutch size	43	1968-2011	35	None					
Brood size	43	1968-2011	68	None					
Nest failure rate at egg stage	43	1968-2011	39	None					
Nest failure rate at chick stage	43	1968-2011	62	Curvilinear	1.57% nests/day	1.25% nests/day	-20.4%		
Laying date	43	1968-2011	42	Curvilinear	May 3	Apr 23	-10 days		


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend




Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of adult birds surviving to following year - error bars represent 95% confidence limits

Key facts


Conservation listings:	Europe: SPEC category 3 (declining) (BiE04) UK: amber (species level and nominate race oenanthe, European status) (BoCC3)
Long-term trend:	UK: possible decline
Population size:	240,000 (170,000-310,000) pairs in 2009 (APEP13: distance sampling estimate for 2006 (Newson et al. 2008) updated using BBS trend)

Status summary

Although it is a common breeding species in many upland areas, the Wheatear was not monitored at the UK scale until the BBS began in 1994. Gibbons et al. (1993) had by then identified range contractions from lowland Britain since 1968-72, perhaps due to losses of suitable grassland and declines in rabbit abundance. Further loss of range, especially in lowland England, had been recorded by 2008-11 (Balmer et al. 2013). BBS trends show wide fluctuations, with little indication of directional change. Previous estimates of UK population have been revised strongly upward, based on BBS distance-sampling data (Gillings et al. 2007). Nest failure rates at the egg stage have fallen substantially and nest productivity has risen. Wheatear has shown moderate decline across Europe since 1980 (PECBMS 2013a). Following widespread declines during the 1990s, the European status of this species is no longer considered 'secure' (BirdLife International 2004). Accordingly, the species was moved from the green to the amber list in the UK in 2009 (Eaton et al. 2009).

In 2012, BTO conducted a Wales Chat Survey for Whinchats, Stonechats and Wheatears on over 300 sample Welsh 1-km squares. The survey will estimate breeding numbers and distribution and record habitat choice by territorial males.

BBS UK 1994-2012
Wheatear




Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	16	1995-2011	325	-1	-17	17		

Source	10 Period (yrs)	2001-2011 Years	371 Plots	19 Change (%)	6 Lower Limit	35 Upper Limit	Alert	Comment
		2006-2011	432					
BBS England	16	1995-2011	181	17	-7	41		
	10	2001-2011	219	29	5	56		
	5	2006-2011	271	36	21	49		
BBS Scotland	16	1995-2011	79	-4	-24	24		
	10	2001-2011	79	12	-9	33		
	5	2006-2011	86	7	-9	24		
BBS Wales	16	1995-2011	52	-13	-29	7		
	10	2001-2011	57	9	-10	33		
	5	2006-2011	57	11	-7	38		


Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




BBS UK graph



BBS England graph




BBS Scotland graph



BBS Wales graph


Fledglings per breeding attempt Wheatear 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Wheatear




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	16	Curvilinear	3.84 fledglings	4.99 fledglings	29.9%		
Clutch size	43	1968-2011	11	None					Small sample
Brood size	43	1968-2011	57	None					
Nest failure rate at egg stage	43	1968-2011	16	Curvilinear	0.79% nests/day	0.01% nests/day	-98.7%		Small sample
Nest failure rate at chick stage	43	1968-2011	38	None					
Laying date	43	1968-2011	12	None			0 days		Small sample


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend




Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of adult birds surviving to following year - error bars represent 95% confidence limits


Key facts

Conservation listings:	Europe: no SPEC category (concentrated in Europe, conservation status favourable) (BiE04) UK: amber (species level, race <i>occidentalis</i> , 25-50% population decline; race <i>hebridium</i> , >20% of European breeders) (BoCC3) UK Biodiversity Action Plan: priority species
Long-term trend:	UK, England: moderate decline
Population size:	2.5 million territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)

Status summary

Dunnock abundance fell substantially between the mid 1970s and mid 1980s, after a period of population stability. Some recovery has occurred throughout the UK since the late 1990s, but the species still meets amber-list criteria. The BBS Fuller et al. 2005). There has been little variation in survival rates over time (Siriwardenæt al. 1998a). Clutch and brood sizes, and the number of fledglings per breeding attempt all increased as the population fell and appear to have fallen again during the recent period of recovery. Numbers have shown widespread moderate decrease across Europe since 1980 (PECBMS 2013a).

CBC/BBS UK 1966-2012
Dunnock





Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	923	-33	-42	-24	>25	
	25	1986-2011	1461	17	7	29		
	10	2001-2011	2333	8	4	12		
	5	2006-2011	2611	1	-1	3		

CBC/BBS England Source	44 Period (yrs)	1967-2011 Years	763 Plots (201)	-37 Change (%)	-45 Lower limit	-25 Upper limit	>25 Alert	Comment
	10	2001-2011	1901	8	5	11		
	5	2006-2011	2147	2	0	5		
CES adults	27	1984-2011	97	-20	-32	-8		
	25	1986-2011	102	-8	-19	2		
	10	2001-2011	103	-15	-21	-6		
	5	2006-2011	102	-15	-21	-9		
CES juveniles	27	1984-2011	95	-28	-49	4		
	25	1986-2011	99	-15	-35	14		
	10	2001-2011	102	-17	-29	-2		
	5	2006-2011	99	-11	-21	3		
BBS UK	16	1995-2011	2042	22	18	28		
	10	2001-2011	2333	8	5	12		
	5	2006-2011	2611	1	-1	3		
BBS England	16	1995-2011	1654	17	11	23		
	10	2001-2011	1878	8	4	12		
	5	2006-2011	2110	2	-1	4		
BBS Scotland	16	1995-2011	141	60	30	92		
	10	2001-2011	162	21	2	37		
	5	2006-2011	188	6	-6	18		
BBS Wales	16	1995-2011	151	29	12	52		
	10	2001-2011	173	9	0	20		
	5	2006-2011	175	-5	-11	3		
BBS N.Ireland	16	1995-2011	70	62	1	104		
	10	2001-2011	84	-20	-29	-9		
	5	2006-2011	86	-18	-26	-10		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




CES adult abundance 1983-2012
Dunnock




CES adults graph

CES juvenile abundance 1983-2012
Dunnock




CES juveniles graph

BBS UK 1994-2012
Dunnock




BBS UK graph

BBS England 1994-2012
Dunnock




BBS England graph

BBS Scotland 1994-2012
Dunnock




BBS Scotland graph

BBS Wales 1994-2012
Dunnock




BBS Wales graph



Demographic trends


Fledglings per breeding attempt Dunnock 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Dunnock




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	118	Curvilinear	1.68 fledglings	1.76 fledglings	4.4%		
Clutch size	43	1968-2011	108	Linear increase	3.96 eggs	4.15 eggs	4.9%		
Brood size	43	1968-2011	122	Curvilinear	3.41 chicks	3.45 chicks	1.2%		
Nest failure rate at egg stage	43	1968-2011	154	Curvilinear	2.51% nests/day	2.31% nests/day	-8.0%		
Nest failure rate at chick stage	43	1968-2011	124	Curvilinear	2.46% nests/day	2.50% nests/day	1.6%		
Laying date	43	1968-2011	85	None			0 days		
Juvenile to Adult ratio (CES)	27	1984-2011	101	Smoothed trend	102 Index value	100 Index value	-2%		
Juvenile to Adult ratio (CES)	25	1986-2011	106	Smoothed trend	113 Index value	100 Index value	-12%		
Juvenile to Adult ratio (CES)	10	2001-2011	107	Smoothed trend	112 Index value	100 Index value	-10%		
Juvenile to Adult ratio (CES)	5	2006-2011	105	Smoothed trend	89 Index value	100 Index value	12%		


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend




Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend




Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Smoothed long-term trend in ratio of juvenile:adult birds caught - green lines indicate 85% confidence limits



Proportion of adult birds surviving to following year - green bars represent 95% confidence limits

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

House Sparrow

Passer domesticus

Key facts


Conservation listings:	Europe: SPEC category 3 (declining) (BiE04) UK: red (>50% population decline) (BoCC3) UK Biodiversity Action Plan: priority species
Long-term trend:	England: rapid decline
Population size:	5.3 (4.8-5.8) million pairs in 2009 (APEP13: distance sampling estimate for 2006 (Newson et al. 2008) updated using BBS trend)
Migrant status:	Resident
Nesting habitat:	Cavity nester
Primary breeding habitat:	Human habitats
Secondary breeding habitat:	
Breeding diet:	Vegetation
Winter diet:	Vegetation

Status summary

CBC sample sizes did not allow monitoring of House Sparrows until 1976; previously, there had been many farmland plots with high populations that CBC volunteers could not properly quantify without better access to farm buildings and housing. CBC/BBS data indicate a rapid decline in abundance over the last 25 years, as does the BTO's Garden Bird Feeding Survey (Siriwardena et al. 2002, Robinson et al. 2005b). These results are supported by many other studies and anecdotal reports, and have generated considerable conservation concern (see Summers-Smith 2003). The overall national decline since the 1970s masks much heterogeneity by region and habitat, and population processes may be relatively fine-grained: overall, populations in rural areas had declined by 47% by 2000, and those in urban and suburban areas by about 60% (CBC and GBFS data: Robinson et al. 2005b). The BBS PECBMS 2013a). The European status of this species is no longer considered 'secure' (BirdLife International 2004).

CBC/BBS England 1976–2012

House Sparrow




Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	34	1977-2011	673	-68	-77	-57	>50	
	25	1986-2011	903	-43	-57	-23	>25	
	10	2001-2011	1444	2	-2	8		
	5	2006-2011	1603	7	4	11		
BBS UK	16	1995-2011	1580	-1	-7	5		
	10	2001-2011	1753	8	3	13		
	5	2006-2011	1927	8	5	11		
BBS England	16	1995-2011	1302	-12	-18	-6		
	10	2001-2011	1433	2	-3	7		
	5	2006-2011	1585	7	3	10		
BBS Scotland	16	1995-2011	92	38	9	70		
	10	2001-2011	102	16	-4	36		
	5	2006-2011	116	4	-10	20		
BBS Wales	16	1995-2011	122	99	69	137		
	10	2001-2011	140	36	20	56		
	5	2006-2011	143	15	4	31		
BBS N.Ireland	16	1995-2011	53	63	-4	133		
	10	2001-2011	64	29	3	55		
	5	2006-2011	68	14	-3	35		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




CBC/BBS England 1976–2012
House Sparrow




CBC/BBS England graph

BBS UK 1994–2012
House Sparrow




BBS UK graph


BBS England 1994–2012
House Sparrow




BBS England graph



BBS Scotland graph




BBS Wales graph



BBS N.Ireland graph

Demographic trends


Fledglings per breeding attempt House Sparrow 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

House Sparrow




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	103	Curvilinear	2.30 fledglings	2.47 fledglings	7.3%		
Clutch size	43	1968-2011	82	None					
Brood size	43	1968-2011	164	Curvilinear	3.36 chicks	2.86 chicks	-14.8%		
Nest failure rate at egg stage	43	1968-2011	115	Linear decline	1.09% nests/day	0.38% nests/day	-65.1%		
Nest failure rate at chick stage	43	1968-2011	119	Curvilinear	1.51% nests/day	0.69% nests/day	-54.3%		
Laying date	43	1968-2011	64	Linear decline	May 25	May 16	-9 days		


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend




Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of adult birds surviving to following year - error bars represent 95% confidence limits

Causes of change

There is evidence that changes in survival rates due to lack of food resources, because of agricultural intensification, are the main driver of House Sparrow declines in farmland, although changes in breeding performance may also have played a role. Different processes have affected House Sparrows in towns, where breeding performance could be the most important driver of declines, although the evidence for the ecological causes is less clear.

Change factor	Primary driver	Secondary driver
Demographic	Decreased survival	Decreased breeding performance
Ecological	Agricultural intensification	

Further information on causes of change

A temporary drop in first-year survival coincided with the period of steepest decline, but changes in breeding performance, especially reduced nest failure rates at the chick stage, appear to have driven a levelling-off in the long-term population trend (Freeman & Crick 2002). Over the period since 1968, brood size has decreased (see above) but there has also been a major decrease in nest failure rates at the egg and chick stages, so the number of fledglings per breeding attempt has shown a net increase. Further evidence for the role of changing survival in House Sparrow declines has been provided by Hole et al. (2002), who found no evidence of significant differences in most breeding-ecology parameters in declining and stable populations in a farm-scale comparison, while Siriwardena et al. (1999) found that national survival rates were lower during the period of decline in the CBC index. Crick & Siriwardena (2002), using NRS data, showed that breeding performance per nesting attempt had increased and was positively correlated with population growth rate in the wider countryside (although there was no such correlation in gardens). Analysis of

Garden BirdWatch data found higher seasonal peak counts, however, relative to pre-breeding numbers, in the north and west of Britain than in the east and south where population decline is strongest, thus indicating that breeding productivity is influencing population trends (Morrison et al. 2014).

There appear to be different processes affecting urban and agricultural populations. On farmland, changes in farming practices due to intensification of agriculture and the tidying of farmyards have reduced the seed available to farmland populations of House Sparrows during winter, which has resulted in a reduction in survival rates (Siriwardena et al. 1999, Chamberlain et al. 2007, Hole 2001), specifically of first-year birds (Crick et al. 2002). This is supported by a positive effect of supplementary seed in winter on farmland House Sparrow population trends in a landscape-scale experiment in East Anglia (Siriwardena et al. 2007). House Sparrows have probably been deleteriously affected by the decrease in the amount of grain spilt around farm buildings and during the process of harvesting since the 1970s (O'Connor & Shrubbs 1986). The move towards autumn-sowing of cereals has meant that cereal stubble has become much rarer, reducing food resources over winter, although Robinson et al. (2001) found no influence of spring-sown cereal on House Sparrow abundance in predominantly pastoral farmland. Conversely, breeding performance is worse where there is more spring cereal (Crick & Siriwardena 2002), although this may reflect geographical associations with areas where spring sowing remains widespread in the UK (the west and north) rather than direct effects of cropping.

Recent declines have been particularly severe in urban areas (Robinson et al. 2005b, Chamberlain et al. 2007). Increased predation by cats and [Sparrowhawks](#), lack of nest sites, loss of food supplies, pollution and disease have all been cited as factors possibly depressing populations in towns (Crick et al. 2002), but supporting evidence for these is mixed. Within urban areas, Shaw et al. (2008) reviewed available evidence and hypothesised that House Sparrows have disappeared from more affluent areas, where changes to habitat structure such as planting of ornamental shrubs and increased demand for off-street parking is likely to reduce the amount of habitat available to House Sparrows and influenced foraging and predation risk. The conversion of private gardens to continuous housing has also had a negative effect on House Sparrow abundance (Chamberlain et al. 2007). Vincent (2005) found that annual productivity among suburban and rural human habitation in Leicestershire was lower than that measured on farmland House Sparrows in Oxfordshire, the main cause of the difference being starvation of chicks. Low body masses at fledging, and consequently low post-fledging survival, were also recorded in Leicestershire. Although only a two-year study, Peach et al. (2008) measured reproductive success in a declining House Sparrow population along an urbanisation gradient in Leicester and also found that a year in which reproductive success was too low to sustain the population was characterised by lower chick survival and body mass at fledging (a predictor of post-fledging survival). However, there is no direct evidence that invertebrate food supplies have declined in these areas and variation in survival has not been investigated.

Negative correlations between indices of Sparrowhawk presence during its post-organochlorine increase and House Sparrow abundance from the Garden Bird Feeding Survey have been interpreted as evidence that increasing predation rates are depressing House Sparrow populations (Bell et al. 2010). However, more sophisticated analyses of large-scale and extensive national monitoring data provide no evidence that House Sparrow population declines were linked to increases in Sparrowhawks (Newson et al. 2010b).

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Tree Sparrow

Passer montanus

Key facts


Conservation listings:	Europe: SPEC category 3 (declining) (BiE04) UK: red (>50% population decline) (BoCC3) UK Biodiversity Action Plan: click here, priority species
Long-term trend:	England: rapid decline
Population size:	200,000 territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)
Migrant status:	Resident
Nesting habitat:	Cavity nester
Primary breeding habitat:	Farmland
Secondary breeding habitat:	
Breeding diet:	Vegetation
Winter diet:	Vegetation

Status summary

Tree Sparrow abundance nose-dived spectacularly in the UK between the late 1970s and the early 1990s. BBS data indicate a significant increase since 1994, but it should be remembered that, for every Tree Sparrow today there were perhaps around 20 in the 1970s, and any recovery therefore has a very long way to go. Clear range contractions occurred between the first two breeding atlas periods (Gibbons et al. 1993), and have accelerated subsequently: Tree Sparrows have now withdrawn completely from some southern and western regions of Britain, but conversely have spread in Northern Ireland (Balmer et al. 2013). Following declines across western and northwestern Europe during the 1990s, the European status of this species is no longer considered 'secure' (BirdLife International 2004). There has been widespread moderate decrease across Europe since 1980 (PECBMS 2013a).

CBC/BBS England 1966–2012

Tree Sparrow



Smoothed population index, relative to an arbitrary 100 in the year given, with 85% confidence limits in green


Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	44	1967-2011	93	-95	-98	-92	>50	
	25	1986-2011	103	-60	-81	-33	>50	Small CBC sample
	10	2001-2011	149	62	37	102		
	5	2006-2011	169	46	28	67		
BBS UK	16	1995-2011	170	113	71	168		
	10	2001-2011	189	81	42	120		
	5	2006-2011	218	52	24	77		
BBS England	16	1995-2011	136	75	39	119		
	10	2001-2011	148	57	27	89		
	5	2006-2011	168	44	21	66		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




CBC/BBS England 1966–2012
Tree Sparrow




CBC/BBS England graph

BBS UK 1994–2012
Tree Sparrow



BBS UK graph

BBS England 1994–2012
Tree Sparrow




BBS England graph

Demographic trends

Fledglings per breeding attempt


Tree Sparrow 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Tree Sparrow




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	253	Linear increase	2.74 fledglings	3.87 fledglings	41.3%		
Clutch size	43	1968-2011	278	Curvilinear	4.77 eggs	5.17 eggs	8.4%		
Brood size	43	1968-2011	372	Curvilinear	3.81 chicks	4.18 chicks	9.6%		
Nest failure rate at egg stage	43	1968-2011	367	Linear decline	0.85% nests/day	0.32% nests/day	-62.4%		
Nest failure rate at chick stage	43	1968-2011	253	Linear decline	1.50% nests/day	0.54% nests/day	-64.0%		
Laying date	43	1968-2011	281	Linear decline	May 28	May 23	-5 days		


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

Causes of change

The mechanisms underlying the decline in this species are largely unknown, although demographic trends suggest that factors operating during the breeding season are not the main driver.

Change factor	Primary driver	Secondary driver
Demographic	Decreased survival?	
Ecological	Agricultural intensification	

Further information on causes of change

The mechanisms underlying the decline in this species are largely unknown. The number of fledglings per breeding attempt has improved substantially as population sizes have decreased (see above), suggesting that decreases in productivity were not responsible for the decline. This has been driven by declines in daily failure rate at both the nest and chick stages and increases in clutch and brood sizes. It is thus more likely that survival has been the critical demographic measure, although ring-recovery analyses have produced equivocal results, perhaps because of small sample sizes (Siriwardena et al. 1998, 2000b).

Components of agricultural intensification, such as reductions in winter stubble, have been implicated in the decline, although direct evidence supporting such ideas is largely incidental. Tree Sparrows aggregate in areas where seed food is available during the winter and they have declined at the same time as other farmland seed-eaters (Siriwardena et al. 1998), providing circumstantial evidence for shortage of food. In winter in Scotland (Hancock & Wilson 2003), the highest densities of Tree Sparrows were recorded in cereal stubble fields (undersown with grass) and weedy brassica fodder crops. These habitats remain relatively seed-rich but have declined in area in the UK (Fuller 2000, Hancock & Wilson 2003). Field & Anderson (2004) also state that anecdotal evidence suggests that many Tree Sparrow colonies are strongly associated with winter seed food sources, and provision of new seed sources is frequently associated with the establishment of new breeding colonies. Although Siriwardena et al. (2007) did not find a significant positive relationship between winter food supply and breeding population trajectory in areas provisioned by RSPB Bird Aid, this may be due to the fact that the BBS trends for this species are increasing; therefore winter food may not currently be limiting, as the remaining populations are in small remnants of suitable habitat and many are subject to active conservation action (e.g. provision of nest boxes).

During the breeding season, Field & Anderson (2004) found that wetland-edge habitats played a key role in providing invertebrate prey to allow successful chick rearing throughout the long breeding season and suggest that it is possible that large areas of UK farmland that were formerly occupied no longer provide these invertebrate resources, due to the effects of intensification in the late 20th century.

Yellow Wagtail

Motacilla flava

Key facts


Conservation listings:	Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe) (BiE04) UK: red (species level, races flavissima and flava) (BoCC3) UK Biodiversity Action Plan: priority species
Long-term trend:	UK, England: rapid decline
Population size:	15,000 territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)
Migrant status:	Long-distance migrant
Nesting habitat:	Ground nester
Primary breeding habitat:	Farmland
Secondary breeding habitat:	
Breeding diet:	Animal
Winter diet:	Animal

Status summary

Britain holds almost the entire world population of the distinctive race flavissima, so population changes in the UK are of global conservation significance. Yellow Wagtails have been in rapid decline since the early 1980s, according to CBC/BBS and especially WBS/WBBS and, after a shift from the green to the amber list in 2002, the species was moved to the red list in 2009 (Eaton et al. 2009). Gibbons et al. (1993) identified a range contraction towards a core area in central England, concurrent with the early years of decline. Further range contraction has occurred extensively since then, especially in the west and south and in parts of East Anglia (Balmer et al. 2013). The significant decrease in brood size warrants NRS concern (Leech & Barimore 2008). The European trend, which comprises several races of the species, has been of moderate decline since 1980, though with little change since 1990 (PECBMS 2013a).

CBC/BBS UK 1966–2012

Yellow Wagtail



Smoothed population index, relative to an arbitrary 100 in the year given, with 85% confidence limits in green


Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	80	-73	-88	-39	>50	
	25	1986-2011	117	-65	-78	-51	>50	Small CBC sample
	10	2001-2011	152	-25	-34	-13	>25	
	5	2006-2011	159	4	-9	18		
CBC/BBS England	44	1967-2011	79	-72	-88	-46	>50	
	25	1986-2011	114	-64	-77	-48	>50	Small CBC sample
	10	2001-2011	149	-25	-33	-13		
	5	2006-2011	155	4	-10	16		
WBS/WBBS waterways	36	1975-2011	24	-95	-99	-89	>50	
	25	1986-2011	22	-93	-97	-85	>50	
	10	2001-2011	23	-49	-70	-10	>25	
	5	2006-2011	20	-17	-45	25		
BBS UK	16	1995-2011	157	-45	-53	-34	>25	
	10	2001-2011	152	-25	-33	-9		
	5	2006-2011	158	3	-10	16		
BBS England	16	1995-2011	153	-45	-52	-34	>25	
	10	2001-2011	149	-24	-34	-12		
	5	2006-2011	154	3	-11	19		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




CBC/BBS UK 1966–2012
Yellow Wagtail




CBC/BBS UK graph

CBC/BBS England 1966–2012
Yellow Wagtail





CBC/BBS England graph

WBS/WBBS 1974–2012
Yellow Wagtail




WBS/WBBS waterways graph



Demographic trends


Fledglings per breeding attempt Yellow Wagtail 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Yellow Wagtail




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Brood size	43	1968-2011	12	Linear decline	4.83 chicks	4.28 chicks	-11.3%		Small sample


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

Causes of change

Agricultural intensification is the ultimate cause of population declines. However, the mechanisms underlying the decline remain unclear.

Change factor	Primary driver	Secondary driver
Demographic	Unknown	
Ecological	Agricultural intensification	

Further information on causes of change

Changes in agricultural practices have been proposed as the main reason for declines via their impact on the quality of foraging and breeding habitats. The magnitude of Yellow Wagtail decline appears to vary between habitats, being strongest in wet grassland and marginal upland areas (Henderson et al. 2004, Wilson & Vickery 2005). Chamberlain & Fuller (2000, 2001) found that there were greater range contractions in regions dominated by pastoral agriculture. The decline in pastoral habitats has been proposed to be due to agricultural intensification, specifically farmland drainage, the conversion of pasture to arable land, changes in grazing and cutting regimes, the loss of insects associated with cattle and changes to grassland ecosystems in marginal upland areas (Gibbons et al. 1993, Chamberlain & Fuller 2000, 2001, Flyckt 1999, Vickery et al. 2001, Nelson et al. 2003, Bradbury & Bradter 2004, Henderson et al. 2004). Such changes are likely to have reduced the quality of grasslands as a nesting and foraging habitat. A detailed study on Yellow Wagtail breeding ecology by Bradbury & Bradter (2004) provided good evidence of the species' breeding requirements on grassland. Territories were associated with a greater proportion of bare earth in the sward, the presence of shallow-edged ponds or wet ditches in the field, and a greater probability of a prolonged winter/spring flood, although the relative importance of these and how they impact upon demographic processes was indecipherable.

Data from eastern England suggest a strong avoidance of grassland and preference for spring-sown crops (Mason & Macdonald 2000), though breeding can also be successful in landscapes dominated by winter cereals (Kirby et al. 2012). A detailed autecological study by Gilroy et al. (2008) provides good evidence that, on arable land, soil penetrability had a significant influence on the abundance of Yellow Wagtails, together with crop type and soil type, as these influenced invertebrate capture rates. There was a strong relationship between Yellow Wagtails and soil penetrability, suggesting a potential causative link between soil degradation and population decline (Gilroy et al. 2008). Breeding-season length may also be limited in cereal-dominated areas, as Yellow Wagtails avoid autumn-sown cereals late in the season (Gilroy et al. 2009, 2010). Predation was also considered and it was found that predation rate was closer nearer to tramlines and field-edges (Morris & Gilroy 2008). It is uncertain how important nest predation in tramlines is as a limiting factor for Yellow Wagtail populations but no studies have reported predation as a major driver of population decline for this species. Work carried out by Benton et al. (2002) showed that, in Scotland, arthropod abundance was significantly related to agricultural change and that this was also linked to measures of farmland bird density. Although Yellow Wagtail does not breed on Scottish farmland, it is an obligate insectivore, so this evidence adds support to the hypothesis that reduced food availability due to agricultural change may have contributed to the declines in this species.

Yellow Wagtails are long-distance migrants, moving to wintering grounds in western Africa south of the Sahara. Factors relating to conditions on the wintering grounds

may also play a role (Bradbury & Bradter 2004, Heldbjerg & Fox 2008, Stevens et al. 2010) but evidence for this is lacking.

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>


Key facts

Conservation listings:	Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe) (BiE04) UK: amber (25-50% population decline) (BoCC3)
Long-term trend:	UK waterways: rapid decline
Population size:	38,000 pairs in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)

Status summary

Grey Wagtails occur at highest densities along fast-flowing upland streams. WBS/WBBS shows a fluctuating population size along waterways, with a fall during the late 1970s and early 1980s from an initial high point in 1974, some increase since the late 1990s, and another recent fall. The species was moved from the green to the amber list in 2002, because of a 41% decline recorded between 1975 and 1999. The BBS trend matches WBS/WBBS closely: there was an initial increase but since 2002 the trend has been steeply downward, especially in Scotland. The trends for Grey Wagtail are very similar to those for Leech & Barimore 2008). Nest failure rates have dropped substantially, and there has been linear increase in the number of fledglings per breeding attempt, suggesting that reduced survival is the likely driver of decline. Numbers have shown widespread moderate decrease across Europe since 1990 (PECBMS 2013a).

WBS/WBBS 1974-2012
Grey Wagtail




Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
WBS/WBBS waterways	36	1975-2011	96	-58	-67	-44	>50	
	25	1986-2011	115	-25	-36	-9	>25	
	10	2001-2011	170	-41	-49	-33	>25	
	5	2006-2011	150	-42	-48	-34	>25	

BBS UK Source	Period (yrs)	1995-2011 Years	Points (n)	Change (%)	Lower limit	Upper limit	>25 Alert	Comment
	10	2001-2011	260	-53	-58	-44	>50	
	5	2006-2011	279	-51	-54	-40	>50	
BBS England	16	1995-2011	148	-19	-31	-1		
	10	2001-2011	177	-42	-49	-32	>25	
	5	2006-2011	196	-41	-46	-29	>25	
BBS Scotland	16	1995-2011	33	-32	-53	0	>25	
	10	2001-2011	37	-64	-74	-45	>50	
	5	2006-2011	40	-56	-69	-38	>50	


Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




WBS/WBBS waterways graph



BBS UK graph




BBS England graph



BBS Scotland graph

Fledglings per breeding attempt


Grey Wagtail 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Grey Wagtail




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	54	Linear increase	2.60 fledglings	3.37 fledglings	29.4%		
Clutch size	43	1968-2011	38	Curvilinear	4.74 eggs	4.66 eggs	-1.8%		
Brood size	43	1968-2011	80	Curvilinear	4.03 chicks	4.08 chicks	1.3%		
Nest failure rate at egg stage	43	1968-2011	57	Linear decline	1.79% nests/day	0.96% nests/day	-46.4%		
Nest failure rate at chick stage	43	1968-2011	56	Linear decline	2.19% nests/day	0.74% nests/day	-66.2%		
Laying date	43	1968-2011	60	Curvilinear	May 5	Apr 27	-8 days		


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend




Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

Key facts

Conservation listings:	Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe) (BiE04) UK: green (species level); amber (race yarrellii, >20% of European breeders) (BoCC3)
Long-term trend:	UK: uncertain
Population size:	470,000 (410,000-520,000) pairs in 2009 (APEP13: distance sampling estimate for 2006 (Newson et al. 2008) updated using BBS trend)

Status summary

Britain and Ireland together hold almost the entire world population of the distinctive dark-backed race yarrellii (Pied Wagtail), and for this reason population changes in the UK are of global conservation significance. The CBC shows that a strong increase occurred up to the mid 1970s, such that populations have shown shallow increase overall since 1967. Since 1974, however, the results of monitoring have been somewhat conflicting: CBC/BBS and WBS/WBBS trends fluctuate in parallel but, whereas little overall change is evident in the CBC/BBS index, WBS/WBBS has shown a rapid decline, suggesting a strong influence of factors specific to linear waterways. The BBS Siriwardena et al. 1998a). Average clutch and brood sizes have declined a little, raising NRS concern (Leech & Barimore 2008), but this has been counteracted by a large fall in nest failure rates. The number of fledglings per breeding attempt has shown a strong linear increase. The European long-term trend, which includes the nominate race of the species (White Wagtail), has shown little change since 1980 (PECBMS 2013a).







Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	535	50	9	107		
	25	1986-2011	874	6	-14	43		
	10	2001-2011	1401	-23	-29	-16		


Source	5 Period (yrs)	2006-2011 Years	1508 Plots	-16 Change (%)	-24 Lower Limit	-8 Upper Limit	Alert	Comment
CBC/BBS England		1967-2011	112					
	25	1986-2011	669	2	-20	38		
	10	2001-2011	1072	-28	-32	-24	>25	
	5	2006-2011	1164	-22	-25	-18		
WBS/WBBS waterways	36	1975-2011	112	-67	-75	-60	>50	
	25	1986-2011	132	-41	-52	-30	>25	
	10	2001-2011	197	-30	-38	-22	>25	
	5	2006-2011	180	-19	-28	-12		
BBS UK	16	1995-2011	1256	-13	-19	-5		
	10	2001-2011	1401	-23	-28	-16		
	5	2006-2011	1508	-16	-22	-6		
BBS England	16	1995-2011	959	-15	-21	-9		
	10	2001-2011	1072	-27	-32	-23	>25	
	5	2006-2011	1164	-22	-25	-17		
BBS Scotland	16	1995-2011	129	-13	-33	5		
	10	2001-2011	133	-19	-36	1		
	5	2006-2011	142	-3	-26	32		
BBS Wales	16	1995-2011	113	-9	-22	5		
	10	2001-2011	126	-15	-27	0		
	5	2006-2011	124	-20	-30	-9		
BBS N.Ireland	16	1995-2011	44	11				
	10	2001-2011	54	-9				
	5	2006-2011	57	-16				

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.







BBS UK graph




BBS England graph



BBS Scotland graph




BBS Wales graph



BBS N.Ireland graph

Fledglings per breeding attempt


Pied Wagtail 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Pied Wagtail




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	88	Linear increase	2.99 fledglings	3.58 fledglings	19.9%		
Clutch size	43	1968-2011	65	Linear decline	5.08 eggs	4.93 eggs	-2.9%		
Brood size	43	1968-2011	136	Linear decline	4.50 chicks	4.32 chicks	-3.9%		
Nest failure rate at egg stage	43	1968-2011	89	Linear decline	1.88% nests/day	0.64% nests/day	-66.0%		
Nest failure rate at chick stage	43	1968-2011	98	Linear decline	1.26% nests/day	0.86% nests/day	-31.7%		
Laying date	43	1968-2011	88	Curvilinear	May 18	May 16	-2 days		


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglinton, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Tree Pipit

Anthus trivialis

Key facts


Conservation listings:	Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe) (BiE04) UK: red (>50% population decline) (BoCC3) UK Biodiversity Action Plan: priority species
Long-term trend:	England: rapid decline
Population size:	88,000 (55,000-121,000) pairs in 2009 (APEP13: distance sampling estimate for 2006 (Newson et al. 2008) updated using BBS trend)
Migrant status:	Long-distance migrant
Nesting habitat:	Ground nester
Primary breeding habitat:	Woodland
Secondary breeding habitat:	
Breeding diet:	Animal
Winter diet:	Animal

Status summary

Tree Pipits occur in greatest abundance in Wales, northern England and Scotland, and thus the marked CBC decline between the first two atlas periods may reflect the range contraction that occurred then in central and southeast England (Gibbons et al. 1993). Since 1994, CBC/BBS data for the species have shown a further severe decrease, especially in England. Recent atlas data show further losses of range, especially in eastern England (Balmer et al. 2013). Brood size has increased since 1966 but nest losses at chick stage have increased greatly and there has been a decrease in fledglings per breeding attempt. Laying dates have shifted earlier by over a week. Although the species has no European conservation listing as yet, numbers have shown widespread moderate decrease across Europe since 1980 (PECBMS 2013a), and the mean change across all European countries during the 1990s was a significant decline (Sanderson et al. 2006). The species was moved from the green to the amber list of UK Birds of Conservation Concern in 2002, and in 2009 to red, on the strength of its UK population decline (Eaton et al. 2009).

CBC/BBS England 1966–2012

Tree Pipit




Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	44	1967-2011	49	-86	-92	-74	>50	
	25	1986-2011	60	-86	-92	-77	>50	Small CBC sample
	10	2001-2011	77	-28	-43	-9	>25	
	5	2006-2011	89	-14	-29	2		
BBS UK	16	1995-2011	138	1	-21	32		
	10	2001-2011	144	-6	-21	10		
	5	2006-2011	166	12	-4	29		
BBS England	16	1995-2011	74	-45	-61	-22	>25	
	10	2001-2011	77	-28	-44	-11	>25	
	5	2006-2011	89	-14	-29	2		
BBS Scotland	16	1995-2011	33	84	22	150		
	10	2001-2011	36	15	-14	57		
	5	2006-2011	46	35	1	77		
BBS Wales	16	1995-2011	32	-9	-37	33		
	10	2001-2011	31	-11	-38	20		
	5	2006-2011	32	4	-17	41		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




CBC/BBS England 1966-2012
Tree Pipit




CBC/BBS England graph

BBS UK 1994-2012
Tree Pipit





BBS UK graph

BBS England 1994-2012
Tree Pipit




BBS England graph




Demographic trends

Fledglings per breeding attempt Tree Pipit 1966–2012



Laying date 1966–2012

Tree Pipit




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	13	Curvilinear	1.54 fledglings	1.24 fledglings	-19.5%		
Clutch size	43	1968-2011	11	None					Small sample
Brood size	43	1968-2011	29	Curvilinear	4.30 chicks	4.49 chicks	4.4%		Small sample
Nest failure rate at egg stage	43	1968-2011	14	Curvilinear	4.80% nests/day	4.35% nests/day	-9.4%		Small sample
Nest failure rate at chick stage	43	1968-2011	22	Curvilinear	3.30% nests/day	5.31% nests/day	60.9%		Small sample
Laying date	43	1968-2011	20	Curvilinear	May 27	May 19	-8 days		Small sample


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

Causes of change

The availability of suitably structured habitat is important and lack of this may have contributed to the decline, possibly through a decrease in nest survival, although evidence for this is based largely on one site. This species being a long-distance migrant, problems on its wintering grounds should not be ruled out.

Change factor	Primary driver	Secondary driver
Demographic	Decreased breeding success	
Ecological	Changes in woodland	

Further information on causes of change

A detailed, eight-year study in Thetford Forest conducted by Burton (2009) provides good evidence that there was a significant decrease in daily nest survival during the chick stage and that overall nesting success was lowest in clearfells and recently planted stands. Overall nesting success appeared to be determined at the habitat scale, and Burton suggested that this may have been because the broad differences in cover between habitats affected the likelihood of nest predation (the main cause of nest failure). Charman et al. (2009) also found that Tree Pipits have high failure rates at the chick stage and implicate predation. It should be noted that records from Thetford Forest, in southeast England, probably contribute over half the nest records for this species each year: thus these trends may not be representative of the UK as a whole.

This species prefers open ground within woodlands and upland grazed woods lacking understorey, and also occupies clearfells, restocks, new plantations, heaths and commons where trees provide songposts (Fuller 1995, Burton 2007, Charman et al. 2009). The species' decline has been greatest in lowland England, particularly in the wider countryside in woodland and common land (Gibbons et al. 1993) and, accordingly, several authors have proposed that the population decline may be linked to the changing forest structure as new plantations mature, and the reduced management of lowland woods (Fuller et al. 2005, Amar et al. 2006, Charman et al. 2009). Data provided by the Repeat Woodland Bird Survey (RWBS) gives reliable evidence that sub-canopy vegetation increased markedly in almost all regions covered between the 1980s and the early 2000s and analyses found that declines of Tree Pipit occurred in woods with higher maximum tree height and increased foliage (Amar et al. 2006, Smart et al. 2007). Fuller & Moreton (1987) and Burton (2007) provide evidence, respectively, for associations with young coppice and, within coniferous plantations, for young restocks, and a disassociation with closed-canopy woodlands. Amar et al. (2006) state that the lack of new plantations and restocks in southern Britain may be contributing to the decline of this species, although specific analyses providing evidence for this are lacking. They also found that Tree Pipit declined more in sites with more tracks, suggesting disturbance can be an issue (Amar et al. 2006, Smart et al. 2007). Targeted management, such as the provision of large blocks of habitat and the retention of mature trees for use as songposts, was found to be beneficial (Burton 2007).

In upland habitats, Fuller et al. (2006) provided evidence showing that both overgrazing and agricultural abandonment of marginal habitats may have detrimental effects on Tree Pipits.

Hewson et al. (2007) analysed the RWBS and BBS/CBC data and found declines in all of the seven long-distance migrant species considered, including Tree Pipit. Thus, although specific evidence relating to factors operating on the wintering grounds is lacking, these cannot be ruled out as causes of population decline.

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>


Key facts

Conservation listings:	Europe: no SPEC category (concentrated in Europe, conservation status favourable) (BiE04) UK: amber (25-50% population decline) (BoCC3)
Long-term trend:	England: moderate decline
Population size:	2.0 (1.8-2.3) million pairs in 2009 (APEP13: distance sampling estimate for 2006 (Newson et al. 2008) updated using BBS trend)

Status summary

The CBC/BBS trend has been downward since the mid 1970s. Moorland, the key Meadow Pipit habitat, was not covered well by the CBC, leading to some doubt about the significance of the early results for this species, but BBS now provides more representative monitoring that, in England at least, confirms the picture presented by CBC. The species has accordingly been moved from the green to the amber list. The BBS Gibbons et al. 1993). Experiments in central Scotland have indicated that Meadow Pipit breeding abundance can be improved by reduced grazing intensity and by mixing cattle and sheep (Evans et al. 2006). Nest failure rates during the chick stage have declined markedly, which may reflect the loss of birds from suboptimal habitat, and the number of fledglings per breeding attempt has increased. A widespread moderate decline is evident across Europe since 1980 (PECBMS 2013a, Lehikoinen et al. 2014).

CBC/BBS England 1966-2012
Meadow Pipit



Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	44	1967-2011	188	-48	-75	-22	>25	
	25	1986-2011	302	-26	-44	-9	>25	
	10	2001-2011	501	-12	-20	-4		
	5	2006-2011	580	-2	-8	3		

BBS UK Source	Period (yrs)	1995-2011 Years	PBS (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
	10	2001-2011	884	-22	-26	-18		
	5	2006-2011	984	-10	-15	-5		
BBS England	16	1995-2011	432	-14	-24	-3		
	10	2001-2011	501	-12	-20	-2		
	5	2006-2011	580	-3	-8	3		
BBS Scotland	16	1995-2011	200	-29	-37	-21	>25	
	10	2001-2011	196	-24	-30	-16		
	5	2006-2011	207	-9	-15	0		
BBS Wales	16	1995-2011	86	-10	-22	0		
	10	2001-2011	93	-23	-31	-15		
	5	2006-2011	92	-12	-20	-7		
BBS N.Ireland	16	1995-2011	64	-29	-41	-7	>25	
	10	2001-2011	73	-51	-58	-46	>50	
	5	2006-2011	73	-51	-56	-43	>50	

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




CBC/BBS England 1966–2012
Meadow Pipit




CBC/BBS England graph

BBS UK 1994–2012
Meadow Pipit




BBS UK graph

BBS England 1994–2012
Meadow Pipit




BBS England graph


BBS Scotland 1994–2012
Meadow Pipit



BBS Scotland graph




BBS Wales graph



BBS N.Ireland graph

Demographic trends


Fledglings per breeding attempt Meadow Pipit 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Meadow Pipit




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	46	Curvilinear	2.07 fledglings	2.32 fledglings	12.5%		
Clutch size	43	1968-2011	37	None					
Brood size	43	1968-2011	74	None					
Nest failure rate at egg stage	43	1968-2011	46	None					
Nest failure rate at chick stage	43	1968-2011	65	Curvilinear	3.09% nests/day	1.77% nests/day	-42.7%		
Laying date	43	1968-2011	39	Curvilinear	May 19	May 18	-1 days		


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend


Key facts

Conservation listings:	Europe: no SPEC category (concentrated in Europe, conservation status favourable) (BiE04) UK: green (species level); amber (race gengleri, >20% of European breeders) (BoCC3)
Long-term trend:	UK, England: shallow increase
Population size:	6.2 million territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)

Status summary

Chaffinch abundance has increased rapidly since the early 1970s, according to CBC/BBS and CES, but numbers seemed to stabilise for a period during the 1990s. This relative stability was associated with a reduction in annual survival, which could be density-dependent (Siriwardena et al. 1999). There was also some evidence of improved breeding performance during the early years of population increase, with larger broods, fewer egg-stage nest failures, and more fledglings per breeding attempt, but these trends are now either cancelled out or reversed. The BBS Robinson et al. 2010). The trend towards earlier laying may be partly explained by recent climate change (Crick & Sparks 1999). Chaffinches are well adapted to suburban and garden habitats, as well as to highly fragmented woodland and hedgerows, occurring less in the open-field, arable habitats that have been affected most by agricultural intensification, so it is possible that they have benefited by environmental changes from which other seed-eating passerines have suffered. Numbers have shown widespread moderate increase across Europe since 1980, though with little change since 1990 (PECBMS 2013a).

CBC/BBS UK 1966-2012
Chaffinch





Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	1091	29	16	47		
	25	1986-2011	1754	12	6	20		

Source	10 Period (yrs)	2001-2011 Years 2006-2011	2798 Plots (n)	5 Change (%)	2 Lower limit	8 Upper limit	Alert	Comment
CBC/BBS England	44	1967-2011	864	30	14	49		
	25	1986-2011	1380	11	4	18		
	10	2001-2011	2193	2	-1	4		
	5	2006-2011	2493	-4	-6	-3		
CES adults	27	1984-2011	78	0	-47	54		
	25	1986-2011	81	-7	-43	29		
	10	2001-2011	84	-16	-31	-6		
	5	2006-2011	80	-23	-32	-15		
CES juveniles	27	1984-2011	61	28	-20	138		
	25	1986-2011	64	71	6	241		
	10	2001-2011	68	55	15	96		
	5	2006-2011	63	16	-8	52		
BBS UK	16	1995-2011	2461	12	8	16		
	10	2001-2011	2798	5	1	7		
	5	2006-2011	3147	0	-2	2		
BBS England	16	1995-2011	1897	12	8	16		
	10	2001-2011	2151	2	-1	5		
	5	2006-2011	2423	-4	-6	-2		
BBS Scotland	16	1995-2011	240	16	5	26		
	10	2001-2011	263	14	4	24		
	5	2006-2011	303	8	0	15		
BBS Wales	16	1995-2011	197	-3	-13	11		
	10	2001-2011	223	7	-1	16		
	5	2006-2011	225	1	-6	9		
BBS N.Ireland	16	1995-2011	90	47	13	64		
	10	2001-2011	106	-7	-14	4		
	5	2006-2011	110	3	-5	12		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




CES adult abundance 1983-2012
Chaffinch




CES adults graph

CES juvenile abundance 1983-2012
Chaffinch




CES juveniles graph

BBS UK 1994-2012
Chaffinch




BBS UK graph

BBS England 1994-2012
Chaffinch




BBS England graph

BBS Scotland 1994-2012
Chaffinch




BBS Scotland graph

BBS Wales 1994-2012
Chaffinch




BBS Wales graph



Demographic trends


Fledglings per breeding attempt Chaffinch 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Chaffinch




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	129	Curvilinear	1.60 fledglings	1.38 fledglings	-13.4%		
Clutch size	43	1968-2011	99	None					
Brood size	43	1968-2011	160	None					
Nest failure rate at egg stage	43	1968-2011	189	Curvilinear	2.96% nests/day	3.78% nests/day	27.7%		
Nest failure rate at chick stage	43	1968-2011	129	Curvilinear	3.00% nests/day	3.21% nests/day	7.0%		
Laying date	43	1968-2011	122	Linear decline	May 12	May 1	-11 days		
Juvenile to Adult ratio (CES)	27	1984-2011	84	Smoothed trend	52 Index value	100 Index value	93%		
Juvenile to Adult ratio (CES)	25	1986-2011	88	Smoothed trend	76 Index value	100 Index value	31%		
Juvenile to Adult ratio (CES)	10	2001-2011	93	Smoothed trend	91 Index value	100 Index value	9%		
Juvenile to Adult ratio (CES)	5	2006-2011	87	Smoothed trend	96 Index value	100 Index value	4%		


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend




Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend




Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Smoothed long-term trend in ratio of juvenile:adult birds caught - green lines indicate 85% confidence limits



Proportion of adult birds surviving to following year - green bars represent 95% confidence limits

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>


Key facts

Conservation listings:	Europe: no SPEC category (not concentrated in Europe, conservation status favourable) (BiE04) UK: amber (25-50% population decline) (BoCC3) UK Biodiversity Action Plan: click here , priority species
Long-term trend:	UK, England: moderate decline
Population size:	220,000 territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)

Status summary

The UK Bullfinch population entered a long period of decline in the mid 1970s, following a period of relative stability. The decline was initially very steep, and more so in farmland than in wooded habitats, but became shallower and eventually ended around 2000, since when there has been substantial increase. CES and CBC/BBS both suggest there are large fluctuations around the overall long-term trend. The BBS Siriwardena et al. 1999, 2000b, 2001), although agricultural intensification and a reduction in the structural and floristic diversity of woodland are suspected to have played a part through losses of food resources and nesting cover (Fuller et al. 2005). Alongside these factors, Proffitt et al. (2004) and Marquiss (2007) mention the constraints on survival outside the breeding season and the possible role of higher Leech & Barimore 2008), and the trend in fledglings per breeding attempt is downward overall. Numbers have shown widespread moderate decrease across Europe since 1980 (PECBMS 2013a). The UK conservation listing was downgraded from red to amber in 2009 (Eaton et al. 2009), since when further population recovery has occurred.

CBC/BBS UK 1966–2012 Bullfinch





Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	328	-36	-48	-22	>25	
	25	1986-2011	464	0	-13	15		
	10	2001-2011	677	34	26	44		

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	44	1967-2011	264	-39	-52	-22	>25	
	25	1986-2011	365	-7	-18	10		
	10	2001-2011	522	28	20	39		
	5	2006-2011	578	25	18	32		
CES adults	27	1984-2011	81	-8	-29	20		
	25	1986-2011	84	-12	-31	10		
	10	2001-2011	83	19	-4	45		
	5	2006-2011	81	15	3	30		
CES juveniles	27	1984-2011	65	-18	-45	30		
	25	1986-2011	68	-2	-29	42		
	10	2001-2011	68	0	-24	28		
	5	2006-2011	68	25	-5	52		
BBS UK	16	1995-2011	594	12	3	22		
	10	2001-2011	677	34	25	44		
	5	2006-2011	745	23	16	31		
BBS England	16	1995-2011	456	8	-1	19		
	10	2001-2011	516	26	18	38		
	5	2006-2011	568	24	16	31		
BBS Scotland	16	1995-2011	41	52	5	97		
	10	2001-2011	46	57	28	102		
	5	2006-2011	56	44	16	86		
BBS Wales	16	1995-2011	62	-5	-25	19		
	10	2001-2011	71	13	-7	34		
	5	2006-2011	66	0	-17	17		
BBS N.Ireland	16	1995-2011	31	35	-26	50		
	10	2001-2011	38	67	21	123		
	5	2006-2011	43	18	-7	43		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




CES adult abundance 1983-2012
Bullfinch




CES adults graph

CES juvenile abundance 1983-2012
Bullfinch




CES juveniles graph

BBS UK 1994-2012
Bullfinch




BBS UK graph

BBS England 1994-2012
Bullfinch




BBS England graph

BBS Scotland 1994-2012
Bullfinch




BBS Scotland graph

BBS Wales 1994-2012
Bullfinch




BBS Wales graph



Demographic trends


Fledglings per breeding attempt Bullfinch 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Bullfinch




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	33	Curvilinear	1.49 fledglings	1.37 fledglings	-7.8%		
Clutch size	43	1968-2011	36	None					
Brood size	43	1968-2011	37	Linear decline	4.18 chicks	3.88 chicks	-7.2%		
Nest failure rate at egg stage	43	1968-2011	51	None					
Nest failure rate at chick stage	43	1968-2011	34	Curvilinear	3.05% nests/day	3.48% nests/day	14.1%		
Laying date	43	1968-2011	34	Linear increase	May 26	Jun 1	6 days		
Juvenile to Adult ratio (CES)	27	1984-2011	85	Smoothed trend	106 Index value	100 Index value	-5%		
Juvenile to Adult ratio (CES)	25	1986-2011	89	Smoothed trend	102 Index value	100 Index value	-2%		
Juvenile to Adult ratio (CES)	10	2001-2011	87	Smoothed trend	110 Index value	100 Index value	-9%		
Juvenile to Adult ratio (CES)	5	2006-2011	85	Smoothed trend	105 Index value	100 Index value	-5%		


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend




Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend




Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Smoothed long-term trend in ratio of juvenile:adult birds caught - green lines indicate 85% confidence limits



Proportion of adult birds surviving to following year - green bars represent 95% confidence limits

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>


Key facts

Conservation listings:	Europe: no SPEC category (concentrated in Europe, conservation status favourable) (BiE04) UK: green (species level, race chloris); amber (race harrisoni, >20% of European breeders) (BoCC3)
Long-term trend:	UK, England: fluctuating, with no long-term trend
Population size:	1.7 (1.6-1.8) million pairs in 2009 (APEP13: distance sampling estimate for 2006 (Newson et al. 2008) updated using BBS trend)

Status summary

Greenfinch abundance varied little up to the mid 1990s, and there was little change in either survival or breeding performance during this period (Siriwardena et al. 1998b, 2000b). More recent CBC/BBS data indicate population increases widely across the UK, followed by a sudden sharp fall induced by a widespread and severe outbreak of the respiratory disease trichomonosis that began in 2005 (Robinson et al. 2010b, Lawson et al. 2012b). The BBS Crick & Sparks 1999). There has been a moderate increase across Europe since 1980 but with no detectable change since 1990 (PECBMS 2013a).

CBC/BBS UK 1966–2012 Greenfinch





Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	786	-20	-35	1		
	25	1986-2011	1268	-1	-16	13		
	10	2001-2011	2071	-33	-36	-30	>25	
	5	2006-2011	2283	-35	-38	-34	>25	
CBC/BBS England	44	1967-2011	666	-13	-33	9		

Source	25 Period (yrs)	1986-2011 Years 2001-2011	1071 Plots (1739)	-1 Change (%)	-16 Lower limit	13 Upper limit	Alert >25	Comment
	5	2006-2011	1928	-35	-38	-34	>25	
CES adults	27	1984-2011	41	12	-51	203		
	25	1986-2011	43	-21	-58	75		
	10	2001-2011	48	-38	-53	-24	>25	
	5	2006-2011	45	-36	-52	-25	>25	
CES juveniles	27	1984-2011	30	162	-11	780		
	25	1986-2011	31	91	-36	531		
	10	2001-2011	38	88	9	173		
	5	2006-2011	37	22	-13	80		
BBS UK	16	1995-2011	1792	-16	-21	-12		
	10	2001-2011	2048	-33	-36	-31	>25	
	5	2006-2011	2245	-35	-37	-33	>25	
BBS England	16	1995-2011	1511	-15	-20	-9		
	10	2001-2011	1724	-33	-35	-30	>25	
	5	2006-2011	1903	-35	-38	-33	>25	
BBS Scotland	16	1995-2011	104	-21	-37	0		
	10	2001-2011	115	-25	-36	-13		
	5	2006-2011	127	-29	-39	-18	>25	
BBS Wales	16	1995-2011	114	-17	-33	12		
	10	2001-2011	132	-39	-47	-26	>25	
	5	2006-2011	134	-39	-45	-31	>25	
BBS N.Ireland	16	1995-2011	51	-10	-39	45		
	10	2001-2011	63	-52	-63	-41	>50	
	5	2006-2011	65	-52	-61	-43	>50	

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




CES adult abundance 1983-2012
Greenfinch




CES adults graph

CES juvenile abundance 1983-2012
Greenfinch




CES juveniles graph

BBS UK 1994-2012
Greenfinch




BBS UK graph

BBS England 1994-2012
Greenfinch




BBS England graph

BBS Scotland 1994-2012
Greenfinch




BBS Scotland graph

BBS Wales 1994-2012
Greenfinch




BBS Wales graph



Demographic trends


Fledglings per breeding attempt Greenfinch 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Greenfinch




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	91	Curvilinear	2.14 fledglings	2.02 fledglings	-5.6%		
Clutch size	43	1968-2011	89	Linear decline	4.76 eggs	4.56 eggs	-4.2%		
Brood size	43	1968-2011	111	Linear decline	4.10 chicks	3.75 chicks	-8.6%		
Nest failure rate at egg stage	43	1968-2011	123	Curvilinear	2.61% nests/day	1.92% nests/day	-26.4%		
Nest failure rate at chick stage	43	1968-2011	92	None					
Laying date	43	1968-2011	91	Linear decline	May 25	May 8	-17 days		
Juvenile to Adult ratio (CES)	27	1984-2011	46	Smoothed trend	95 Index value	100 Index value	5%		
Juvenile to Adult ratio (CES)	25	1986-2011	49	Smoothed trend	63 Index value	100 Index value	59%		
Juvenile to Adult ratio (CES)	10	2001-2011	55	Smoothed trend	51 Index value	100 Index value	96%		
Juvenile to Adult ratio (CES)	5	2006-2011	53	Smoothed trend	64 Index value	100 Index value	56%		


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend




Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Smoothed long-term trend in ratio of juvenile:adult birds caught - green lines indicate 85% confidence limits

Key facts


Conservation listings:	Europe: SPEC category 2 (declining) (BiE04) UK: red (species level, race cannabina); amber (race autochthona, >20% of European breeders, European status) (BoCC3) UK Biodiversity Action Plan: click here , priority species
Long-term trend:	England: rapid decline
Population size:	430,000 territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend for England)
Migrant status:	Short-distance migrant
Nesting habitat:	Above-ground nester
Primary breeding habitat:	Farmland
Secondary breeding habitat:	
Breeding diet:	Vegetation
Winter diet:	Vegetation

Status summary

Linnet abundance fell rapidly in the UK in the late 1960s, and again between the mid 1970s and mid 1980s, but decrease has been followed by a long period of relative stability. Numbers have fallen further in Britain since the start of BBS in 1994, but there has been some increase in Northern Ireland. The BBS PECBMS 2013a), and the European status of this species is no longer considered 'secure' (BirdLife International 2004).

CBC/BBS England 1966–2012

Linnet




Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	44	1967-2011	449	-71	-79	-63	>50	
	25	1986-2011	696	-5	-23	13		
	10	2001-2011	1019	-9	-13	-1		
	5	2006-2011	1077	17	9	24		
BBS UK	16	1995-2011	1179	-19	-24	-12		
	10	2001-2011	1248	-12	-18	-5		
	5	2006-2011	1298	4	-1	10		
BBS England	16	1995-2011	954	-19	-24	-11		
	10	2001-2011	1002	-8	-13	-1		
	5	2006-2011	1048	17	9	23		
BBS Scotland	16	1995-2011	89	-16	-32	14		
	10	2001-2011	93	-20	-38	10		
	5	2006-2011	98	-21	-33	2		
BBS Wales	16	1995-2011	92	-30	-48	-8	>25	
	10	2001-2011	100	-40	-54	-29	>25	
	5	2006-2011	96	-17	-31	-3		
BBS N.Ireland	16	1995-2011	37	41	-11	110		
	10	2001-2011	44	-6	-27	44		
	5	2006-2011	48	-16	-32	13		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




CBC/BBS England 1966–2012
Linnet




CBC/BBS England graph

BBS UK 1994–2012
Linnet




BBS UK graph


BBS England 1994–2012
Linnet




BBS England graph



BBS Scotland graph



BBS Wales graph




BBS N.Ireland graph

Demographic trends

Fledglings per breeding attempt


Linnet 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Linnet




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	122	Linear decline	2.69 fledglings	2.36 fledglings	-12.4%		
Clutch size	43	1968-2011	123	Linear decline	4.74 eggs	4.62 eggs	-2.5%		
Brood size	43	1968-2011	141	Curvilinear	4.11 chicks	4.11 chicks	0.1%		
Nest failure rate at egg stage	43	1968-2011	172	Linear increase	1.85% nests/day	2.35% nests/day	27.0%		
Nest failure rate at chick stage	43	1968-2011	122	Linear increase	1.53% nests/day	2.27% nests/day	48.4%		
Laying date	43	1968-2011	127	None			0 days		


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend




Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of adult birds surviving to following year - error bars represent 95% confidence limits

Causes of change

There is convincing evidence that nest failure rates rose during the principal period of population decline and this represents the most likely demographic mechanism driving the observed decreases in abundance. The most likely ecological driver of this pattern is habitat impoverishment due to agricultural intensification.

Change factor	Primary driver	Secondary driver
Demographic	Decreased breeding success	
Ecological	Agricultural intensification	

Further information on causes of change

Siriwardena et al. (1999, 2000b) provide convincing evidence that nest failure rates at the egg stage rose during the principal period of population decline and this represents the most likely demographic mechanism driving the observed decrease in abundance. They found an obvious change in the egg-stage failure rate of Linnnet nests after 1975 and this was detectable in the total fledglings produced, suggesting that the deterioration in breeding performance had an important role in driving the species' concurrent decline in abundance (Siriwardena et al. 2000b). Moorcroft & Wilson (2000) concur that the severe decline during the 1970s and 1980s occurred via a reduction in breeding success, attributing this to a reduction in the availability of breeding-season food supplies on arable farmland caused by agricultural intensification. However, they state that the precise demographic mechanism involved is unclear: instead of breeding performance per attempt, they suggest reductions in the number of nesting attempts being made by individual females or a reduction in immediate post-fledging survival due to resource limitations as more likely, although these hypotheses were not tested. BTO monitoring data do not permit analysis of these parameters but it is plausible that such effects occurred in parallel with the breeding success effects

indicated by NRS results. Nevertheless, all these patterns are consistent with the results of Siriwardena et al. (1999), who reported that index change was not significantly correlated with adult and first-year survival. They found no significant trend-specific difference in survival, and survival rates in periods of decline were higher than those in periods of increase.

After 1986, egg-stage nest survival increased and this led to a slight increase in breeding performance, although, as with the earlier decline, greater numbers of breeding attempts or increased post-fledging survival may also have contributed to the ending of population decline (Siriwardena et al. 2000b, Wilson et al. 1996, Moorcroft et al. 1997). Increases in the crop area of oilseed rape are thought to have improved Linnet breeding success by compensating for the herbicide-mediated decline in many farmland weeds that were traditionally important in this species' summer diet (Moorcroft et al. 1997). Both the number of breeding attempts possible in a season and post-fledging survival could have increased in response to this improvement in food supplies, as could chick survival. Oddly, Siriwardena et al. (2001b) identified a significant negative effect of rape on breeding performance through the egg-stage daily nest failure rate and no positive effect on success through the nestling stage in a further analysis of nest record data. This is clearly inconsistent with the results of intensive work on Linnets (Wilson et al. 1996, Moorcroft et al. 1997), perhaps reflecting the different geographical biases affecting nest records and this particular intensive study. Nevertheless, it suggests that environmental effects on Linnet breeding success show complex spatial variation and that the knock-on effects on trends in abundance could also be difficult to characterise.

The current long-term pattern, spanning the Linnet's periods of decrease and relative stability, is of linear increase in nest failure rates and linear decrease in the number of fledglings per breeding attempt (see above).

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Lesser Redpoll

Acanthis cabaret

Key facts


Conservation listings:	Europe (A. cabaret/flammea): no SPEC category (not concentrated in Europe, conservation status favourable) (BiE04) UK: red (>50% population decline) (BoCC3) UK Biodiversity Action Plan: priority species
Long-term trend:	England: rapid decline
Population size:	220,000 pairs in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)
Migrant status:	Short-distance migrant
Nesting habitat:	Above-ground nester
Primary breeding habitat:	Woodland
Secondary breeding habitat:	
Breeding diet:	Animal
Winter diet:	Vegetation

Status summary

Lesser Redpolls were abundant and widespread in lowland Britain in the 1970s, and frequent then on CBC and CES plots, but, concurrent with a sustained period of severe decline, have withdrawn completely as breeding birds from large areas of lowland England (Balmer et al. 2013). Uncertainty about the representativeness of the monitoring data prior to the establishment of BBS initially denied the species a place among birds of conservation concern, since it was thought possible that the population may have withdrawn from the lowlands to northern and western UK regions, where monitoring prior to 1994 was less effective. Since a range contraction of 11% between 1968-72 and 1988-91 was evident in all parts of the UK (Gibbons et al. 1993), however, it is perhaps more likely that decrease was general. Accordingly the species was moved from green to amber in 2002 and in 2009 to the red list. Since Acanthis cabaret is currently treated by BOU as a separate species from the Common Redpoll A. flammea, and has a restricted range that lies wholly within western Europe, it arguably warrants a European conservation listing at the next review. The taxonomic status of cabaret remains controversial, however (Stoddart 2013). Recent UK data show a general strong increase although, especially in lowland areas, the population remains severely depleted. A rapid increase has been recorded in the Republic of Ireland since 1998 (Crowe 2012).

CBC/BBS England 1966-2012

Lesser Redpoll







Smoothed population index, relative to an arbitrary 100 in the year given, with 85% confidence limits in green

Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	44	1967-2011	47	-83	-94	-53	>50	
	25	1986-2011	48	-88	-95	-78	>50	Small CBC sample
	10	2001-2011	70	41	-16	106		
	5	2006-2011	81	66	22	115		
BBS UK	16	1995-2011	159	55	26	96		
	10	2001-2011	179	31	5	57		
	5	2006-2011	207	47	29	72		
BBS England	16	1995-2011	62	14	-25	71		
	10	2001-2011	70	40	-23	106		
	5	2006-2011	81	66	23	126		
BBS Scotland	16	1995-2011	46	55	12	129		
	10	2001-2011	51	20	-8	63		
	5	2006-2011	62	73	37	111		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.





Demographic trends

Productivity and survival trends for this species are not currently produced by BTO

Causes of change

Although sample sizes are small, declines in both survival and productivity appear to have led to the Lesser Redpoll decline. Evidence for the ecological drivers behind this is largely circumstantial but they are thought to include maturation of woodland and a reduction in birch seed food supplies.

Change factor	Primary driver	Secondary driver
Demographic	Decreased survival	Decreased breeding success
Ecological	Changes in woodland	

Further information on causes of change

Though samples are too small to continue presenting a trend, CES data indicated a rapid long-term decline in productivity and there is evidence that survival rates have fallen (Siriwardena et al. 1998).

There is very little evidence available regarding the ecological drivers behind the decline of this species. In southern Britain, at least, the decrease may be attributable to a reduction in the amount of suitable young forest growth (Fuller et al. 2005). Amar et al. (2006) and Smart et al. (2007) both found relationships with lichen and bracken cover, although these studies were limited to broadleaved woodlands. Evans (1966) and Cramp & Perrins (1994) point to the importance of birch to the species, which could potentially explain the relationships found by Amar et al. (2006) and Smart et al. (2007). Birch seeds are an important component of this species' diet. Amaret al. (2006) state that birch has declined in many woodlands as they have matured, and this could raise the possibility of winter food as a factor in the species decline, although this evidence is circumstantial and given that species with similar winter diet, such as [Siskin](#), are faring better, may be unlikely.

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts


Conservation listings:	Europe: no SPEC category (not concentrated in Europe, conservation status favourable) (BiE04) UK: green (BoCC3)
Long-term trend:	UK: fluctuating, with no long-term trend
Population size:	40,000 (31,000-53,000) pairs in 2009 (APEP13: distance sampling estimate for 2006 (Newson et al. 2008) updated using BBS trend)

Status summary

The UK breeding population of Crossbills is difficult to assess in any one season, even by special survey, and is exceptionally variable between years. The core of the population lies in the taiga forests across Eurasia, from where birds periodically erupt westwards and southwards in search of better feeding conditions. After arrivals in Britain, many thousands of birds may stay to breed, perhaps for a few years, before survivors and their offspring return to the Continent (Newton 2006). The spur to movements is a failure of the cone crop, especially in Norway spruce *Picea abies*, which is this species' main food (Summers 1999). Crossbills begin breeding in January, sometimes even earlier, and by the start of the BBS period in April most sightings are of highly mobile family parties. In irruption years, BBS sightings may include many birds from the Continent, which often begin to arrive in late May or during June. The BBS trend therefore reflects post-breeding rather than breeding numbers, and on a wider geographical scale than just the UK. Atlas data for 2008-11 confirm that Crossbills are currently at a high level of abundance (Balmer et al. 2013) but it is not clear whether recent increase is part of any long-term trend.

BBS UK 1994–2012

Crossbill




Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	16	1995-2011	56	130	46	231		
	10	2001-2011	68	181	97	279		

Source	5 Period (yrs)	2006-2011 Years	83 Plots (n)	131 Change (%)	62 Lower limit	218 Upper limit	Alert	Comment
--------	----------------------	--------------------	--------------------	----------------------	----------------------	-----------------------	-------	---------

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.



Demographic trends

Productivity and survival trends for this species are not currently produced by BTO

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>


Key facts

Conservation listings:	Europe: no SPEC category (not concentrated in Europe, conservation status favourable) (BiE04) UK: green (species level); amber (race britannica, >20% of European breeders) (BoCC3)
Long-term trend:	England: moderate increase
Population size:	1.2 (1.1-1.3) million pairs in 2009 (APEP13: distance sampling estimate for 2006 (Newson et al. 2008) updated using BBS trend)

Status summary

Goldfinch abundance fell sharply from the mid 1970s until the mid 1980s, but the decline was both preceded and followed by significant population increases. The recent upturn has lifted the species from the amber list of conservation concern into the green category, and has been accompanied by an increase in its use of gardens for winter feeding. The BBS Siriwardena et al. 1999). There have been no clear changes in productivity as measured by NRS and CES. A strong trend towards earlier laying has been identified and may be partly explained by recent climate change (Crick & Sparks 1999). There has been widespread moderate increase across Europe since 1980, though with little change since 1990 (PECBMS 2013a). A strong increase has been recorded in the Republic of Ireland since 1998 (Crowe 2012).

CBC/BBS England 1966-2012 Goldfinch



Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	44	1967-2011	550	91	41	139		
	25	1986-2011	899	244	183	297		
	10	2001-2011	1538	83	75	92		
	5	2006-2011	1803	58	53	64		

CES adults Source	27 Period (yrs)	1984-2011 Years 1986-2011	31 Plots 33	69 Change (%)	-7 Lower limit	197 Upper limit	Alert	Comment
	10	2001-2011	39	52	14	98		
	5	2006-2011	41	46	15	87		
CES juveniles	27	1984-2011	22	-1	-61	406		
	25	1986-2011	23	119	16	409		
	10	2001-2011	26	82	16	178		
	5	2006-2011	27	136	43	261		
BBS UK	16	1995-2011	1553	109	97	122		
	10	2001-2011	1842	80	71	88		
	5	2006-2011	2119	52	46	60		
BBS England	16	1995-2011	1282	100	88	116		
	10	2001-2011	1518	84	75	93		
	5	2006-2011	1770	57	53	63		
BBS Scotland	16	1995-2011	88	164	84	237		
	10	2001-2011	102	95	48	141		
	5	2006-2011	116	54	20	96		
BBS Wales	16	1995-2011	126	79	45	130		
	10	2001-2011	146	12	-4	30		
	5	2006-2011	148	16	-2	32		
BBS N.Ireland	16	1995-2011	47	929				
	10	2001-2011	64	197				
	5	2006-2011	73	64				

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.




CBC/BBS England 1966-2012
Goldfinch




CBC/BBS England graph

CES adult abundance 1983-2012
Goldfinch








CES adults graph

CES juvenile abundance 1983-2012
Goldfinch




CES juveniles graph



Fledglings per breeding attempt


Goldfinch 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Goldfinch




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	29	None					
Clutch size	43	1968-2011	21	None					Small sample
Brood size	43	1968-2011	35	None					
Nest failure rate at egg stage	43	1968-2011	37	None					
Nest failure rate at chick stage	43	1968-2011	30	None					Small sample
Laying date	43	1968-2011	24	Curvilinear	Jun 5	May 21	-15 days		Small sample
Juvenile to Adult ratio (CES)	27	1984-2011	36	Smoothed trend	315 Index value	100 Index value	-68%	>50	
Juvenile to Adult ratio (CES)	25	1986-2011	38	Smoothed trend	591 Index value	100 Index value	-83%	>50	
Juvenile to Adult ratio (CES)	10	2001-2011	45	Smoothed trend	196 Index value	100 Index value	-49%	>25	
Juvenile to Adult ratio (CES)	5	2006-2011	45	Smoothed trend	85 Index value	100 Index value	17%		


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend




Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Smoothed long-term trend in ratio of juvenile:adult birds caught - green lines indicate 85% confidence limits

Key facts


Conservation listings:	Europe: no SPEC category (concentrated in Europe, conservation status favourable) (BiE04) UK: green (BoCC3)
Long-term trend:	UK: increase
Population size:	420,000 pairs in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)

Status summary

The maturing of new conifer plantations has aided the spread of breeding Siskins throughout the UK, from their previous stronghold in the Scottish Highlands, since about 1950. Their habit of using garden feeders, especially in late winter, has developed since the 1960s and, despite many of the birds involved migrating to the Baltic region to breed, may also have helped to boost the UK breeding population. The 1988-91 Breeding Atlas identified a major expansion of the breeding range into southern Britain (Gibbons et al. 1993) and subsequently there have been considerable further range gains, especially in the south and west (Balmeret al. 2013). Progressively more CBC plots became occupied during the 1970s and 1980s (Marchant et al. 1990), but samples were still insufficient for annual monitoring until BBS began in 1994. Results since then show parallel fluctuations of extraordinary amplitude in England and Scotland. To some extent, this may reflect the occasional large continental influxes affecting spring numbers on a broad UK scale. The overall trend across Europe since 1980 has been a moderate decrease (PECBMS 2013a).

BBS UK 1994–2012

Siskin




Population changes in detail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	16	1995-2011	165	89	42	129		
	10	2001-2011	193	109	73	152		
	5	2006-2011	246	93	64	123		

BBS England Source	16 Period (yrs)	1995-2011 Years 2001-2011	58 Plots (72)	96 Change (74)	5 Lower Limit	350 Upper Limit	Alert	Comment
	5	2006-2011	98	96	37	195		
BBS Scotland	16	1995-2011	73	81	28	129		
	10	2001-2011	82	90	45	148		
	5	2006-2011	104	88	48	126		


Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.



BBS UK graph



BBS England graph



BBS Scotland graph

Demographic trends

Productivity and survival trends for this species are not currently produced by BTO

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Key facts


Conservation listings:	Europe: no SPEC category (concentrated in Europe, conservation status favourable) (BiE04) UK: red (>50% population decline) (BoCC3) UK Biodiversity Action Plan: priority species
Long-term trend:	UK, England: rapid decline
Population size:	710,000 territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)
Migrant status:	Resident
Nesting habitat:	Ground nester
Primary breeding habitat:	Farmland
Secondary breeding habitat:	
Breeding diet:	Animal
Winter diet:	Vegetation

Status summary

Yellowhammer abundance began to decline on farmland in the mid 1980s. The downward trend has continued to at least 2009, although with substantial increase in Scotland since 2003. The BBS Balmer et al. 2013). The species, listed as green in 1996, has been red listed since 2002. Numbers have shown widespread moderate decrease across Europe since 1980 (PECBMS 2013a).

CBC/BBS UK 1966–2012




Yellowhammer




Smoothed population index, relative to an arbitrary 100 in the year given, with 85% confidence limits in green


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	550	-55	-65	-44	>50	
	25	1986-2011	855	-50	-56	-43	>25	
	10	2001-2011	1263	-4	-10	2		
	5	2006-2011	1393	1	-5	6		
CBC/BBS England	44	1967-2011	480	-59	-68	-49	>50	
	25	1986-2011	747	-56	-61	-51	>50	
	10	2001-2011	1102	-13	-17	-9		
	5	2006-2011	1212	-4	-7	-1		
BBS UK	16	1995-2011	1177	-13	-19	-8		
	10	2001-2011	1251	-4	-9	2		
	5	2006-2011	1373	1	-3	6		
BBS England	16	1995-2011	1027	-22	-26	-18		
	10	2001-2011	1096	-13	-18	-9		
	5	2006-2011	1202	-4	-8	-1		
BBS Scotland	16	1995-2011	105	31	7	55		
	10	2001-2011	112	39	23	60		
	5	2006-2011	127	24	11	41		
BBS Wales	16	1995-2011	35	-48	-64	-33	>25	
	10	2001-2011	34	-27	-42	-11	>25	
	5	2006-2011	32	-20	-32	-1		

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.






BBS England graph




BBS Scotland graph



BBS Wales graph

Demographic trends


Fledglings per breeding attempt Yellowhammer 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012


Yellowhammer




More on demographic trends


Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	49	Curvilinear	0.82 fledglings	1.16 fledglings	41.3%		
Clutch size	43	1968-2011	43	None					
Brood size	43	1968-2011	66	Curvilinear	2.98 chicks	3.00 chicks	0.7%		
Nest failure rate at egg stage	43	1968-2011	63	Curvilinear	4.96% nests/day	3.04% nests/day	-38.7%		
Nest failure rate at chick stage	43	1968-2011	50	Curvilinear	4.34% nests/day	3.83% nests/day	-11.8%		
Laying date	43	1968-2011	26	Linear increase	May 31	Jun 7	7 days		Small sample

For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).






Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

Causes of change

Declines in annual survival have been proposed as the demographic mechanism for decline, due to winter resource limitation, although ring-recovery data are sparse and so most evidence for this is indirect.

Change factor	Primary driver	Secondary driver
Demographic	Decreased survival	
Ecological	Agricultural intensification	

Further information on causes of change

Yellowhammer is unique among farmland birds in that its population was stable until the mid 1980s, followed by a decline, suggesting that it alone was affected by some change that occurred in the 1980s (Siriwardena et al. 1998a). Although long-term demographic trends presented here show no linear changes over time (see above), there is some evidence that survival rates decreased during the initial period of decline (Siriwardena et al. 1998b, 2000a, Kyrkos 1997), and that breeding performance tended to improve (Siriwardena et al. 2000b). However, declines in clutch size, brood size and nest success were formerly of NRS concern (Leech & Barimore 2008).

Best estimates of the variation in adult and first-year Yellowhammer survival (from ring recoveries) suggest that it has been sufficient to explain the species' decline (Kyrkos 1997). Reductions in winter seed availability as a result of agricultural intensification (for example, the loss of winter stubbles and a reduction in weed densities) are widely believed to have contributed to the population decline, presumably through impacts on survival rates. Siriwardena et al. (2007), found that Yellowhammer declines were less steep in areas where the species received more overwinter provisioning, providing experimental evidence for winter resource limitation. Food availability (and therefore, as a conservation measure, supplementary feeding) in late winter appears to be particularly important because demand for seed food is greatest at this time and this is also when the food supply resulting from agri-environment conservation measures is at its lowest (Siriwardena et al. 2007). Further evidence comes from Gillings et al. (2005), who used two complementary extensive bird surveys undertaken at the same localities in summer and winter to show that the areas of extensive stubble in winter were correlated with better population performance, presumably because overwinter survival is relatively high. This is supported by another study, in Oxfordshire (Wilson et al. 1996), which found that the only habitat type for which a clear preference was displayed in winter was stubble.

In terms of changes to habitat, Kyrkos et al. (1998) found that Yellowhammer breeding density decreased with increasing proportion of farmland under grassland. It may be that modern improved grassland has neither the weed density required by adult Yellowhammers nor sufficient invertebrate prey for birds feeding nestlings. The dense sward structure of highly fertilised leys may also reduce access to invertebrate prey (Perkins et al. 2000). This is supported by the results of Douglas et al. (2010a) who found that foraging in grass margins was increased by experimental mowing, showing that access to prey in dense vegetation limits feeding activity. Siriwardena et al. (2000b, 2000c) provide further evidence that grazing supported the lowest breeding performance, although the best breeding performance was associated with mixed farmland, suggesting that loss of heterogeneity in the landscape may be a factor in the decline, although they state that this is unlikely to be the main mechanism behind

the declines. Bradbury & Stoate (2000) further suggest that loss or degradation of hedges and field margins, loss of stubbles and intensification of grassland management may have reduced nest-site and food availability for farmland Yellowhammers.

Increased use of pesticides may have also played a role in decreasing breeding success. Boatman et al. (2004) used an experimental set-up to look at the effect of pesticides on breeding performance, and further evidence was provided by Morris et al. (2005), who showed that increased use of pesticides results in reduced invertebrate abundance, lower brood production and fewer chicks fledging. Hart et al. (2006) also demonstrated how insecticide applications can depress Yellowhammer breeding productivity. Whittingham et al. (2005) found that the local availability of rotational set-aside was a good predictor of sites chosen for breeding territories, which could reflect the benefits of both sparse vegetation (access to bare ground for foraging) and lack of pesticide use.

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>


Key facts

Conservation listings:	Europe: no SPEC category (not concentrated in Europe, conservation status favourable) (BiE04) UK: amber (25-50% population decline to 2006) (BoCC3) UK Biodiversity Action Plan: click here , priority species
Long-term trend:	UK: moderate decline England: shallow decline
Population size:	250,000 territories in 2009 (APEP13: 1988-91 Atlas estimate updated using CBC/BBS trend)

Status summary

Both CBC/BBS and WBS/WBBS indices declined rapidly during the 1970s, after an earlier increase, but Reed Bunting abundance has fluctuated without a clear trend since the 1980s. Since 1994, results from BBS indicate significant population increase, though with a downturn in the last few seasons. The BBS Peach et al. 1999). This is supported by a moderate decline in CES productivity and by a major increase in failure rates at the egg stage, which has raised NRS concern (Leech & Barimore 2008). Farmland densities are four times higher in oilseed rape than in cereals or setaside and this crop is crucial in reducing the dependency of the species on wetlands (Gruar et al. 2006). The initial decline placed Reed Bunting on the red list but in 2009, with evidence from BBS of some recovery in numbers, the species was moved from red to amber. There has been widespread moderate decrease across Europe since 1980 (PECBMS 2013a).

CBC/BBS UK 1966-2012
Reed Bunting






Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	248	-27	-42	-10	>25	
	25	1986-2011	359	10	-9	32		
	10	2001-2011	575	14	4	28		


Source	Period (yrs)	2006-2011 Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	44	1967-2011	194	-22	-40	-4		
	25	1986-2011	274	21	-3	50		
	10	2001-2011	431	30	19	41		
	5	2006-2011	504	3	-3	9		
WBS/WBBS waterways	36	1975-2011	85	-59	-71	-41	>50	
	25	1986-2011	101	-3	-25	30		
	10	2001-2011	143	11	1	25		
	5	2006-2011	135	-2	-11	11		
CES adults	27	1984-2011	59	-62	-72	-48	>50	
	25	1986-2011	62	-59	-70	-44	>50	
	10	2001-2011	63	-22	-35	-5		
	5	2006-2011	65	-25	-35	-14	>25	
CES juveniles	27	1984-2011	44	90	-17	311		
	25	1986-2011	47	-5	-42	44		
	10	2001-2011	47	-13	-34	20		
	5	2006-2011	47	17	-9	50		
BBS UK	16	1995-2011	476	19	9	33		
	10	2001-2011	555	16	6	29		
	5	2006-2011	631	-7	-13	-1		
BBS England	16	1995-2011	358	26	12	43		
	10	2001-2011	416	30	18	40		
	5	2006-2011	479	2	-5	9		
BBS Scotland	16	1995-2011	58	19	-10	59		
	10	2001-2011	68	0	-19	34		
	5	2006-2011	79	-21	-35	-5		
BBS N.Ireland	16	1995-2011	32	-17	-44	35		
	10	2001-2011	38	-16	-39	26		
	5	2006-2011	39	-28	-42	-12	>25	

Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.







WBS/WBBS waterways graph




CES adults graph




CES juveniles graph




BBS UK graph



BBS England graph




BBS Scotland graph



Demographic trends


Fledglings per breeding attempt Reed Bunting 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012


Reed Bunting




More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	47	None					
Clutch size	43	1968-2011	44	None					
Brood size	43	1968-2011	61	None					
Nest failure rate at egg stage	43	1968-2011	51	Curvilinear	0.63% nests/day	1.19% nests/day	88.9%		
Nest failure rate at chick stage	43	1968-2011	51	None					
Laying date	43	1968-2011	48	None			0 days		
Juvenile to Adult ratio (CES)	27	1984-2011	62	Smoothed trend	227 Index value	100 Index value	-56%	>50	
Juvenile to Adult ratio (CES)	25	1986-2011	65	Smoothed trend	310 Index value	100 Index value	-68%	>50	
Juvenile to Adult ratio (CES)	10	2001-2011	67	Smoothed trend	154 Index value	100 Index value	-35%	>25	
Juvenile to Adult ratio (CES)	5	2006-2011	69	Smoothed trend	101 Index value	100 Index value	-1%		


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend




Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend




Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Smoothed long-term trend in ratio of juvenile:adult birds caught - green lines indicate 85% confidence limits



Proportion of adult birds surviving to following year - green bars represent 95% confidence limits

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>

Corn Bunting

Emberiza calandra

Key facts


Conservation listings:	Europe: SPEC category 2 (declining) (BiE04) UK: red (>50% population decline, historical decline) (BoCC3) UK Biodiversity Action Plan: click here , priority species
Long-term trend:	UK, England: rapid decline
Population size:	11,000 (9,000-13,000) territories in 2009 (APEP13: 1993 estimate (Donald & Evans 1995) updated using CBC/BBS trend)
Migrant status:	Resident
Nesting habitat:	Ground nester
Primary breeding habitat:	Farmland
Secondary breeding habitat:	
Breeding diet:	Animal
Winter diet:	Vegetation

Status summary

Following an earlier, historical decrease, Corn Buntings declined very steeply between the mid 1970s and mid 1980s, with local extinctions across large sections of their former range. Subsequently the decline has continued, but at a reduced rate. Corn Buntings have declined rapidly across Europe since 1980, though with no detectable change since 1990 (PECBMS 2013a), and have declined to extinction in Ireland (Taylor & O'Halloran 2002). With declines across much of its European range, this previously 'secure' species is now provisionally evaluated as 'declining' (BirdLife International 2004).

CBC/BBS UK 1966–2012

Corn Bunting







Smoothed population index, relative to an arbitrary 100 in the year given, with 85% confidence limits in green

Population changes in detail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	44	1967-2011	73	-87	-94	-77	>50	
	25	1986-2011	106	-65	-82	-46	>50	Small CBC sample
	10	2001-2011	139	-9	-25	7		
	5	2006-2011	152	-8	-21	7		
CBC/BBS England	44	1967-2011	70	-85	-94	-74	>50	
	25	1986-2011	101	-63	-81	-44	>50	Small CBC sample
	10	2001-2011	133	-2	-18	17		
	5	2006-2011	146	-8	-20	7		
BBS UK	16	1995-2011	142	-34	-45	-23	>25	
	10	2001-2011	138	-9	-27	11		
	5	2006-2011	151	-5	-18	9		
BBS England	16	1995-2011	136	-30	-43	-19	>25	
	10	2001-2011	133	-3	-23	17		
	5	2006-2011	145	-5	-18	6		


Tables show changes with their 90% confidence limits. Alerts are flagged for significant changes only. See [here](#) for more information.





Demographic trends


Fledglings per breeding attempt Corn Bunting 1966–2012



Mean number of fledglings produced per nest - green bars represent standard error and black line shows long-term trend

Laying date 1966–2012

Corn Bunting




Mean laying date in Julian days (1st April = Day 90) - green bars represent standard error and black line shows long-term trend


More on demographic trends

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2011	Change	Alert	Comment
Fledglings per breeding attempt	43	1968-2011	10	None					
Brood size	43	1968-2011	15	Curvilinear	3.32 chicks	3.59 chicks	8.3%		Small sample
Nest failure rate at egg stage	43	1968-2011	12	None					Small sample
Nest failure rate at chick stage	43	1968-2011	15	Curvilinear	4.50% nests/day	2.48% nests/day	-44.9%		Small sample
Laying date	43	1968-2011	17	Linear decline	Jun 27	Jun 13	-14 days		Small sample


For details of analytical methods for the Nest Record Scheme, the Constant Effort Sites (CES) and the Retrapping Adults for Survival (RAS) scheme, please follow links [here](#).




Mean number of eggs per nest - green bars represent standard error and black line shows long-term trend



Mean number of chicks per nest - green bars represent standard error and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend



Proportion of nests failing per day during incubation - green bars represent 95% confidence limits and black line shows long-term trend

Causes of change

Changes in farming practice are believed to have been responsible for declines, through impacts on reduced seed and/or invertebrate abundance. The demographic causes are unclear and there is conflicting evidence as to whether breeding or wintering effects have been the primary driver.

Change factor	Primary driver	Secondary driver
Demographic	Unknown	
Ecological	Agricultural intensification	

Further information on causes of change

National-scale evidence gives no indication of a historical role for breeding success, but there are contemporary local correlations between agricultural practices and breeding success, including a notable effect on numbers of breeding attempts. Causes of change may be different in different populations, as some of this species' breeding habitats are completely different and isolated from each other. There is no way to test for effects of survival. Conversely, it is easy to test for effects on breeding success, especially locally and with respect to contemporary as opposed to historical land-use. This leads to a big imbalance in the evidence available.

Breeding performance per nesting attempt increased considerably while population numbers have been declining (Crick 1997, Siriwardena et al. 2000a), but it is also reported that fewer birds now raise a second brood, thus reducing productivity overall (Brickle & Harper 2002). Brood size has decreased since 1990 (see graph above). Ring-recovery sample sizes do not permit an analysis of survival rates, meaning that it is impossible to test for effects of survival (Siriwardena et al. 1998a, 2000a). Any decrease there has been in survival rates is probably a result of the reduction in winter seed availability that has followed from agricultural intensification (Donald 1997, Wilson et al. 2007). Donald & Evans (1994) found that 60% of Corn Buntings fed on winter stubbles, which were the only field type for which a consistent preference was detected.

Spring-sown cereals have been found to be a particularly important habitat for Corn Bunting (Brickle & Harper 2000, Fox & Heldbjerg 2008), and hence its reduction may have contributed to declines, as they provide long-lasting stubbles during the winter and abundant food in the form of surface grain when first sown. In the breeding season, spring cereals were among the most frequently used habitats for nesting and for collecting chick food; territory associations with overhead wires (for songposts) and fallow (positive in early summer, negative in late summer) became stronger in later years as the population declined (Perkins et al. 2012). Siriwardena et al. (2000b) provide evidence that mixed farming at the territory scale supported better breeding performance. However, Donald & Forrest (1995) found little evidence for breeding-season effects in their study using CBC data and suggest that numbers are more likely to have declined due to reduced winter food supplies resulting particularly from the loss of spring tillage, increased pesticide usage and improved harvesting and storage techniques.

A reduction in food availability has been implicated in the declines of this species. In arable-dominated areas in Scotland, Perkins et al. (2011) provide evidence showing that AES management (agri-environment schemes) that increased food availability reversed population declines. However, where a high proportion of Corn Buntings

nested in grasslands, an additional AES option that delayed mowing was essential to achieving population increase. Setchfield et al. (2012) have further demonstrated that AES management of cereals can boost productivity and emphasise the importance of delayed harvest to the number and success of late nests.

As part of a PhD study, Brickle (1999) modelled the population dynamics of Corn Buntings in Sussex, concluding that productivity was the most likely cause of decline in the South Downs, also finding evidence of indirect effects of pesticides. Brickle & Harper (1999) identified the main food items of chicks, most of which have declined in abundance on lowland farmland (Campbell et al. 1997). Boatman et al. (2004) further analysed the data from Brickle et al. (2000) and found that arthropod abundance in the vicinity of the nest had a significant effect on the survival of broods, although this was based only on two years' data, whilst Ewald et al. (2002) found that densities of Corn Bunting were higher where the number of pesticide applications was low. Brickle et al. (2000) found that chick weight and nest survival at the nestling stage were respectively positively and negatively correlated with invertebrate food availability, and chick food abundance was negatively correlated with the number of insecticide applications to cereal fields. However, the authors state that the contribution of this reduction in breeding performance to the Corn Bunting's decline depends on the mortality rates for fledged chicks and older birds, information on which is sparse.

Brickle & Harper (2002) found that, although predation accounted for the majority of nest failures in their Corn Bunting study population, there was a seasonal decline in the nest survival rate during incubation, which was largely due to increased losses through farming operations. Furthermore, they speculated that harvesting of cereal crops may reduce the availability of suitable breeding habitat late in the season, thus curtailing the length of the breeding season, and preventing double-brooding. A reduction in fecundity via these mechanisms provides one explanation for the collapse of the Corn Bunting population (Donald 1997, Brickle & Harper 2002).

This report should be cited as: Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds. BTO Research Report No. 652. BTO, Thetford. <https://www.bto.org/birdtrends>



Images: Grey Wagtail, by Edmund Fellowes / BTO; Lapwing, by Sarah Kelman / BTO

BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds.

This report is a “one-stop-shop” for information about the population status of our common terrestrial birds. With one page per species, readers can quickly find all the key information about trends in population size and breeding performance as measured by BTO monitoring schemes. It provides an overview of trends for the period 1966-2013.

This report is the third in a series, prepared within the Partnership between the British Trust for Ornithology (BTO) and the Joint Nature Conservation Committee (JNCC) (on behalf of Natural England, Scottish Natural Heritage, Countryside Council for Wales and the Environment & Heritage Service of Northern Ireland) as part of its programme of research into nature conservation.

It is the result of the sustained long-term fieldwork efforts of many thousands of the BTO's volunteer supporters. Without their enthusiasm for collecting these hard-won facts, the cause of conservation in the UK would be very much the poorer.

Baillie, S.R., Marchant, J.H., Leech, D.I., Massimino, D., Eglington, S.M., Johnston, A., Noble, D.G., Barimore, C., Kew, A.J., Downie, I.S., Risely, K. & Robinson, R.A. (2014). BirdTrends 2013: trends in numbers, breeding success and survival for UK breeding birds.. *BTO Research Report 652*, BTO, Thetford, UK.