

Brood patch codes

Code State of brood patch

- 0 absent
- 1 starting
- well defined, breast muscle and gut still visible through the skin
- 3 the skin of the belly is opaque, thickened or engorged, veined and red with broad, undulating wrinkles
- 4 skin of belly has thin wrinkles, engorgement of skin has gone, but the skin is still stretched, and the breast muscle and gut are visible again through the skin
- 5 feathering over
- P present

Moult codes

Code Stage of moult

- J = wholly **J**uvenile plumage
- P = undergoing Post-juvenile body moult (see codes M, A if in full post-juvenile moult)
- B = undergoing active Body moult not a post-juvenile moult and not part of a full moult of wings and tail
- Old plumage, no body or main moult (includes birds that have completed partial postiuvenile moult)
- S = Starting body moult: main moult yet to start wings and tail old
- **E** = **E**nding body moult: main moult finished wings and tail new
- N = New plumage following main moult, no body or main moult (used for birds that have recently completed moult, not used after 31 December)
- M^1 = active Moult of wings and tail (main moult)
- $\underline{\mathbf{A}} = \mathbf{A}$ rrested wing moult (primaries or secondaries)
- T = Partial post-juvenile moult involving Tail or Tertials

¹If a moult code of M is entered, then computer variables ACT1\$ or ACT2\$ should be set to M if these are not already in use for other higher priority activity codes.

Primary moult scores

Score Stage of growth

- 0 an old feather
- 1 old feather missing or new feather completely in pin
- new feather just emerging from the sheath, up to one third grown
- 3 new feather between one and two thirds grown
- 4 new feather more than two thirds grown, but waxy sheath still at its base
- 5 new feather fully grown with no trace of sheath at its base

Post-juvenile moult in the alula



Code Moult stage

- 0 both old
- 1 largest old, other new
- 2 largest new, other old
- both new

Sexing method codes

Code Description

- $\mathbf{D} = \mathbf{D}\mathbf{N}\mathbf{A}$
- E = internal cloacal Examination when used to confirm sexing eg in wildfowl
- C = Cloacal protuberance (males only)
- $\mathbf{B} = \mathbf{B}$ rood patch present (females only)
- **A** = **A**ctivity bird sexed by observing activity or behaviour
- S = Size difference ie a measurement such as wing length is the only criterion of sexing

Note: Use only the codes given above and no others.

Activity Codes

Code Activity

- A = Artificial bait provided intentionally or unintentionally (including rubbish tips, agricultural products eg grain and carrion)¹
- T = Tape-lured
- $L = \text{at } \hat{\mathbf{L}}$ ighthouse as a result of night-time attraction
- = at Sea on board ship or other man-made structure
- N = Nesting adult bird currently known to be breeding
- C = at Colony (not necessarily breeding) record for full-grown only, **not** for pulli
- $\mathbf{M} = \text{undergoing } \mathbf{M} \text{oult of wing or tail feathers (normal moult, not just odd feathers)}$
- $\mathbf{R} = \text{at or joining } \mathbf{R} \text{oost}$
- P = on Passage species which is not regular as resident, winter or summer, certainly not a local breeding or wintering bird

Note: Code A is not available in B-RING. Use only the codes given above and no others